arXiv:math/0106157v1 [math.SG] 19 Jun 2001

Gromov–Witten invariants of symplectic quotients and adiabatic limits

A. Rita Gaio Dietmar A. Salamon
Fac. Ciências-Porto ETH-Zürich

19 June 2001

Abstract

We study pseudoholomorphic curves in symplectic quotients as adiabatic limits of solutions of a system of nonlinear first order elliptic partial differential equations in the ambient symplectic manifold. The symplectic manifold carries a Hamiltonian group action. The equations involve the Cauchy-Riemann operator over a Riemann surface, twisted by a connection, and couple the curvature of the connection with the moment map. Our main theorem asserts that the genus zero invariants of Hamiltonian group actions defined by these equations are related to the genus zero Gromov–Witten invariants of the symplectic quotient (in the monotone case) via a natural ring homomorphism from the equivariant cohomology of the ambient space to the quantum cohomology of the quotient.

1 Introduction

The main theorem of this paper asserts that under certain hypotheses there is a ring homomorphism from the equivariant cohomology of a symplectic manifold M with a Hamiltonian G-action to the quantum cohomology of the symplectic quotient $\bar{M}$ such that the following diagram commutes

$$\begin{tikzcd}
\mathrm{H}^*(M_{\mathrm{G}}) \arrow[rr, "\varphi"] \arrow[dr, "\Phi_B"'] & & \mathrm{QH}^*(\bar{M}) \arrow[dl, "\mathrm{GW}_{\bar{B}}"] \\
& \mathbb{Z} &
\end{tikzcd}$$

Here $\mathrm{G}W_{\bar{B}}$ denotes the genus zero Gromov–Witten invariants of $\bar{M}$ with fixed marked points associated to a homology class $\bar{B} \in H_2(\bar{M};\mathbb{Z})$, and Φ_B denotes the genus zero invariants of Hamiltonian group actions associated to the equivariant homology class $B = \kappa(\bar{B}) \in H_2(M_{\mathrm{G}};\mathbb{Z})$. The latter invariant was introduced in [CGS, CGMS, M]. The homomorphism φ is defined indirectly as a consequence of a comparison theorem for the two invariants. A more direct definition in terms of vortices over the complex plane with values in M will be given elsewhere. The proof of the comparison theorem is based on an adiabatic limit analysis which relates the solutions of the equations used in the definition of the invariants Φ to pseudoholomorphic curves in the symplectic quotient. Our hypotheses are that the moment map is proper, that M is convex at infinity, and that the quotient $\bar{M}$ is smooth. These hypotheses are needed to even state the result. In addition we assume that there are no holomorphic spheres in the ambient manifold (and hence M is necessarily noncompact) and that $\bar{M}$ is monotone. These hypotheses are of technical nature and it might be possible to remove them. But this would require more analysis than is carried out in the present paper. Before stating the main results more precisely (Theorem A and Corollary A') we begin with a brief discussion of the invariants introduced in [CGS, CGMS, M].

Invariants of Hamiltonian group actions

Let (M,ω) be a symplectic manifold (not necessarily compact) and G be a compact connected Lie group with Lie algebra $\mathfrak{g}$. We fix an invariant inner product $\langle\cdot,\cdot\rangle$ on $\mathfrak{g}$ and identify $\mathfrak{g}$ with its dual $\mathfrak{g}^*$. We assume that G acts on M by Hamiltonian symplectomorphisms and that the action is generated by an equivariant moment map $\mu : M \to \mathfrak{g}$. This means that, for every $\eta \in \mathfrak{g}$, the vector field $X_\eta \in \mathrm{Vect}(M)$ that generates the action is determined by $\iota(X_\eta)\omega = d\langle\mu,\eta\rangle$. Let $\pi : P \to \Sigma$ be a principal G-bundle over a compact oriented Riemann surface $(\Sigma, j_\Sigma, \mathrm{dvol}_\Sigma)$. We fix a smooth family $\Sigma \to \mathcal{J}_{\mathrm{G}}(M,\omega) : z \mapsto J_z$ of G-invariant and ω-compatible almost complex structures on M. This determines a family of metrics $\langle\cdot,\cdot\rangle_z := \omega(\cdot, J_z\cdot)$. The invariants are derived from the equations

$$\bar{\partial}_{J,A}(u) = 0, \qquad *F_A + \mu(u) = 0, \tag{1}$$

for a pair (u,A), where $u : P \to M$ is a G-equivariant map and A is a connection on P. Here $\bar{\partial}_{J,A}$ is the nonlinear Cauchy-Riemann operator twisted by A and F_A is the curvature of A. Both terms in the second identity in (1) are sections of the Lie algebra bundle $\mathfrak{g}_P := P \times_{\mathrm{Ad}} \mathfrak{g}$. In contrast to the theory of pseudoholomorphic curves, equations (1) involve the volume form dvol_Σ (via the Hodge $*$-operator in the second equation) and not just the complex structure j_Σ. Equations (1) are invariant under the action of the gauge group $\mathcal{G} = \mathcal{G}(P)$ (of equivariant maps from P to G) by

$$g^*(u, A) = (g^{-1}u, g^{-1}dg + g^{-1}Ag).$$

¿From a geometric point of view, the solutions of (1) correspond to the absolute minima of the energy functional

$$E(u, A) = \frac{1}{2}\int_\Sigma \left(|d_Au|^2 + |F_A|^2 + |\mu(u)|^2\right) \mathrm{dvol}_\Sigma$$

in a given homotopy class. If the pair (u, A) is a solution of (1) then it has energy

$$E(u, A) = \int_\Sigma \left(u^*\omega - d\langle\mu(u), A\rangle\right) =: \langle[\omega - \mu], [u]\rangle$$

and this number is an invariant of the equivariant homology class represented by the map u. We impose the following hypothesis throughout this paper.

(H1) *The moment map μ is proper, zero is a regular value of μ, and* G *acts freely on $\mu^{-1}(0)$.*

Under this hypothesis the quotient

$$\bar{M} := M/\!\!/\mathrm{G} := \mu^{-1}(0)/\mathrm{G}$$

is a compact symplectic manifold. The induced symplectic form will be denoted by $\bar{\omega}$. The equivariant homology class $[u] \in H_2(M_{\mathrm{G}}; \mathbb{Z})$ is defined by the following diagram, which also shows how it is related to the characteristic class $[P] \in H_2(\mathrm{BG}; \mathbb{Z})$ and to the class $[\bar{u}] \in H_2(\bar{M}; \mathbb{Z})$ in the case $\mu \circ u \equiv 0$. Note that, since G is connected, the equivariant homology class $[u]$ determines the isomorphism class of the bundle P. We denote $M_{\mathrm{G}} := M \times_{\mathrm{G}} \mathrm{EG}$.

$$\begin{tikzcd}
{[\Sigma]} & \in & \mathrm{H}_2(\Sigma;\mathbb{Z}) \arrow[r, "\bar{u}_*"] & \mathrm{H}_2(\bar{M};\mathbb{Z}) \arrow[d, "\kappa"] & \ni & {[\bar{u}]} \\
 & & \mathrm{H}_2(P_{\mathrm{G}};\mathbb{Z}) \arrow[u, "\simeq"] \arrow[r, "u_*"] \arrow[d] & \mathrm{H}_2(M_{\mathrm{G}};\mathbb{Z}) \arrow[dl] & \ni & {[u]} \\
{[P]} & \in & \mathrm{H}_2(\mathrm{BG};\mathbb{Z}) & & &
\end{tikzcd}$$

Fix a homology class $\bar{B} \in H_2(\bar{M};\mathbb{Z})$, let $B := \kappa(\bar{B}) \in H_2(M_{\mathrm{G}};\mathbb{Z})$, and denote the space of solutions of (1) that represent this homology class by

$$\widetilde{\mathcal{M}}_{B,\Sigma} := \{(u,A) \in C^\infty_{\mathrm{G}}(P,M) \times \mathcal{A}(P) \,|\, [u] = B,\ u \text{ and } A \text{ satisfy (1)}\}\,.$$

Here $P \to \Sigma$ denotes a principal G-bundle whose characteristic class $[P] \in H_2(\mathrm{BG};\mathbb{Z})$ is determined by B as above. The quotient by the action of the gauge group will be denoted by

$$\mathcal{M}_{B,\Sigma} := \widetilde{\mathcal{M}}_{B,\Sigma}/\mathcal{G}.$$

We impose another hypothesis which guarantees compactness [CGMS].

(H2) *There exists a* G*-invariant almost complex structure* $J \in \mathcal{J}_{\mathrm{G}}(M,\omega)$*, a proper* G*-invariant function* $f : M \to [0,\infty)$*, and a constant* $c > 0$ *such that*

$$f(x) \geq c \qquad \Longrightarrow \qquad \langle \nabla_\xi \nabla f(x), \xi \rangle > 0$$

for every nonzero vector $\xi \in T_xM$ *and*

$$f(x) \geq c \qquad \Longrightarrow \qquad df(x)JX_{\mu(x)}(x) \geq 0.$$

Moreover, $\int_{S^2} v^*\omega = 0$ *for every smooth map* $v : S^2 \to M$*.*

This hypothesis implies that $\sup_P (f \circ u) \leq c$ for every solution (u,A) of (1) over any Riemann surface and in any homology class (see [CGMS]). In [CGS] it is shown that (H2) follows from (H1) in the case of linear actions on $\mathbb{C}^n$. In [CGMS] it is shown that the moduli space $\mathcal{M}_{B,\Sigma}$ is a smooth compact manifold of dimension

$$\dim\, \mathcal{M}_{B,\Sigma} = \left(\frac{1}{2}\dim\, M - \dim\, \mathrm{G}\right)\chi(\Sigma) + 2\langle c_1^{\mathrm{G}}(TM), B\rangle$$

for a generic J, provided that $(H1-2)$ are satisfied, B is a nontorsion homology class, and the area of Σ is sufficiently large. The latter condition, together with the energy identity, guarantees that every solution of (1) is somewhere close to the zero set of the moment map. The class $c_1^{\mathrm{G}}(TM) \in H^2(M_{\mathrm{G}};\mathbb{Z})$ in the dimension formula denotes the equivariant first Chern class of the complex vector bundle (TM,J).

Consider the evaluation map $\mathrm{ev}_{\mathrm{G}} : \mathcal{M}_{B,\Sigma} \to M_{\mathrm{G}}$, defined by

$$\mathrm{ev}_{\mathrm{G}}([u,A]) := [u(p_0), \Theta_0(u,A)],$$

where $p_0 \in P$ is fixed and $\Theta_0 : \widetilde{\mathcal{M}}_{B,\Sigma} \to \mathrm{EG}$ is a smooth map such that

$$\Theta_0(g^{-1}u, g^*A) = g(p_0)^{-1}\Theta_0(u, A).$$

This means that Θ_0 is a classifying map for the principal G-bundle $\mathcal{P}_{B,\Sigma} \to \mathcal{M}_{B,\Sigma}$ obtained as the quotient of $\widetilde{\mathcal{M}}_{B,\Sigma}$ by the based gauge group $\mathcal{G}_0 := \{g \in \mathcal{G} \,|\, g(p_0) = \mathbb{1}\}$. Let $\alpha \in H^*(M_{\mathrm{G}}; \mathbb{Z})$ be a class of degree $\deg(\alpha) = \dim \mathcal{M}_{B,\Sigma}$ and define

$$\Phi_{B,\Sigma}(\alpha) := \int_{\mathcal{M}_{B,\Sigma}} \mathrm{ev}_{\mathrm{G}}^*\alpha.$$

In [CGMS] it is shown that this integer is independent of the almost complex structure J, the metric on Σ, and the point p_0 used to define it.

Now let $D := \{z \in \mathbb{C} \,|\, |z| \le 1\}$ and consider the space of maps $v : D \to M$ that map the boundary ∂D to a G-orbit in $\mu^{-1}(0)$:

$$\mathcal{V} := \left\{v : D \to M \,|\, \exists g : \mathbb{R}/2\pi\mathbb{Z} \to \mathrm{G} \;\exists x \in \mu^{-1}(0) \;\forall \theta \in \mathbb{R} \;\; v(e^{i\theta}) = g(\theta)x\right\}.$$

Let $\mathrm{m} : \mathcal{V} \to \mathbb{Z}$ denote the function which assigns to each element $v \in \mathcal{V}$ the Maslov index of the loop of symplectic matrices obtained from the linear maps $g(\theta) : T_xM \to T_{g(\theta)x}M$ in a trivialization along v. Every smooth map $\bar{v} : S^2 \to \bar{M}$ lifts to a map $v \in \mathcal{V}$ and in this case the Maslov index $\mathrm{m}(v)$ is equal to the first Chern number $\langle c_1(T\bar{M}), \bar{v}_*[S^2]\rangle$. The minimal Maslov number will be denoted by

$$N := \inf_{v \in \mathcal{V},\, \mathrm{m}(v) > 0} \mathrm{m}(v).$$

This is a lower bound for the minimal Chern number of $\bar{M}$. We impose a third hypothesis.

(H3) *There exists a constant* $\tau > 0$ *such that*

$$\int_D v^*\omega = \tau \mathrm{m}(v)$$

for every $v \in \mathcal{V}$.

This hypothesis implies that the quotient $\bar{M}$ is a monotone symplectic manifold and that the energy of every holomorphic sphere in $\bar{M}$ is an integer multiple of $\hbar := \tau N$. The main result of this paper asserts that under hypotheses (H1-3) the invariant Φ_{B,S^2} agrees with the corresponding genus zero Gromov–Witten invariant of $\bar{M}$, provided that the cohomology classes α_i have degrees less than $2N$.

The main theorem

Theorem A. *Assume* (H1-3) *and let* $\bar{B} \in H_2(\bar{M};\mathbb{Z})$ *and* $\alpha_1, \dots, \alpha_k \in H^*_{\mathrm{G}}(M;\mathbb{Z})$ *be given such that*

$$\deg(\alpha_i) < 2N$$

for $i = 1, \dots, k$ *and*

$$\sum_{i=1}^{k} \deg(\alpha_i) := \left(\frac{1}{2}\dim M - \dim \mathrm{G}\right)\chi(\Sigma) + 2\langle c_1^{\mathrm{G}}(TM), B\rangle,$$

where $B := \kappa(\bar{B}) \in H_2^{\mathrm{G}}(M;\mathbb{Z})$*. Then*

$$\Phi_{B,S^2}(\alpha_1 \smile \cdots \smile \alpha_k) = \mathrm{G}W_{\bar{B},S^2}(\bar{\alpha}_1, \dots, \bar{\alpha}_k),$$

where $\bar{\alpha}_i := \kappa(\alpha_i) \in H^*(\bar{M};\mathbb{Z})$*.*

Remarks. (i) In the definition of $\Phi_{B,\Sigma}$ the point $p_0 \in P$ at which the map u is evaluated is fixed and the cohomology class $\mathrm{ev}_{\mathrm{G}}^*\alpha \in H^*(\mathcal{M}_{B,\Sigma};\mathbb{Z})$ is independent of the choice of the point p_0 used in the definition of ev_{G}. The Gromov–Witten invariants in Theorem A are also to be understood with fixed marked points on S^2 in the definitions of the evaluation maps, and with almost complex structures that are allowed to depend on the base point $z \in S^2$.

(ii) If, in addition to $(H3)$, we assume $[\bar{\omega}] = \tau c_1(T\bar{M})$ then the proof of Theorem A goes through word by word for the higher genus case (with fixed marked points). In general, the extension to general Riemann surfaces requires a refined version of the compactness theorem in Section 12 which takes account of the preservation of the homotopy class in the limit, as in Gromov compactness. With similar refined arguments one should be able to deal with the case of varying marked points or of varying complex structures on Σ.

(iii) The assertion of Theorem A does not continue to hold in the case $\deg(\alpha_i) \geq 2N$. For example, consider the standard action of S^1 on $\mathbb{C}^n$, let $P \to S^2$ be an S^1-bundle of degree $d \geq 0$, and denote by $c \in H^2(\mathrm{B}S^1;\mathbb{Z}) = H^2_{S^1}(\mathbb{C}^n;\mathbb{Z})$ the positive generator. Then the minimal Chern number is $N = n$, the dimension of $\mathcal{M}_{d,S^2}$ is $2nd + 2n - 2$, and we have $\Phi_{d,S^2}(c^m) = 1$ whenever $m = nd + n - 1$. The corresponding Gromov–Witten invariant (for a k-tuple of classes $c^{m_1}, \dots, c^{m_k}$ with $m_1 + \cdots + m_k = m$) counts holomorphic spheres of degree d in $\mathbb{C}P^{n-1}$ passing at k given points $z_1, \dots, z_k \in S^2$ through generic copies of $\mathbb{C}P^{n-1-m_i}$ for $i = 1, \dots, k$. Thus the Gromov–Witten invariant is zero whenever $\deg(c^{m_i}) = 2m_i \geq 2n$ for some i.

Equivariant and quantum cohomology

Consider the monotone case. The quantum cohomology $\mathrm{QH}^*(\bar{M})$ of $\bar{M}$ is the ring of all formal sums of the form

$$\bar{\alpha} = \sum_{\bar{B} \in H_2(\bar{M};\mathbb{Z})} \bar{\alpha}_{\bar{B}} e^{\bar{B}},$$

where $\bar{\alpha}_{\bar{B}} \in H^*(\bar{M};\mathbb{Z})/\text{torsion}$, such that

$$\#\left\{\bar{B} \in H_2(\bar{M};\mathbb{Z}) \,|\, \bar{\alpha}_{\bar{B}} \neq 0,\, \langle[\bar{\omega}], \bar{B}\rangle \leq c\right\} < \infty$$

for all $c > 0$. The degree convention is $\deg(e^{\bar{B}}) := 2\langle c_1(T\bar{M}), \bar{B}\rangle$. Choose an integral basis $\bar{e}_0, \ldots, \bar{e}_n$ of $H^*(\bar{M};\mathbb{Z})/\text{torsion}$ and let $\bar{e}_i^*$ denote the dual basis in the sense that

$$\int_{\bar{M}} \bar{e}_i \smile \bar{e}_j^* = \delta_{ij}.$$

Then the product structure on $\mathrm{QH}^*(\bar{M})$ is defined by

$$\bar{\alpha}_1 * \bar{\alpha}_2 := \sum_{\bar{B}_1,\bar{B}_2,\bar{B}} \sum_{i=0}^{n} \mathrm{G}W_{\bar{B}-\bar{B}_1-\bar{B}_2,S^2}(\bar{\alpha}_{1\bar{B}_1}, \bar{\alpha}_{2\bar{B}_2}, \bar{e}_i^*)\bar{e}_i e^{\bar{B}}.$$

The sum is over all quadruples $i, \bar{B}_1, \bar{B}_2, \bar{B}$ such that

$$\deg(\bar{\alpha}_1) + \deg(\bar{\alpha}_2) = \deg(\bar{e}_i) + 2c_1(\bar{B}),$$

where we abbreviate $c_1(\bar{B}) := \langle c_1(T\bar{M}), \bar{B}\rangle$.

The Gromov–Witten invariant associated to a Riemann surface Σ, with a fixed complex structure j_Σ and fixed marked points $z_1, \ldots, z_k$, can be extended to a map $\mathrm{G}W_{\bar{B},\Sigma} : \mathrm{QH}^*(\bar{M}) \otimes \cdots \otimes \mathrm{QH}^*(\bar{M}) \to \mathbb{Z}$ by the formula

$$\mathrm{G}W_{\bar{B},\Sigma}(\bar{\alpha}_1, \ldots, \bar{\alpha}_k) := \sum_{\bar{B}_i} \mathrm{G}W_{\bar{B}-\bar{B}_1-\cdots-\bar{B}_k,\Sigma}(\bar{\alpha}_{1\bar{B}_1}, \ldots, \bar{\alpha}_{k\bar{B}_k}).$$

With this convention the gluing formula for the Gromov–Witten invariants [MS1, RT] can be expressed in the form

$$\mathrm{G}W_{\bar{B},\Sigma}(\bar{\alpha}_1, \ldots, \bar{\alpha}_k) = \mathrm{G}W_{\bar{B},\Sigma}(\bar{\alpha}_1 * \cdots * \bar{\alpha}_k). \tag{2}$$

We abbreviate $H^*(X) := H^*(X;\mathbb{Z})/\text{torsion}$.

Corollary A'. *Assume* $(H1-3)$ *and suppose that* $H^*(M_{\mathrm{G}})$ *is generated by classes of degree less than* $2N$*. Then there exists a unique (surjective) ring homomorphism* $\varphi : H^*(M_{\mathrm{G}}) \longrightarrow \mathrm{QH}^*(\bar{M})$ *such that, for every* $\alpha \in H^*(M_{\mathrm{G}})$ *and every* $\bar{B} \in H_2(\bar{M};\mathbb{Z})$,

$$\deg(\alpha) < 2N \qquad \Longrightarrow \qquad \varphi(\alpha) = \kappa(\alpha),$$

$$\Phi_{\kappa(\bar{B}),S^2}(\alpha) = \mathrm{GW}_{\bar{B},S^2}(\varphi(\alpha)).$$

Proof. Let $\alpha \in H^*(M_{\mathrm{G}})$ and choose $\alpha_{ij} \in H^*(M_{\mathrm{G}})$ such that $\deg(\alpha_{ij}) < 2N$ and

$$\alpha = \sum_{i=1}^{k} \alpha_{i1} \smile \cdots \smile \alpha_{i\ell}. \tag{3}$$

Define

$$\varphi(\alpha) := \sum_{i=1}^{k} \kappa(\alpha_{i1}) * \cdots * \kappa(\alpha_{i\ell}). \tag{4}$$

We prove that $\varphi(\alpha)$ is independent of the choice of α_{ij}. To see this, note that, since the cohomology of $\bar{M}$ is generated by classes of degree less than $2N$, so is the quantum cohomology. This means that a quantum cohomology class $\bar{\alpha} \in \mathrm{QH}^*(\bar{M})$ is zero if and only if $\mathrm{GW}_{\bar{B},S^2}(\bar{\alpha},\bar{\beta}_1,\ldots,\bar{\beta}_m) = 0$ for every $\bar{B} \in H_2(\bar{M};\mathbb{Z})$ and all $\bar{\beta}_1,\ldots\bar{\beta}_m \in H^*(\bar{M})$ such that $\deg(\bar{\beta}_j) < 2N$ for all j. Now suppose that the expression on the right of (4) is nonzero. Then, by what we have just observed, there exist cohomology classes $\bar{\beta}_1,\ldots,\bar{\beta}_m$ of degrees less than $2N$ and a homology class $\bar{B} \in H_2(\bar{M};\mathbb{Z})$ such that

$$\sum_{i=1}^{k} \mathrm{GW}_{\bar{B},S^2}(\kappa(\alpha_{i1}),\ldots,\kappa(\alpha_{i\ell}),\bar{\beta}_1,\ldots,\bar{\beta}_m) \neq 0.$$

Since the homomorphism $\kappa : H^*(M_{\mathrm{G}}) \to H^*(\bar{M})$ is surjective (cf. [K]), there exist classes $\beta_j \in H^*(M_{\mathrm{G}})$ (of degrees less than $2N$) such that $\kappa(\beta_j) = \bar{\beta}_j$ for every j. Hence, by Theorem A,

$$\sum_{i=1}^{k} \Phi_{\kappa(\bar{B}),S^2}(\alpha_{i1} \smile \cdots \smile \alpha_{i\ell} \smile \beta_1 \smile \cdots \smile \beta_m) \neq 0,$$

and hence $\alpha \neq 0$. This shows that φ is well defined. The map φ is obviously a ring homomorphism. The formula $\Phi_{B,S^2}(\alpha) = \mathrm{GW}_{\bar{B},S^2}(\varphi(\alpha))$ follows immediately from Theorem A and the gluing formula (2) for the Gromov–Witten invariants. $\square$

The homomorphism $\varphi : H^*(M_{\mathrm{G}}) \to \mathrm{QH}^*(\bar{M})$ can be defined geometrically in terms of the vortex equations over $\mathbb{C}$:

$$\partial_s u + L_u\Phi + J(\partial_t u + L_u\Psi) = 0, \qquad \partial_s\Psi - \partial_t\Phi + [\Phi, \Psi] + \mu(u) = 0. \tag{5}$$

For every finite energy solutions of (5) in radial gauge there exist a loop $g : S^1 \to \mathrm{G}$ and a point $x_0 \in \mu^{-1}(0)$ such that

$$\lim_{r\to\infty} u(re^{i\theta}) = g(e^{i\theta})x_0 \tag{6}$$

(see Section 11). Every map $u : \mathbb{C} \to M$ that satisfies (6) determines an equivariant homology class $B = [u] \in H_2(M_{\mathrm{G}}; \mathbb{Z})$. Now the moduli space $\mathcal{M}_B(J)$ of gauge equivalence classes of solutions of (5) and (6) that represent the class B has two evaluation maps $\mathrm{ev}_0 : \mathcal{M}_B \to M_{\mathrm{G}}$ and $\mathrm{ev}_\infty : \mathcal{M}_B \to \bar{M}$. The map φ can be defined by

$$\varphi(\alpha) = \sum_{i=0}^{n} \sum_{\bar{B}} \left(\int_{\mathcal{M}_{\kappa(\bar{B})}} \mathrm{ev}_0^*\alpha \smile \mathrm{ev}_\infty^*\bar{e}_i^* \right) \bar{e}_i e^{\bar{B}}.$$

The details of this construction will be carried out elsewhere.

Outline of the proof of Theorem A

The proof of Theorem A is based on an adiabatic limit argument in which the metric on the Riemann surface is scaled by a large factor ε^{-2}. Then equations (1) have the form

$$\bar{\partial}_{J,A}(u) = 0, \qquad *F_A + \varepsilon^{-2}\mu(u) = 0. \tag{7}$$

The solutions of (7) minimize the ε-dependent energy

$$E^\varepsilon(u, A) = \frac{1}{2}\int_\Sigma \left(|d_A u|^2 + \varepsilon^2|F_A|^2 + \varepsilon^{-2}|\mu(u)|^2\right) \mathrm{dvol}_\Sigma,$$

and the value of this functional at a solution of (7) is independent of ε in a given equivariant homology class. In this paper we examine the limit behaviour of the solutions of (7) as ε tends to zero for Riemann surfaces of any genus. The limit equations have the form

$$\bar{\partial}_{J,A}(u) = 0, \qquad \mu(u) = 0. \tag{8}$$

The solutions of (8) can be interpreted as pseudoholomorphic curves in the symplectic quotient $\bar{M} = \mu^{-1}(0)/\mathrm{G}$ with respect to the induced family of almost complex structures $\bar{J}_z$ (see Section 2). We impose a further hypothesis that is satisfied for a generic family of G-invariant almost complex structures on M:

(H4) *Every nonconstant $\bar{J}$-holomorphic curve $\bar{u} : \Sigma \to \bar{M}$ is regular in the sense that the linearized Cauchy-Riemann operator along $\bar{u}$ is surjective.*

This hypothesis guarantees that the moduli space of holomorphic curves in $\bar{M}$ is smooth.

The proof of Theorem A requires three preliminary theorems which are of interest in their own rights. Theorem B constructs a $\mathcal{G}(P)$-equivariant map

$$(u_0, A_0) \mapsto (u_\varepsilon, A_\varepsilon) =: \widetilde{\mathcal{T}}^\varepsilon(u_0, A_0)$$

which assigns to every regular solution of (8) a nearby solution of (7) for $\varepsilon > 0$ sufficiently small. How small ε must be chosen depends (continuously) on the given pair (u_0, A_0). Theorem C shows that the map $\widetilde{\mathcal{T}}^\varepsilon$ constructed in Theorem B is *locally surjective* in the sense that every solution of (7) that is sufficiently close to a solution (u_0, A_0) of (8) must be in the image of $\widetilde{\mathcal{T}}^\varepsilon$. The neighbourhood in which surjectivity holds depends on ε: it becomes smaller as ε tends to zero. Theorem D strengthens the local surjectivity result of Theorem C. We remove the assumption that the solution of (7) is close to some given solution of (8). However, we consider only solutions of (7) that satisfy a suitable L^∞-bound on the first derivatives and prove that every solution of (7) that satisfies this bound lies in the image of $\widetilde{\mathcal{T}}^\varepsilon$ for ε small. The proof of Theorem A is then based on a bubbling argument in the small ε limit which establishes a one-to-one correspondence between the solutions of (8) and those of (7) in a zero dimensional setting, where additional conditions have been imposed.

In Section 2 we review standard results about the moduli space of pseudoholomorphic curves in the symplectic quotient $M /\!\!/ \mathrm{G}$ and rephrase them in terms of solutions of (8). Theorems B, C, and D will be stated in Section 3. The remaining sections are devoted to the proofs of the four main theorems.

While the general outline of the proof of Theorem A is analogous to the proof of the Atiyah–Floer conjecture in [DS1, DS2] there are several new ingredients in the present paper.

In [DS2] the moduli space of Floer connecting orbits is a finite set, while the moduli space $\mathcal{M}^0$ of pseudoholomorphic curves is, in general, a (noncompact) manifold of positive dimension. Hence, in constructing the map $\mathcal{T}^\varepsilon$ from (a compact subset of) $\mathcal{M}^0$ to $\mathcal{M}^\varepsilon$, care must be taken to establish that the constants depend continuously on the point in $\mathcal{M}^0$. This refers to the linear and quadratic estimates needed in the proof of Theorem B (Sections 4 and 5). Secondly, we extend the estimates of [DS1] for the $(1, p, \varepsilon)$-norms to the $(2, p, \varepsilon)$-norms (see Lemmata 4.5-4.8 and Proposition 5.1). For the linear estimates the extension to the higher derivatives is quite subtle because of the ε-dependent norms. Another new ingredient arises from the presence of nonlinearities in the highest order terms of the Cauchy–Riemann equations. This requires more delicate quadratic estimates (Proposition 5.1) for the proof of local uniqueness (Theorem 6.2).

The proof of the ε-local slice theorem in Section 7 (which can be viewed as a simpler analogue of Theorem B) is considerably harder than the analogous result in [DS2], since a) we must establish estimates for the $(2, p, \varepsilon)$-norms, b) we must prove that the constants depend continuously on the point in $\mathcal{M}^0$, c) the manifold M is not an affine space so we must deal with additional lower order terms, and d) we give a proof of the linear estimate in Lemma 7.3 for $p > 2$ (the analogue in [DS2] was only established for $p = 2$).

The proof of local surjectivity (Theorem C) requires a subtle tubular neighbourhood theorem for the moduli spaces $\mathcal{M}^\varepsilon$. The result is quantitative with constants independent of ε. In particular, the proof involves an estimate for the derivative of the map $(u_0, A_0) \to (\xi_0, \alpha_0, g)$ given by the ε-local slice theorem. This is where the estimates for the $(2, p, \varepsilon)$-norms are needed. As a result the entire adiabatic limit argument has to be carried out for these higher norms. In comparison, the analogous result in [DS2, Proposition 6.3] can be disposed of with a simple time shift argument and only requires estimates in the $(1, p, \varepsilon)$-norm.

Another new ingredient in the present paper is the apriori estimate in Lemma 9.1. It asserts that every solution of (7) which satisfies a certain L^∞ bound must be $\varepsilon^{3/2}$-close to the zero set of the moment map. As a consequence we obtain in Theorem 10.1 a much stronger surjectivity result for the map $\mathcal{T}^\varepsilon$; compare with [DS2, Theorem 8.1]. This strenghtened form of [DS2, Theorem 8.1] is needed to close a gap in the proof of [DS2, Theorem 9.1], namely to prove that the holomorphic sphere in the symplectic quotient appearing in the bubbling argument on page 634 in [DS2] is nonconstant. At the same time the bubbling argument in the proof of [DS2, Theorem 9.1] can be simplified: it suffices to work with the sequence $c_\nu := \sup(|d_{A_\nu} u_\nu| + \varepsilon_\nu^{-1}|F_{A_\nu}|)$ instead of $c_\nu := \sup(|d_{A_\nu} u_\nu| + \varepsilon_\nu^{-1}|F_{A_\nu}|^{1/2})$. The modified bubbling argument is carried out in the present context in Section 12.

In Section 11 we establish the asymptotic behaviour and the quantization of the energy for solutions of the nonlinear vortex equations on the complex plane. (An analogous result for anti-self-dual instantons is used without proof in [DS2].) In Section 12 we construct a classifying map on an open set in $C^\infty_{\mathrm{G}}(P, M) \times \mathcal{A}(P)$, which contains the moduli spaces $\mathcal{M}^\varepsilon$ for all $\varepsilon \in [0, \varepsilon_0]$, with values in a finite dimensional approximation of EG; and we prove C^1-convergence for the composition of the resulting evaluation map with $\mathcal{T}^\varepsilon$. All these results are needed in the proof of our main theorem.

2 Pseudoholomorphic curves

For $z \in \Sigma$ let $\bar{J}_z$ denote the almost complex structure on $\bar{M}$ induced by J_z, let $\bar{P} \longrightarrow \bar{M}$ denote the principal G-bundle $\bar{P} := \mu^{-1}(0) \subset M$, and let $\bar{A}$ denote the connection on $\bar{P}$ determined by ω and J. If (u, A) is a solution of (8) then u descends to a $\bar{J}$-holomorphic curve $\bar{u} : \Sigma \to \bar{M}$ and A is the pullback of $\bar{A}$ under $\bar{u}$. Two gauge equivalent solutions descend to the same map $\bar{u}$ and every $\bar{J}$-holomorphic curve $\bar{u} : \Sigma \to \bar{M}$ lifts to a solution of (8) for some principal G-bundle P (isomorphic to the pullback of $\bar{P}$ under $\bar{u}$).

Fix a homology class $\bar{B} \in H_2(\bar{M}; \mathbb{Z})$, let $B := \kappa(\bar{B}) \in H_2^{\mathrm{G}}(M; \mathbb{Z})$, and consider the space

$$\widetilde{\mathcal{M}}^0_{B,\Sigma} := \{(u, A) \in C^\infty_{\mathrm{G}}(P, M) \times \mathcal{A}(P) \,|\, [u] = B \text{ and (8) holds}\}.$$

This space is invariant under the action of the gauge group $\mathcal{G}(P)$. Under our standing hypothesis (H4) the quotient

$$\mathcal{M}^0_{B,\Sigma} := \widetilde{\mathcal{M}}^0_{B,\Sigma} / \mathcal{G}(P)$$

is a smooth manifold of dimension

$$\dim \mathcal{M}^0_{B,\Sigma} = \left(\frac{1}{2} \dim M - \dim \mathrm{G}\right) \chi(\Sigma) + 2\langle c_1^{\mathrm{G}}(TM), B\rangle$$

(see [MS1, Theorem 3.3.4]). Note that $\mathcal{M}_{B,\Sigma}$ and $\mathcal{M}^0_{B,\Sigma}$ have the same dimension.

For later reference we now introduce explicit notation for a local parametrization of $\mathcal{M}^0_{B,\Sigma}$ by the kernel of the linearized operator. Linearizing equations (8) at a solution (u, A) gives rise to the Cauchy–Riemann operator

$$\mathcal{D}^0 := \mathcal{D}^0_{(u,A)} : \Omega^0(\Sigma, H_u) \to \Omega^{0,1}(\Sigma, H_u)$$

given by

$$\mathcal{D}^0\xi_0 := \pi_u D_{(u,A)}\xi_0,$$

where $D_{(u,A)} : \Omega^0(\Sigma, u^*TM/\mathrm{G}) \to \Omega^{0,1}(u^*TM/\mathrm{G})$ is the operator (109) in Appendix B. The bundle $H_u \to \Sigma$ and the projection $\pi_u : u^*TM \to H_u$ are defined as follows. Consider the bundle $H \longrightarrow \Sigma \times \mu^{-1}(0)$ with fibres

$$H_{z,x} := \ker\, d\mu(x) \cap \ker\, d\mu(x)J_z.$$

There is an orthogonal decomposition

$$T_xM = \operatorname{im} L_x \oplus H_{z,x} \oplus \operatorname{im} J_zL_x$$

for every $(z, x) \in \Sigma \times \mu^{-1}(0)$, where $L_x : \mathfrak{g} \to T_xM$ the infinitesimal action, i.e.

$$L_x\eta := X_\eta(x).$$

Its dual operator with respect to the metric determined by J_z is given by

$$L_x^* = L_x^{*z} = d\mu(x)J_z(x).$$

Now let $u : P \to \mu^{-1}(0)$ be an equivariant map and consider the pullback of H under the map $\tilde{u} : P \to \Sigma \times \mu^{-1}(0)$, given by $\tilde{u}(p) := (\pi(p), u(p))$. This pullback is a G-equivariant vector bundle over P and its quotient

$$H_u := \tilde{u}^*H/\mathrm{G} \longrightarrow \Sigma$$

is naturally isomorphic to the pullback of the tangent bundle $T\bar{M}$ under the induced map $\bar{u} : \Sigma \to \bar{M}$. Let $\pi_u : u^*TM/\mathrm{G} \to H_u$ denote the orthogonal projection onto the harmonic part. Thus $\pi_u[\xi] := [\pi_u\xi]$ where the lifted projection $u^*TM \to u^*H$ (also denoted by π_u) is given by

$$\pi_u\xi := \xi - L_u(L_u^*L_u)^{-1}L_u^*\xi + JL_u(L_u^*L_u)^{-1}L_u^*J\xi \tag{9}$$

for a G-equivariant section $\xi : P \to u^*TM$.

Theorem 2.1. *Assume* (H1) *and* (H4) *and fix a constant* $p > 2$. *For every* $(\bar{u}_0, \bar{A}_0) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}$ *there exist a sequence of positive constants* $\delta, c, c_1, c_2, \ldots$ *and a map*

$$\mathcal{F}^0 := \mathcal{F}^0_{(\bar{u}_0,\bar{A}_0)} : B^0_\delta \to \widetilde{\mathcal{M}}^0_{B,\Sigma}, \qquad B^0_\delta := \{\xi_0 \in \ker \mathcal{D}^0_{(\bar{u}_0,\bar{A}_0)} \mid \|\xi_0\|_{L^p} < \delta\},$$

such that the following holds.
(i) *If* $\xi_0 \in \mathcal{B}^0_\delta$ *then there exists a unique pair of sections* $\xi_1 \in \Omega^0(\Sigma, H_{\bar{u}_0})$ *and* $\xi_2 \in \Omega^0(\Sigma, \mathrm{im}\, JL_{\bar{u}_0}/\mathrm{G})$ *such that*

$$\xi_1 \in \mathrm{im}\,(\mathcal{D}^0_{(\bar{u}_0,\bar{A}_0)})^*, \qquad \|\xi_1\|_{W^{1,p}} + \|\xi_2\|_{W^{1,p}} \leq c\,\|\xi_0\|_{W^{1,p}}\,.$$

and the pair (u_0, A_0)*, given by*

$$u_0 := \exp^{\mu^{-1}(0)}_{\bar{u}_0}(\xi_0 + \xi_1) = \exp_{\bar{u}_0}(\xi_0 + \xi_1 + \xi_2), \quad A_0 := -(L^*_{u_0} L_{u_0})^{-1} L^*_{u_0} du_0,$$

satisfies (8). The pair (u_0, A_0) *is the image of* ξ_0 *under* $\mathcal{F}^0$.
(ii) *For every integer* $k \geq 1$ *and every* $\xi_0 \in B^0_\delta$ *we have*

$$\|\xi_1\|_{W^{k,p}} + \|\xi_2\|_{W^{k,p}} \leq c_k\,\|\xi_0\|^2_{W^{k,p}}\,, \qquad \left\|A_0 - \bar{A}_0\right\|_{W^{k,p}} \leq c_k\,\|\xi_0\|_{W^{k,p}}\,,$$

where ξ_1, ξ_2, *and* A_0 *are as in (i).*
(iii) *The map* $\mathcal{F}^0$ *is smooth and* $d\mathcal{F}^0(0)\xi_0 = (\xi_0, \alpha_0)$, *where* $\alpha_0 \in \Omega^1(\Sigma, \mathfrak{g}_P)$ *is uniquely determined by the equation*

$$D\bar{\partial}_{J,\bar{A}_0}(\bar{u}_0)\xi_0 + X_{\alpha_0}(\bar{u}_0)^{0,1} = 0.$$

Theorem 2.1 is a standard result in the theory of holomorphic curves (cf. [MS1]). It follows from Fredholm theory and an infinite dimensional version of the implicit function theorem. In most applications the moduli space $\mathcal{M}^0_{B,\Sigma}$ is not compact. However, it can be exhausted by the compact subsets

$$\mathcal{M}^0_{B,\Sigma}(c_0) := \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)/\mathcal{G}(P),$$

where $c_0 > 0$ and

$$\widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0) := \left\{(u, A) \in \widetilde{\mathcal{M}}^0_{B,\Sigma} \mid \|d_A u\|_{L^\infty} + \|F_A\|_{L^\infty} \leq c_0\right\}.$$

Note that $\widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$ is invariant under the action of $\mathcal{G}(P)$. For later reference we prove the following lemma.

Lemma 2.2. *Fix a reference connection $\hat{A} \in \mathcal{A}(P)$. Then, for every $c_0 > 0$ and every integer $\ell \in \mathbb{N}$, there exists a constant $c = c(c_0, \ell) > 0$ such that, for every $(u_0, A_0) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$, we have*

$$\inf_{g \in \mathcal{G}(P)} \left(\|g^{-1}u\|_{C^\ell} + \|g^*A_0 - \hat{A}\|_{C^\ell} \right) \leq c.$$

Proof. Suppose, by contradiction, that there exists a sequence $(u_\nu, A_\nu) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$ such that $\|g^{-1}u_\nu\|_{C^\ell} + \|g^*A_\nu - \hat{A}\|_{C^\ell} \geq \nu$ for every ν and every $g \in \mathcal{G}(P)$. By [MS1, Theorem B.4.2] there exists a subsequence, still denoted by (u_ν, A_ν), such that the induced maps $\bar{u}_\nu : \Sigma \to M/\!\!/\mathrm{G}$ converge in the C^∞-topology to a smooth $\bar{J}$-holomorphic curve. The limit curve represents the same homotopy class as the approximating curves and hence can be represented by a pair $(u, A) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$. Since the sequence $\bar{u}_\nu$ converges to $\bar{u} : \Sigma \to M/\!\!/\mathrm{G}$ in the $C^{\ell+1}$-topology, there exists a constant $\nu_0 \geq 0$ such that, for every $\nu \geq \nu_0$, there exist a gauge transformation g_ν and a section $\xi_\nu \in \Omega^0(\Sigma, H_u)$ such that

$$g_\nu^{-1}u_\nu = \exp_u(\xi_\nu), \qquad \lim_{\nu \to \infty} \|\xi_\nu\|_{C^{\ell+1}} = 0.$$

The formulae

$$g_\nu^* A_\nu = -(L^*_{g_\nu^{-1}u_\nu} L_{g_\nu^{-1}u_\nu})^{-1} L_{g_\nu^{-1}u_\nu} d(g_\nu^{-1}u_\nu), \qquad A = -(L_u^* L_u)^{-1} L_u du,$$

show that $g_\nu^*(u_\nu, A_\nu)$ converges to (u, A) in the C^ℓ topology. This contradicts the choice of the sequence (u_ν, A_ν) and hence proves the lemma. □

Theorem 2.3. *Assume (H1) and (H4), let $\bar{B} \in H_2(M; \mathbb{Z})$ be a nontorsion homology class, and let $(\Sigma, \mathrm{dvol}_\Sigma, j_\Sigma)$ be a compact Riemann surface. Then, for every $c_0 > 0$ and every $p > 2$, there exist positive constants c and δ such that the following holds. If $\bar{u} : \Sigma \to \bar{M}$ is a smooth map such that $[\bar{u}] = \bar{B}$ and*

$$\|d\bar{u}\|_{L^\infty} \leq c_0, \qquad \|\bar{\partial}_{\bar{J}}(\bar{u})\|_{L^p} \leq \delta$$

then there exists a section $\bar{\xi} \in \Omega^0(\Sigma, \bar{u}^*T\bar{M})$ such that

$$\bar{\partial}_{\bar{J}}(\exp_{\bar{u}}(\bar{\xi})) = 0, \qquad \|\bar{\xi}\|_{W^{1,p}} \leq c\|\bar{\partial}_{\bar{J}}(\bar{u})\|_{L^p}.$$

Proof. This is again a standard result for pseudoholomorphic curves and the proof is almost word by word the same as that of [DS2, Theorem 2.5]. Here is a sketch. One argues by contradiction. If the result were false, there would be a sequence of smooth maps $\bar{u}_i : \Sigma \to \bar{M}$ that satisfies

$$\sup_i \|d\bar{u}_i\|_{L^\infty} < \infty, \qquad \lim_{i\to\infty} \|\bar{\partial}_{\bar{J}}(\bar{u}_i)\|_{L^p} = 0,$$

but which does not satisfy the conclusion of the theorem for any constant c. This means that the $W^{1,p}$-distance of $\bar{u}_i$ to the space of $\bar{J}$-holomorphic curves is not controlled uniformly by the L^p-norm of $\bar{\partial}_{\bar{J}}(\bar{u}_i)$. Now, by the Arzéla–Ascoli and Banach–Alaoglu theorems, a suitable subsequence of $\bar{u}_i$ converges, strongly with respect to the sup-norm and weakly in $W^{1,p}$, to a $\bar{J}$-holomorphic curve $\bar{u}$. It follows from standard elliptic regularity for $\bar{J}$-holomorphic curves that $\bar{u}_i$ then converges strongly with respect to the $W^{1,p}$-norm. To see this, write $\bar{u}_i = \exp_{\bar{u}}(\bar{\xi}_i)$ and observe that

$$\begin{aligned} \|\bar{\xi}_i\|_{W^{1,p}} &\leq c_1 \left(\|D_{\bar{u}}\bar{\xi}_i\|_{L^p} + \|\bar{\xi}_i\|_{L^p}\right) \\ &\leq c_2 \left(\|\bar{\partial}_{\bar{J}}(\bar{u}_i)\|_{L^p} + \|\bar{\xi}_i\|_{W^{1,p}}\|\bar{\xi}_i\|_{L^\infty}\right) + c_1\|\bar{\xi}_i\|_{L^p}. \end{aligned}$$

Here the first inequality is the elliptic estimate for the Cauchy–Riemann operator $D_{\bar{u}}$ and the second is the quadratic estimate for $\bar{\partial}_{\bar{J}}$. With this established it follows from hypothesis (H4) and the implicit function theorem for the operator $\bar{\partial}_{\bar{J}}$ that there exists a sequence of $\bar{J}$-holomorphic curves $\bar{u}_{0i}$ whose $W^{1,p}$-distance to $\bar{u}_i$ is bounded above by a fixed constant times the L^p-norm of $\bar{\partial}_{\bar{J}}(\bar{u}_i)$ (see [DS2, Theorem 2.1]). This shows that the sequence $\bar{u}_i$ does after all satisfy the conclusion of the theorem, in contradiction to our assumption. □

3 Adiabatic limits

Before stating our main results we introduce some notation. Fix an equivariant homology class $\bar{B} \in H_2(\bar{M};\mathbb{Z})$, let $B := \kappa(\bar{B})$, and denote

$$\mathcal{B} := \left\{(u, A) \in C^\infty_{\mathrm{G}}(P, M) \times \mathcal{A}(P) \,|\, [u] = B\right\}.$$

This space is an infinite dimensional Frêchet manifold with tangent space

$$T_{(u,A)}\mathcal{B} := \Omega^0(\Sigma, u^*TM/\mathrm{G}) \times \Omega^1(\Sigma, \mathfrak{g}_P).$$

It carries an action of the gauge group $\mathcal{G} = \mathcal{G}(P)$ by $g^*(u, A) = (g^{-1}u, g^*A)$. Consider the vector bundle $\mathcal{E} \to \mathcal{B}$ with fibres

$$\mathcal{E}_{(u,A)} := \Omega^{0,1}(\Sigma, u^*TM/\mathrm{G}) \oplus \Omega^0(\Sigma, \mathfrak{g}_P)$$

and the $\mathcal{G}$-equivariant section $\mathcal{F}^\varepsilon : \mathcal{B} \to \mathcal{E}$ given by

$$\mathcal{F}^\varepsilon(u, A) := (\bar\partial_{J,A}(u), *F_A + \varepsilon^{-2}\mu(u)).$$

The zero set of this section is the space

$$\widetilde{\mathcal{M}}^\varepsilon_{B,\Sigma} := \{(u, A) \in \mathcal{B} \,|\, u \text{ and } A \text{ satisfy (7)}\}.$$

Its quotient by the action of the gauge group will be denoted by

$$\mathcal{M}^\varepsilon_{B,\Sigma} := \widetilde{\mathcal{M}}^\varepsilon_{B,\Sigma}/\mathcal{G}(P).$$

The following theorem asserts the existence of solutions of (7) for sufficiently small ε near every regular solution of (8). The result is quantitative and the estimates are expressed in terms of suitable ε-dependent norms. Moreover, an operator $\mathcal{D}^\varepsilon : T_{(u,A)}\mathcal{B} \to \mathcal{E}_{(u,A)} \oplus \Omega^0(\Sigma, \mathfrak{g}_P)$ appears. This operator is the augmented vertical differential of $\mathcal{F}^\varepsilon$. The operator and the norms will be defined in Section 4.

Theorem B. *Assume (H1) and (H4) and let $\bar B \in H_2(\bar M; \mathbb{Z})$ be a nontorsion homology class. Then, for every $c_0 > 0$ and every $p > 2$, there exist positive constants ε_0, c, and δ such that for every $\varepsilon \in (0, \varepsilon_0]$ there exists a $\mathcal{G}(P)$-equivariant map*

$$\widetilde{\mathcal{T}}^\varepsilon : \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0) \to \widetilde{\mathcal{M}}^\varepsilon_{B,\Sigma}$$

that satisfies the following conditions.

(a) *If $(u_0, A_0) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$ then*

$$\widetilde{\mathcal{T}}^\varepsilon(u_0, A_0) = (\exp_{u_0}(\xi_\varepsilon), A_0 + \alpha_\varepsilon),$$

where $\zeta_\varepsilon = (\xi_\varepsilon, \alpha_\varepsilon) \in T_{(u_0,A_0)}\mathcal{B}$ satisfies

$$\|\zeta_\varepsilon\|_{2,p,\varepsilon;(u_0,A_0)} \le c\varepsilon^2,$$

$$-d^*_{A_0}\alpha_\varepsilon + \varepsilon^{-2}L^*_{u_0}\xi_\varepsilon = 0, \qquad \zeta_\varepsilon \in \mathrm{im}\,(\mathcal{D}^\varepsilon_{(u_0,A_0)})^*. \tag{10}$$

(b) *If $(u_0, A_0) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$ and $(u, A) = (\exp_{u_0}(\xi), A_0 + \alpha) \in \widetilde{\mathcal{M}}^\varepsilon_{B,\Sigma}$ where $\zeta = (\xi, \alpha) \in T_{(u_0,A_0)}\mathcal{B}$ satisfies (10) and*

$$\|\zeta\|_{1,p,\varepsilon;(u_0,A_0)} \le \delta\varepsilon^{2/p+1/2},$$

then $(u, A) = \widetilde{\mathcal{T}}^\varepsilon(u_0, A_0)$.

The map $\widetilde{\mathcal{T}}^{\varepsilon}$ of Theorem A descends to a map between the quotient spaces which we denote by

$$\mathcal{T}^{\varepsilon}:\mathcal{M}^0_{B,\Sigma}(c_0)\longrightarrow\mathcal{M}^{\varepsilon}_{B,\Sigma}.$$

Assertion (a) is proved by a Newton type iteration (see Section 6). It requires linear and quadratic estimates for the ε-dependent norms with constants that are independent of ε. These estimates are proved in Sections 4 and 5. Assertion (b) is a strengthened form of the corresponding uniqueness statement. Here the neighbourhood in which uniqueness holds is larger than in the existence result (namely it is of radius $c\varepsilon^{2/p+1/2}$ instead of $c\varepsilon^2$). The uniqueness statement shows that the maps $\mathcal{T}^{\varepsilon}$ are independent of c_0 in the sense that two such maps corresponding to different values of c_0 (but the same value of ε) agree on the intersection of their domains. The next theorem shows that $\mathcal{T}^{\varepsilon}$ is locally surjective.

Theorem C. *Assume* $(H1)$ *and* $(H4)$ *and let* $\bar{B}\in H_2(\bar{M};\mathbb{Z})$ *be a nontorsion homology class. Then, for every* $c_0>0$ *and every* $p>2$*, there exist positive constants* ε_0 *and* δ *such that the following holds for every* $\varepsilon\in(0,\varepsilon_0]$*. If*

$$(\bar{u}_0,\bar{A}_0)\in\widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0-1),\qquad(u,A)=(\exp_{\bar{u}_0}(\bar{\xi}),\bar{A}_0+\bar{\alpha})\in\widetilde{\mathcal{M}}^{\varepsilon}_{B,\Sigma},$$

where $\bar{\zeta}=(\bar{\xi},\bar{\alpha})\in T_{(\bar{u}_0,\bar{A}_0)}\mathcal{B}$ *satisfies*

$$\left\|\bar{\zeta}\right\|_{1,p,\varepsilon;(\bar{u}_0,\bar{A}_0)}\le\delta\varepsilon^{2/p+1/2},$$

then $(u,A)\in\widetilde{\mathcal{T}}^{\varepsilon}(\widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0))$*.*

This result is restated more precisely in Theorem 8.1 in Section 8. There it is proved that

$$g^*(u,A)=\widetilde{\mathcal{T}}^{\varepsilon}(u_0,A_0)$$

for some gauge transformation g and some pair (u_0,A_0) in the image of the map $\mathcal{F}^0$ of Theorem 2.1. Moreover, it is shown that the distances of g to $\mathbb{1}$ (in the $(2,p,\varepsilon)$-norm) and of (u_0,A_0) to $(\bar{u}_0,\bar{A}_0)$ (in any norm) are controlled by the $(1,p,\varepsilon)$-norm of $\bar{\zeta}$.

Theorem C strengthens the local uniqueness result of Theorem B (b) in that condition (10) is no longer required. The proof relies on an ε-dependent local slice theorem (Section 7) and on the construction of a tubular neighbourhood of the moduli space $\mathcal{M}^0_{B,\Sigma}(c_0)$ in which the normal bundle is the intersection of the ε-dependent local slice with the image of the adjoint operator $\mathcal{D}^{\varepsilon *}$ (Section 8).

The next theorem strengthens the local surjectivity result of Theorem C. It does not require the solution (u, A) of (7) to be close to any solution of (8). However, it only applies to solutions that satisfy a uniform L^∞-bound on $d_A u$ and for which u takes values in the compact set

$$M^C := \{x \in M \,|\, |\mu(x)| \le C,\ |\eta| \le C|L_x\eta|\ \forall\ \eta \in \mathfrak{g}\}\,.$$

Theorem D. *Assume* $(H1)$ *and* $(H4)$ *and let* $\bar{B} \in H_2(\bar{M};\mathbb{Z})$ *be a nontorsion homology class. Then, for every* $C > 0$*, there exist positive constants* ε_0 *and* c_1 *such that the following holds for every* $\varepsilon \in (0, \varepsilon_0]$*. If* $(u, A) \in \widetilde{\mathcal{M}}^{\varepsilon}_{B,\Sigma}$ *such that*

$$\|d_A u\|_{L^\infty} \le C, \qquad u(P) \subset M^C \tag{11}$$

then $(u, A) \in \widetilde{\mathcal{T}}^{\varepsilon}(\widetilde{\mathcal{M}}^0_{B,\Sigma}(c_1))$.

Under hypotheses $(H1-2)$ the moduli space $\mathcal{M}^\varepsilon_{B,\Sigma}$ is compact [CGMS]. In this case all solutions of (7) satisfy

$$\|d_A u\|_{L^\infty} + \|\mu(u)\|_{L^\infty} \le C_\varepsilon$$

for some ε-dependent constant C_ε. However, this does not guarantee surjectivity because, on the one hand, the constant C_ε may diverge to infinity as ε tends to zero and, on the other hand, the solutions of (7) may not all satisfy the second condition in the definiton of M^C, namely that the image of u belongs to the set of regular points of μ. There may be sequences $(\varepsilon_i, u_i, A_i)$ of solutions of (7) with $\varepsilon_i \to 0$ such that either $u_i(P)$ intersects the set of singular points of μ or $d_{A_i}u_i$ does not stay bounded, and then bubbling occurs in the small ε limit. Under the hypotheses of Theorem A we shall prove that such bubbling cannot occur and establish a bijection between suitable zero dimensional moduli spaces.

4 Linear estimates

The estimates in this section follow the ones in [DS2, Section 4]. In adapting the proofs to the present context we encounter additional zeroth order terms.

These arise from the Levi-Civita connection and the almost complex structure on M; they are not present in [DS2] where M is replaced by the space of connections over a Riemann surface and the almost complex structure by the Hodge $*$-operator. We extend the results of [DS2] by including estimates for the second derivatives. Moreover, in the present case it is crucial that the constants depend continuously on the pair (u, A). In [DS2] the moduli space is a finite set and so the question of continuous dependence does not arise.

For $u \in C^\infty_{\mathrm{G}}(P, M)$ we introduce the spaces

$$\begin{aligned}\mathcal{X}_u &:= \Omega^0(\Sigma, u^*TM/\mathrm{G}) \times \Omega^1(\Sigma, \mathfrak{g}_P),\\ \mathcal{X}'_u &:= \Omega^{0,1}(\Sigma, u^*TM/\mathrm{G}) \times \Omega^0(\Sigma, \mathfrak{g}_P) \times \Omega^0(\Sigma, \mathfrak{g}_P).\end{aligned}$$

Thus $\mathcal{X}_u = T_{(u,A)}\mathcal{B}$ and $\mathcal{X}'_u = \mathcal{E}_{(u,A)} \times \Omega^0(\Sigma, \mathfrak{g}_P)$ for every $A \in \mathcal{A}(P)$. If the map u is understood from the context then we shall omit the subscript u. It is convenient to introduce the norms

$$\begin{aligned}\|\xi\|_{1,p,\varepsilon;(u,A)} &:= \|\xi\|_{L^p} + \varepsilon \left\|\tilde{\nabla}_A \xi\right\|_{L^p},\\ \|\xi\|_{2,p,\varepsilon;(u,A)} &:= \|\xi\|_{L^p} + \varepsilon \left\|\tilde{\nabla}_A \xi\right\|_{L^p} + \varepsilon^2 \left\|\tilde{\nabla}_A{}^* \tilde{\nabla}_A \xi\right\|_{L^p},\\ \|\alpha\|_{1,p,\varepsilon;A} &:= \|\alpha\|_{L^p} + \varepsilon \|d_A\alpha\|_{L^p} + \varepsilon \|d_A^*\alpha\|_{L^p},\\ \|\alpha\|_{2,p,\varepsilon;A} &:= \|\alpha\|_{L^p} + \varepsilon \|d_A\alpha\|_{L^p} + \varepsilon \|d_A^*\alpha\|_{L^p} + \varepsilon^2 \|d_A^*d_A\alpha + d_Ad_A^*\alpha\|_{L^p}\end{aligned}$$

for $\xi \in \Omega^{0,k}(\Sigma, u^*TM/\mathrm{G})$, $\alpha \in \Omega^k(\Sigma, \mathfrak{g}_P)$, $k = 0, 1$, and $1 \le p \le \infty$. Here $\tilde{\nabla}_A$ denotes the Hermitian connection on u^*TM/G defined by (108). For $\ell = 0, 1, 2$, $1 \le p \le \infty$, and

$$\zeta = (\xi, \alpha) \in \mathcal{X}_u, \qquad \zeta' = (\xi', \varphi', \psi') \in \mathcal{X}'_u$$

we consider the norms

$$\begin{aligned}\|\zeta\|_{\ell,p,\varepsilon;(u,A)} &:= \|\xi\|_{\ell,p,\varepsilon;(u,A)} + \varepsilon \|\alpha\|_{\ell,p,\varepsilon;(u,A)},\\ \|\zeta'\|_{\ell,p,\varepsilon;(u,A)} &:= \|\xi'\|_{\ell,p,\varepsilon;(u,A)} + \varepsilon \|\varphi'\|_{\ell,p,\varepsilon;(u,A)} + \varepsilon \|\psi'\|_{\ell,p,\varepsilon;(u,A)},\end{aligned}$$

where $\|\xi\|_{0,p,\varepsilon;(u,A)} := \|\xi\|_{L^p}$. These norms are gauge invariant, e.g.

$$\left\|(g^{-1}\xi, g^{-1}\alpha g)\right\|_{\ell,p,\varepsilon;(g^{-1}u,g^*A)} = \|(\xi, \alpha)\|_{\ell,p,\varepsilon;(u,A)}.$$

If the pair (u, A) is understood from the context we shall drop it to simplify the notation. In particular, we abbreviate

$$\|\zeta\|_{\infty,\varepsilon} := \|\zeta\|_{0,\infty,\varepsilon;(u,A)}.$$

The augmented vertical differential of $\mathcal{F}^\varepsilon$ at a zero $(u, A) \in \mathcal{B}$ is the operator

$$\mathcal{D}^\varepsilon = \mathcal{D}^\varepsilon_{(u,A)} : \mathcal{X}_u \to \mathcal{X}'_u$$

given by

$$\mathcal{D}^\varepsilon \begin{pmatrix} \xi \\ \alpha \end{pmatrix} = \begin{pmatrix} D\xi + (L_u\alpha)^{0,1} \\ \varepsilon^{-2}L_u^*\xi - d_A^*\alpha \\ \varepsilon^{-2}d\mu(u)\xi + *d_A\alpha \end{pmatrix}, \tag{12}$$

where $D = D_{(u,A)} : \Omega^0(\Sigma, u^*TM/\mathrm{G}) \to \Omega^{0,1}(\Sigma, u^*TM/\mathrm{G})$ is the Cauchy–Riemann operator defined by (109). The second coordinate in the definition of $\mathcal{D}^\varepsilon$ corresponds to the local slice condition for the $\mathcal{G}$-action. For the definition of the adjoint operator it is convenient to use the ε-dependent inner products associated to the $(0, 2, \varepsilon)$-norms. In addition we use twice the standard inner product on the space $\Omega^{0,1}(\Sigma, u^*TM/\mathrm{G})$. Then the adjoint of $\mathcal{D}^\varepsilon$ is given by

$$\mathcal{D}^{\varepsilon *} \begin{pmatrix} \xi' \\ \varphi' \\ \psi' \end{pmatrix} = \begin{pmatrix} 2D^*\xi' + L_u\varphi' + JL_u\psi' \\ 2\varepsilon^{-2}L_u^*\xi' - d_A\varphi' - *d_A\psi' \end{pmatrix}$$

for $(\xi', \varphi', \psi') \in \mathcal{X}'$. The sole purpose of the factor 2 is to render the off-diagonal terms in the operator $\mathcal{D}^{\varepsilon *}\mathcal{D}^\varepsilon$ of zeroth order.

Remark 4.1 (Local coordinates). Let $v : U \to \Sigma$ be a holomorphic coordinate chart defined on an open set $U \subset \mathbb{C}$ and let $\tilde{v} : U \to P$ be a lift of v. In this trivialization the map u, the connection A, the vector field ξ along u, and the 1-form α are represented by

$$\begin{aligned} u^{\mathrm{loc}} &:= u \circ \tilde{v}, & A^{\mathrm{loc}} &:= \tilde{v}^{\,*}A = \Phi\, ds + \Psi\, dt, \\ \xi^{\mathrm{loc}} &:= \xi \circ \tilde{v}, & \alpha^{\mathrm{loc}} &:= \tilde{v}^{\,*}\alpha = \varphi\, ds + \psi\, dt, \end{aligned}$$

where $\Phi, \Psi, \varphi, \psi$ are Lie algebra valued functions on U. The volume form on U is given by

$$\lambda^2\, ds \wedge dt := v^*\mathrm{dvol}_\Sigma$$

for some function $\lambda : U \to (0, \infty)$ and the metric has the form $\lambda^2(ds^2 + dt^2)$. From now on we shall drop the superscript "loc" and introduce the notation

$$\begin{aligned} v_s &:= \partial_s u + X_\Phi(u), & v_t &:= \partial_t u + X_\Psi(u), \\ \nabla_{A,s}\xi &:= \nabla_s\xi + \nabla_\xi X_\Phi(u), & \nabla_{A,t}\xi &:= \nabla_t\xi + \nabla_\xi X_\Psi(u), \\ \tilde{\nabla}_{A,s}\xi &:= \nabla_{A,s}\xi - \tfrac{1}{2}J(\nabla_{v_s}J + \partial_s J)\xi, & \tilde{\nabla}_{A,t}\xi &:= \nabla_{A,t}\xi - \tfrac{1}{2}J(\nabla_{v_t}J + \partial_t J)\xi, \\ \nabla_{A,s}\eta &:= \partial_s\eta + [\Phi, \eta], & \nabla_{A,t}\eta &:= \partial_t\eta + [\Psi, \eta], \end{aligned}$$

for $\eta : U \to \mathfrak{g}$ and a vector field $\xi : U \to u^*TM$ along u. Then

$$d_A u = v_s\, ds + v_t\, dt, \qquad \nabla_A \xi = \nabla_{A,s}\xi\, ds + \nabla_{A,t}\xi\, dt,$$

and

$$\begin{aligned} *F_A &= \lambda^{-2}\left(\partial_s\Psi - \partial_t\Phi + [\Phi,\Psi]\right), \\ *d_A\alpha &= \lambda^{-2}\left(\nabla_{A,s}\psi - \nabla_{A,t}\varphi\right), \\ d_A^*\alpha &= -\lambda^{-2}(\nabla_{A,s}\varphi + \nabla_{A,t}\psi). \end{aligned}$$

In local coordinates a $(0,1)$-form on Σ with values in u^*TM/G has the form $\frac{1}{2}(\xi' ds - J\xi' dt)$, where $\xi'(s,t) \in T_{u(s,t)}M$. In particular,

$$\begin{aligned} \bar\partial_{J,A}(u) &= \frac{1}{2}(v_s + Jv_t)\, ds + \frac{1}{2}(v_t - Jv_s)\, dt, \\ (\nabla_A\xi)^{0,1} &= \frac{1}{2}\left(\nabla_{A,s}\xi + J\nabla_{A,t}\xi\right) ds + \frac{1}{2}\left(\nabla_{A,t}\xi - J\nabla_{A,s}\xi\right) dt, \end{aligned}$$

We represent a $(0,1)$-form by twice the coefficient of ds. Then

$$D\xi = \tilde\nabla_{A,s}\xi + J\tilde\nabla_{A,t}\xi + \frac{1}{4}N(\xi, v_s - Jv_t) + \frac{1}{2}(J\partial_s J - \partial_t J)\xi,$$

where $N(\xi_1,\xi_2) = 2J((\nabla_{\xi_2}J)\xi_1 - (\nabla_{\xi_1}J)\xi_2)$ denotes the Nijenhuis tensor, and

$$2D^*\xi' = \frac{1}{\lambda^2}\left(-\tilde\nabla_{A,s}\xi' + J\tilde\nabla_{A,t}\xi' - \frac{1}{2}J(\nabla_{\xi'}J)(v_s - Jv_t) + \frac{1}{2}(J\partial_s J - \partial_t J)\xi\right).$$

The Weitzenböck formula has the form $D^*D\xi = \frac{1}{2}\tilde\nabla_A{}^*\tilde\nabla_A\xi + \text{l.o.t.}$ In the Kähler case we have $\nabla_A = \tilde\nabla_A$, $\nabla J = 0$, and $\partial_s J = \partial_t J = 0$. Hence in this case

$$\begin{aligned} D^*D\xi &= -\frac{1}{2\lambda^2}\left(\nabla_{A,s}\nabla_{A,s}\xi + \nabla_{A,t}\nabla_{A,t}\xi\right) \\ &\quad -\frac{1}{2\lambda^2}JR(v_s,v_t)\xi - \frac{1}{2\lambda^2}J\nabla_\xi X_{\partial_s\Psi - \partial_t\Phi + [\Phi,\Psi]}(u). \end{aligned}$$

In local coordinates the operators $\mathcal{D}^\varepsilon$ and $\mathcal{D}^{\varepsilon *}$ have the form

$$\mathcal{D}^\varepsilon\zeta = \begin{pmatrix} D\xi + L_u\varphi + JL_u\psi \\ \lambda^{-2}(\nabla_{A,s}\varphi + \nabla_{A,t}\psi) + \varepsilon^{-2}L_u^*\xi \\ \lambda^{-2}(\nabla_{A,s}\psi - \nabla_{A,t}\varphi) + \varepsilon^{-2}d\mu(u)\xi \end{pmatrix},$$

$$\mathcal{D}^{\varepsilon *}\zeta' = \begin{pmatrix} 2D^* + L_u\varphi' + JL_u\psi' \\ -\nabla_{A,s}\varphi' + \nabla_{A,t}\psi' + \varepsilon^{-2}L_u^*\xi' \\ -\nabla_{A,s}\psi' - \nabla_{A,t}\varphi' + \varepsilon^{-2}d\mu(u)\xi' \end{pmatrix}.$$

Proposition 4.2. *For every $p \geq 2$ and every $c_0 > 0$ there exist positive constants ε_0 and c such that*

$$\begin{aligned}
\|\zeta\|_{1,p,\varepsilon;(u,A)} &\leq c\left(\varepsilon\|\mathcal{D}^\varepsilon\zeta\|_{0,p,\varepsilon} + \|\pi_u\xi\|_{L^p}\right), \\
\|\zeta - \pi_u\zeta\|_{1,p,\varepsilon;(u,A)} &\leq c\varepsilon\left(\|\mathcal{D}^\varepsilon\zeta\|_{0,p,\varepsilon} + \|\pi_u\xi\|_{L^p}\right), \\
\|\zeta'\|_{1,p,\varepsilon;(u,A)} &\leq c\left(\varepsilon\|\mathcal{D}^{\varepsilon*}\zeta'\|_{0,p,\varepsilon} + \|\pi_u\xi'\|_{L^p}\right), \\
\|\zeta' - \pi_u\zeta'\|_{1,p,\varepsilon;(u,A)} &\leq c\varepsilon\left(\|\mathcal{D}^{\varepsilon*}\zeta'\|_{0,p,\varepsilon} + \|\pi_u\xi'\|_{L^p}\right),
\end{aligned}$$

for all $(u,A) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$, $\zeta = (\xi,\alpha) \in \mathcal{X}_u$, $\zeta' = (\xi',\varphi',\psi') \in \mathcal{X}'_u$, and $\varepsilon \in (0,\varepsilon_0]$. Here we abbreviate $\mathcal{D}^\varepsilon := \mathcal{D}^\varepsilon_{(u,A)}$ and $\pi_u\zeta := (\pi_u\xi, 0)$ and $\pi_u\zeta' := (\pi_u\xi', 0, 0)$, where π_u is defined by (9).

In this paper we prove Proposition 4.2 only in the case $p = 2$. The proof for $p > 2$ is similar to the proof of an analogous result in [S]. The details for the present case will be carried out elsewhere.

Lemma 4.3. **(i)** *If $\bar\partial_{J,A}(u) = 0$ then*

$$\mathcal{D}^\varepsilon\mathcal{D}^{\varepsilon*}\zeta' = \begin{pmatrix} 2DD^*\xi' + 2\varepsilon^{-2}(L_uL_u^*\xi')^{0,1} + (DJ - JD)L_u\psi' \\ \Delta_\varepsilon\varphi' + [*F_A + \varepsilon^{-2}\mu(u), \psi'] \\ \Delta_\varepsilon\psi' - [*F_A + \varepsilon^{-2}\mu(u), \varphi'] + 2\varepsilon^{-2}L_u^*(DJ - JD)^*\xi' \end{pmatrix}$$

for $\zeta' := (\xi',\varphi',\psi') \in \mathcal{X}'_u$, where $\Delta_\varepsilon := d_A^*d_A + \varepsilon^{-2}L_u^*L_u$.
(ii) *If $\bar\partial_{J,A}(u) = 0$ and $\mu(u) = 0$ then*

$$\mathcal{D}^{\varepsilon*}\mathcal{D}^\varepsilon\zeta = \begin{pmatrix} 2D^*D\xi + \varepsilon^{-2}L_uL_u^*\xi + \varepsilon^{-2}JL_uL_u^*J^*\xi + Q^*\alpha \\ d_A^*d_A\alpha + d_Ad_A^*\alpha + \varepsilon^{-2}L_u^*L_u\alpha + \varepsilon^{-2}Q\xi \end{pmatrix},$$

for $\zeta = (\xi,\alpha) \in \mathcal{X}_u$, where $Q : \Omega^0(\Sigma, u^*TM/\mathrm{G}) \to \Omega^1(\Sigma, \mathfrak{g}_P)$ denotes the zeroth order operator

$$\begin{aligned}
Q\xi &:= 2L_u^*D\xi - d_AL_u^*\xi - *d_Ad\mu(u)\xi \\
&= \rho(\xi, d_Au) - *\rho(J\xi, d_Au) + *L_u^*\dot J\xi + \frac{1}{2}L_u^*N(\xi, \partial_{J,A}(u)).
\end{aligned}$$

Proof. We shall repeatedly use the identities

$$\begin{aligned}
d_A^*\alpha &= -*d_A*\alpha, & *d_Ad_A\varphi &= [*F_A, \varphi], \\
L_u^* &= d\mu(u)J, & d\mu(u)L_u\varphi &= -[\mu(u), \varphi]
\end{aligned}$$

for $\alpha \in \Omega^1(\Sigma, \mathfrak{g}_P)$ and $\varphi \in \Omega^0(\Sigma, \mathfrak{g}_P)$. To prove (i) note that the triple $(\tilde{\xi}', \tilde{\varphi}', \tilde{\psi}') := \mathcal{D}^\varepsilon \mathcal{D}^{\varepsilon *}(\xi', \varphi', \psi')$ is given by

$$\begin{array}{rcl}
\tilde{\xi}' & = & D(2D^*\xi' + L_u\varphi' + JL_u\psi') + (L_u(2\varepsilon^{-2}L_u^*\xi' - d_A\varphi' - *d_A\psi'))^{0,1} \\
& = & 2DD^*\xi' + 2\varepsilon^{-2}(L_uL_u^*\xi')^{0,1} \\
& & + DL_u\varphi' - (L_ud_A\varphi')^{0,1} + DJL_u\psi' - (L_u * d_A\psi')^{0,1}, \\
\tilde{\varphi}' & = & \varepsilon^{-2}L_u^*(2D^*\xi' + L_u\varphi + JL_u\psi) - d_A^*(2\varepsilon^{-2}L_u^*\xi' - d_A\varphi - *d_A\psi) \\
& = & d_A^*d_A\varphi' + \varepsilon^{-2}L_u^*L_u\varphi' + 2\varepsilon^{-2}(DL_u - L_ud_A)^*\xi' + [*F_A + \varepsilon^{-2}\mu(u), \psi'], \\
\tilde{\psi}' & = & \varepsilon^{-2}d\mu(u)(2D^*\xi' + L_u\varphi' + JL_u\psi') + *d_A(2\varepsilon^{-2}L_u^*\xi' - d_A\varphi' - *d_A\psi') \\
& = & d_A^*d_A\psi' + \varepsilon^{-2}L_u^*L_u\psi' - [*F_A + \varepsilon^{-2}\mu(u), \varphi'] \\
& & + 2\varepsilon^{-2}(L_u^*J^*D^*\xi' + d_A^* * L_u^*\xi').
\end{array}$$

The assertion now follows from the fact that

$$J(L_u\alpha)^{0,1} = (L_u * \alpha)^{0,1}, \qquad L_u^*J^*\xi' = - * L_u^*\xi', \tag{13}$$

for $\alpha \in \Omega^1(\Sigma, \mathfrak{g}_P)$ and $\xi' \in \Omega^{0,1}(\Sigma, u^*TM/\mathrm{G})$, and

$$\bar{\partial}_{J,A}(u) = 0 \qquad \Longrightarrow \qquad DL_u\varphi = (L_ud_A\varphi)^{0,1} \tag{14}$$

for $\varphi \in \Omega^0(\Sigma, \mathfrak{g}_P)$. The first equation in (13) follows from the fact that $*\alpha = -\alpha \circ J_\Sigma$ for every 1-form α on Σ (with values in any vector bundle). The second equation in (13) follows from the first by duality. Equation (14) follows from the fact that the section $(u, A) \mapsto \bar{\partial}_{J,A}(u)$ of the vector bundle over $\mathcal{B}$ with fibres $\Omega^{0,1}(\Sigma, u^*TM/\mathrm{G})$ is $\mathcal{G}(P)$-equivariant.

To prove (ii) note that the pair $(\tilde{\xi}, \tilde{\alpha}) := \mathcal{D}^{\varepsilon *}\mathcal{D}^\varepsilon(\xi, \alpha)$ is given by

$$\begin{array}{rcl}
\tilde{\xi} & = & 2D^*(D\xi + (L_u\alpha)^{0,1}) \\
& & + L_u(\varepsilon^{-2}L_u^*\xi - d_A^*\alpha) + JL_u(\varepsilon^{-2}d\mu(u)\xi + *d_A\alpha) \\
& = & 2D^*D\xi + 2\varepsilon^{-2}(L_uL_u^*\xi')^{0,1} + \left(2D^*(L_u\alpha)^{0,1} - L_ud_A^*\alpha + JL_u * d_A\alpha\right), \\
\tilde{\alpha} & = & 2\varepsilon^{-2}L_u^*(D\xi + (L_u\alpha)^{0,1}) \\
& & - d_A(\varepsilon^{-2}L_u^*\xi - d_A^*\alpha) - *d_A(\varepsilon^{-2}d\mu(u)\xi + *d_A\alpha) \\
& = & d_A^*d_A\alpha + d_Ad_A^*\alpha + \varepsilon^{-2}L_u^*L_u\alpha + \varepsilon^{-2}\left(2L_u^*D\xi - d_AL_u^*\xi - *d_Ad\mu(u)\xi\right).
\end{array}$$

Here we have used the fact that $\mu(u) = 0$ and hence $2L_u^*(L_u\alpha)^{0,1} = L_u^*L_u\alpha$. The formula for the operator $Q := 2L_u^*D - d_AL_u^* - *d_Ad\mu(u)$ follows by computing in local coordinates. □

Proof of Proposition 4.2 for $p=2$. Let $\zeta' := (\xi', \varphi', \psi') := \mathcal{D}^\varepsilon \zeta$. Then, by Lemma 4.3, the formula

$$\mathcal{D}^{\varepsilon *}\mathcal{D}^\varepsilon \zeta = \mathcal{D}^{\varepsilon *}\zeta'$$

is equivalent to

$$\begin{aligned} 2D^*D\xi + \varepsilon^{-2}L_uL_u^*\xi + \varepsilon^{-2}JL_uL_u^*J^*\xi + Q^*\alpha &= 2D^*\xi' + L_u\varphi' + JL_u\psi', \\ d_A^*d_A\alpha + d_Ad_A^*\alpha + \varepsilon^{-2}L_u^*L_u\alpha + \varepsilon^{-2}Q\xi &= 2\varepsilon^{-2}L_u^*\xi' - d_A\varphi' - *d_A\psi'. \end{aligned}$$

Take the L^2-inner product of the first equation with ξ and of the second equation with $\varepsilon^2\alpha$. The sum of the resulting identities gives

$$\begin{aligned} &\varepsilon^{-2}\left\|L_u^*\xi\right\|^2 + \varepsilon^{-2}\left\|L_u^*J\xi\right\|^2 + 2\left\|D\xi\right\|^2 + \left\|L_u\alpha\right\|^2 + \varepsilon^2\left\|d_A\alpha\right\|^2 + \varepsilon^2\left\|d_A^*\alpha\right\|^2 \\ &\quad = 2\langle \xi', D\xi\rangle + 2\langle \xi', L_u\alpha\rangle - 2\langle \alpha, Q\xi\rangle \\ &\qquad + \langle \varphi', L_u^*\xi\rangle - \langle \psi', L_u^*J\xi\rangle - \varepsilon^2\langle \varphi', d_A^*\alpha\rangle + \varepsilon^2\langle \psi', *d_A\alpha\rangle \\ &\quad \le 3\left\|\xi'\right\|^2 + \left\|D\xi\right\|^2 + 2^{-1}\left\|L_u\alpha\right\|^2 + \delta\left\|\alpha\right\|^2 + \delta^{-1}\left\|Q\xi\right\|^2 \\ &\qquad + \varepsilon^2\left\|\varphi'\right\|^2 + \varepsilon^2\left\|\psi'\right\|^2 + 2^{-1}\varepsilon^{-2}\left\|L_u^*\xi\right\|^2 + 2^{-1}\varepsilon^{-2}\left\|L_u^*J\xi\right\|^2 \\ &\qquad + 2^{-1}\varepsilon^2\left\|d_A^*\alpha\right\|^2 + 2^{-1}\varepsilon^2\left\|d_A\alpha\right\|^2. \end{aligned}$$

Here all norms are L^2-norms and all inner products are L^2-inner products. Choose $\delta > 0$ so small that $\delta\left\|\alpha\right\|^2 \le 4^{-1}\left\|L_u\alpha\right\|^2$ for all α. Then

$$\begin{aligned} &\varepsilon^{-2}\left\|L_u^*\xi\right\|^2 + \varepsilon^{-2}\left\|L_u^*J\xi\right\|^2 + \left\|D\xi\right\|^2 + \left\|L_u\alpha\right\|^2 + \varepsilon^2\left\|d_A\alpha\right\|^2 + \varepsilon^2\left\|d_A^*\alpha\right\|^2 \\ &\quad \le 12\left\|\xi'\right\|^2 + 4\varepsilon^2\left\|\varphi'\right\|^2 + 4\varepsilon^2\left\|\psi'\right\|^2 + 4\delta^{-1}\left\|Q\right\|_{L^\infty}^2\left\|\xi\right\|^2 \\ &\quad \le 12\left\|\mathcal{D}^\varepsilon(\xi,\alpha)\right\|_{0,2,\varepsilon}^2 + 4\delta^{-1}\left\|Q\right\|_{L^\infty}^2\left\|\xi\right\|_{L^2}^2. \end{aligned}$$

Now the required estimates follow from the inequalities

$$\begin{aligned} \|\tilde{\nabla}_A\pi_u\xi\|_{L^2} &\le c\left(\|\tilde{\nabla}_A\xi\|_{L^2} + \|\xi\|_{L^2}\right), \\ \|\xi - \pi_u\xi\|_{L^2} &\le c'\left(\|L_u^*\xi\|_{L^2} + \|L_u^*J\xi\|_{L^2}\right), \\ \|\tilde{\nabla}_A\xi\|_{L^2} &\le c''\left(\|D\xi\|_{L^2} + \|\xi\|_{L^2}\right). \end{aligned}$$

The first inequality follows from (15) below. In the second inequality the constant c' can be chosen as an upper bound for the norms of the linear maps $L_x(L_x^*L_x)^{-1}$ over all $x \in \mu^{-1}(0)$. The third inequality is the L^2-estimate for the Cauchy–Riemann operator and it follows from the Weitzenböck formula. The constant c'' is gauge invariant and depends continuously on the pair (u, A) with respect to the C^1-norm and hence can be chosen independent of $(u, A) \in \widetilde{\mathcal{M}}_{B,\Sigma}^0(c_0)$. This proves the proposition in the case $p = 2$. □

The next lemma expresses the Sobolev inequalities in terms of the ε-dependent norms.

Lemma 4.4. *For every $p > 2$ and every $c_0 > 0$ there exists a constant $c > 0$ such that*

$$\|\zeta\|_{\infty,\varepsilon} \le c\varepsilon^{-2/p}\|\zeta\|_{1,p,\varepsilon;(u,A)}, \qquad \|\zeta\|_{1,\infty,\varepsilon} \le c\varepsilon^{-2/p}\|\zeta\|_{2,p,\varepsilon;(u,A)}$$

for all $(u, A) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$, $\zeta \in \mathcal{X}_u$, and $\varepsilon \in (0, 1)$.

Proof. Multiply the metric on Σ by ε^{-2}. Then the $W^{k,p}$-norm of (ξ, α) with respect to the rescaled metric is equal to $\varepsilon^{-2/p}$ times the (k, p, ε)-norm of ζ, and the L^∞-norm with respect to the rescaled metric is equal to the (∞, ε)-norm. Hence the estimates follows from the Sobolev embedding theorem for the rescaled metric. The constant is gauge invariant and it depends continuously on u (with respect to the C^1-norm) and A (with respect to the C^0-norm). By Lemma 2.2, the estimate holds with a uniform constant c. $\square$

Lemma 4.5. *For every $p \ge 2$ and every $c_0 > 0$ there exist positive constants ε_0 and c such that*

$$\begin{aligned}
\|\pi_u\mathcal{D}^\varepsilon\zeta - \mathcal{D}^0\pi_u\zeta\|_{k,p,\varepsilon} &\le c\|\xi - \pi_u\xi\|_{k,p,\varepsilon}, \\
\|\pi_u\mathcal{D}^\varepsilon\zeta - \mathcal{D}^\varepsilon\pi_u\zeta\|_{k,p,\varepsilon} &\le c\|\xi\|_{k,p,\varepsilon}, \\
\|\pi_u\mathcal{D}^{\varepsilon*}\zeta' - \mathcal{D}^{0*}\pi_u\zeta'\|_{k,p,\varepsilon} &\le c\|\xi' - \pi_u\xi'\|_{k,p,\varepsilon}, \\
\|\pi_u\mathcal{D}^{\varepsilon*}\zeta' - \mathcal{D}^{\varepsilon*}\pi_u\zeta'\|_{k,p,\varepsilon} &\le c\|\xi'\|_{k,p,\varepsilon}
\end{aligned}$$

for every $(u, A) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$, $\zeta = (\xi, \alpha) \in \mathcal{X}_u$, $\zeta' = (\xi', \varphi', \psi') \in \mathcal{X}'_u$, $\varepsilon \in (0, 1]$, and $k = 0, 1$. Here we abbreviate $\mathcal{D}^\varepsilon := \mathcal{D}^\varepsilon_{(u,A)}$ for $\varepsilon \ge 0$.

Proof. We prove first that, for every vector field $v \in \mathrm{Vect}(\Sigma)$, there exists a constant $c = c(p, c_0, v) > 0$ such that

$$\begin{aligned}
\|\pi_u\tilde{\nabla}_{A,v}\xi - \tilde{\nabla}_{A,v}\pi_u\xi\|_{W^{k,p},A} &\le c\|\xi\|_{W^{k,p},A}, \\
\|\pi_u\tilde{\nabla}_{A,v}\xi - \pi_u\tilde{\nabla}_{A,v}\pi_u\xi\|_{W^{k,p},A} &\le c\|\xi - \pi_u\xi\|_{W^{k,p},A},
\end{aligned} \tag{15}$$

for $(u, A) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$, $\xi \in \Omega^0(\Sigma, u^*TM/\mathrm{G})$ and $k = 0, 1$. Here the $W^{1,p}$-norm labelled by A is understood as the (gauge invariant) $(1, p, \varepsilon)$-norm for $\varepsilon = 1$. To prove (15) we choose local holomorphic coordinates $s + it$ on Σ.

Thus $\xi(s,t) \in T_{(u(s,t)}M$, and v_s, v_t, $\nabla_{A,s}\xi$, and $\nabla_{A,t}\xi$ are as in Remark 4.1. Write

$$\xi = \pi_u\xi + L_u\eta_1 + JL_u\eta_2,$$

where $\eta_i(s,t) \in \mathfrak{g}$. Define $B_s(s,t) : \mathfrak{g} \to T_{u(s,t)}M$ and $B_t(s,t) : \mathfrak{g} \to T_{u(s,t)}M$ by

$$B_s\eta := \nabla_{v_s}X_\eta(u), \qquad B_t\eta := \nabla_{v_t}X_\eta(u).$$

Then

$$\nabla_{A,s}L_u\eta - L_u\nabla_{A,s}\eta = B_s\eta, \qquad \nabla_{A,t}L_u\eta - L_u\nabla_{A,t}\eta = B_t\eta$$

and hence

$$\nabla_{A,s}\pi_u\xi - \pi_u\nabla_{A,s}\xi = \pi_u(B_s\eta_1 + JB_s\eta_2 + (\nabla_{v_s}J + \partial_s J)L_u\eta_2).$$

Since $\eta_1 = (L_u^*L_u)^{-1}L_u^*(\xi - \pi_u\xi)$ and $\eta_2 = -(L_u^*L_u)^{-1}L_u^*J(\xi - \pi_u\xi)$, we have

$$\|\nabla_{A,s}\pi_u\xi - \pi_u\nabla_{A,s}\xi\|_{L^p} \le c\|\xi - \pi_u\xi\|_{L^p}.$$

This proves (15) for the local vector field $\partial/\partial s$. For $\partial/\partial t$ the proof is analogous. Hence the result follows for any linear combination of these vector fields supported in the given coordinate chart, and hence for every vector field on Σ. For $\xi' \in \Omega^{0,1}(\Sigma, u^*TM/\mathrm{G})$ there are similar inequalities.

By (15), there exists a constant $c' = c'(p, c_0) > 0$ such that

$$\begin{aligned}
\|\pi_u(\tilde{\nabla}_A\xi)^{0,1} - (\tilde{\nabla}_{A,v}\pi_u\xi)^{0,1}\|_{W^{k,p},A} &\le c'\|\xi\|_{W^{k,p},A}, \\
\|\pi_u(\tilde{\nabla}_A\xi)^{0,1} - \pi_u(\tilde{\nabla}_{A,v}\pi_u\xi)^{0,1}\|_{W^{k,p},A} &\le c'\|\xi - \pi_u\xi\|_{W^{k,p},A},
\end{aligned}$$

for $(u,A) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$, $\xi \in \Omega^0(\Sigma, u^*TM/\mathrm{G})$, and $k = 0, 1$. Now the operator $D = D_{(u,A)}$ is given by

$$D\xi = (\nabla_A\xi)^{0,1} - J(\nabla_\xi J)\partial_{J,A}(u)$$

and hence

$$\begin{aligned}
\|\pi_u D\xi - D\pi_u\xi\|_{W^{k,p},A} &\le c''\|\xi\|_{W^{k,p},A}, \\
\|\pi_u D(\xi - \pi_u\xi)\|_{W^{k,p},A} &\le c''\|\xi - \pi_u\xi\|_{W^{k,p},A},
\end{aligned} \tag{16}$$

for $(u,A) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$, $\xi \in \Omega^0(\Sigma, u^*TM/\mathrm{G})$, and $k = 0, 1$. Since

$$\pi_u\mathcal{D}^\varepsilon\zeta = \pi_u D\xi, \qquad \mathcal{D}^0\pi_u\zeta = \pi_u D\pi_u\xi, \qquad \mathcal{D}^\varepsilon\pi_u\zeta = (D\pi_u\xi, 0, 0),$$

the required estimates for the operator $\mathcal{D}^\varepsilon$ follow from (16). The proof for the adjoint operator is analogous. $\square$

In the following we use the notation

$$\tilde{\nabla}_{A,v}\zeta := (\tilde{\nabla}_{A,v}\xi, \nabla_{A,v}\alpha), \qquad \tilde{\nabla}_{A,v}\xi' := (\tilde{\nabla}_{A,v}\xi', \nabla_{A,v}\varphi', \nabla_{A,v}\psi')$$

for $v \in \mathrm{Vect}(\Sigma)$, $\zeta = (\xi, \alpha) \in \mathcal{X}_u$, and $\zeta' = (\xi', \varphi', \psi') \in \mathcal{X}'_u$, where $\tilde{\nabla}_A$ is the connection on u^*TM/G defined by (108) and $\nabla_{A,v}\alpha \in \Omega^1(\Sigma, \mathfrak{g}_P)$ is the covariant derivative induced by the connection A and the Levi-Civita connection on Σ.

Lemma 4.6. *For every $p \geq 2$, every $c_0 > 0$, and every equivariant vector field $v \in \mathrm{Vect}(\Sigma)$ there exists a constant $c > 0$ such that*

$$\begin{aligned} \|\mathcal{D}^\varepsilon \tilde{\nabla}_{A,v}\zeta - \tilde{\nabla}_{A,v}\mathcal{D}^\varepsilon\zeta\|_{k,p,\varepsilon} &\leq c\varepsilon^{-1}\|\zeta\|_{k+1,p,\varepsilon}, && (17)\\ \|\mathcal{D}^{\varepsilon *} \tilde{\nabla}_{A,v}\zeta' - \tilde{\nabla}_{A,v}\mathcal{D}^{\varepsilon *}\zeta'\|_{k,p,\varepsilon} &\leq c\varepsilon^{-1}\|\zeta'\|_{k+1,p,\varepsilon} && (18) \end{aligned}$$

for all $(u, A) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$, $\zeta \in \mathcal{X}_u$, $\zeta' \in \mathcal{X}_u$, $\varepsilon \in (0, 1]$, and $k = 0, 1$.

Proof. We compute in local coordinates. Let $\zeta' = (\xi', \varphi', \psi') := \mathcal{D}^\varepsilon\zeta$. Then

$$\begin{aligned} \xi' &= \tilde{\nabla}_{A,s}\xi + J\tilde{\nabla}_{A,t}\xi + \frac{1}{4}N(\xi, v_s - Jv_t) + \frac{1}{2}(J\partial_s J - \partial_t J)\xi + L_u\varphi + JL_u\psi, \\ \varphi' &= \lambda^{-2}\left(\nabla_{A,s}\varphi + \nabla_{A,t}\psi\right) + \varepsilon^{-2}L_u^*\xi, \qquad (19)\\ \psi' &= \lambda^{-2}\left(\nabla_{A,s}\psi - \nabla_{A,t}\varphi\right) + \varepsilon^{-2}d\mu(u)\xi. \end{aligned}$$

Here $A = \Phi ds + \Psi dt$, $\alpha = \varphi ds + \psi dt$, and v_s, v_t, $\nabla_{A,s}\varphi$, $\tilde{\nabla}_{A,s}\xi$, $\nabla_{A,t}\varphi$, and $\tilde{\nabla}_{A,t}\xi$ are as in Remark 4.1. It suffices to prove the estimate for the local operators $\tilde{\nabla}_{A,s}$ and $\tilde{\nabla}_{A,t}$. Let $\zeta'_s = (\xi'_s, \varphi'_s, \psi'_s)$ be defined by (19) with (ξ, φ, ψ) replaced by $(\tilde{\nabla}_{A,s}\xi, \nabla_{A,s}\varphi, \nabla_{A,s}\psi)$. Since $\tilde{\nabla}_A J = 0$ we obtain

$$\begin{aligned} \tilde{\nabla}_{A,s}\xi' - \xi'_s &= J(\tilde{\nabla}_{A,s}\tilde{\nabla}_{A,t}\xi - \tilde{\nabla}_{A,t}\tilde{\nabla}_{A,s}\xi) + \nabla_{v_s}X_\varphi(u) + J\nabla_{v_s}X_\psi(u) \\ &\quad + \frac{1}{4}\tilde{\nabla}_{A,s}(N(\xi, v_s - Jv_t)) - \frac{1}{4}N(\tilde{\nabla}_{A,s}\xi, v_s - Jv_t) \\ &\quad + \frac{1}{2}\tilde{\nabla}_{A,s}((J\partial_s J - \partial_t J)\xi) - \frac{1}{2}(J\partial_s J - \partial_t J)\tilde{\nabla}_{A,s}\xi \\ &\quad - \frac{1}{2}J(\nabla_{v_s}J + \partial_s J)(L_u\varphi - JL_u\psi), \\ \nabla_{A,s}\varphi' - \varphi'_s &= \lambda^{-2}\left(\nabla_{A,s}\nabla_{A,t}\psi - \nabla_{A,t}\nabla_{A,s}\psi\right) + (\partial_s\lambda^{-2})\left(\nabla_{A,s}\varphi + \nabla_{A,t}\psi\right) \\ &\quad + \varepsilon^{-2}\rho(v_s, \xi) - \frac{1}{2}\varepsilon^{-2}d\mu(u)(\nabla_{v_s}J + \partial_s J)\xi, \\ \nabla_{A,s}\psi' - \psi'_s &= -\lambda^{-2}\left(\nabla_{A,s}\nabla_{A,t}\varphi + \nabla_{A,t}\nabla_{A,s}\varphi\right) + (\partial_s\lambda^{-2})\left(\nabla_{A,s}\psi - \nabla_{A,t}\varphi\right) \\ &\quad - \varepsilon^{-2}\rho(v_s, J\xi) - \frac{1}{2}\varepsilon^{-2}L_u^*(\nabla_{v_s}J + \partial_s J)\xi. \end{aligned}$$

Here we have used Lemma C.2. For the vector field $\partial/\partial s$, multiplied by any cutoff function, the estimates (17) and (18) follow from these three identities. The proof for $\partial/\partial t$ is similar, and so is the proof for the adjoint operator. $\square$

Lemma 4.7. *Let $p \geq 2$ and $c_0 > 0$. Suppose that $\mathcal{D}^0 := \mathcal{D}^0_{(u,A)}$ is onto for every $(u, A) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$. Then there exist positive constants ε_0 and c such that the operator $\mathcal{D}^\varepsilon := \mathcal{D}^\varepsilon_{(u,A)}$ is onto for every $(u, A) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$ and every $\varepsilon \in (0, \varepsilon_0]$ and*

$$\begin{aligned} \|\zeta'\|_{k+1,p,\varepsilon} &\leq c\Big(\varepsilon \|\mathcal{D}^{\varepsilon*}\zeta'\|_{k,p,\varepsilon} + \|\pi_u\mathcal{D}^{\varepsilon*}\zeta'\|_{k,p,\varepsilon}\Big), && (20)\\ \|\zeta' - \pi_u\zeta'\|_{k+1,p,\varepsilon} &\leq c\varepsilon \|\mathcal{D}^{\varepsilon*}\zeta'\|_{k,p,\varepsilon}, && (21)\end{aligned}$$

for $\zeta' \in \mathcal{X}'_u$ and $k = 0, 1$.

Proof. By elliptic regularity, there exists a constant $C_0 > 0$ such that

$$\|\xi_0\|_{L^p} \leq C_0\|\mathcal{D}^{0*}\xi_0\|_{L^p} \tag{22}$$

for every $(u, A) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$ and every $\xi_0' \in \Omega^{0,1}(\Sigma, H_u)$. Hence

$$\begin{aligned} \|\pi_u\zeta'\|_{L^p} &\leq C_0\|\mathcal{D}^{0*}\pi_u\zeta'\|_{L^p}\\ &\leq C_0\|\pi_u\mathcal{D}^{\varepsilon*}\zeta'\|_{L^p} + C_0\|\pi_u\mathcal{D}^{\varepsilon*}\zeta' - \mathcal{D}^{0*}\pi_u\zeta'\|_{L^p}\\ &\leq C_0\|\pi_u\mathcal{D}^{\varepsilon*}\zeta'\|_{L^p} + C_0c_1\|\zeta' - \pi_u\zeta'\|_{0,p,\varepsilon}\\ &\leq C_0\|\pi_u\mathcal{D}^{\varepsilon*}\zeta'\|_{L^p} + C_0c_1c_2\varepsilon\left(\|\mathcal{D}^{\varepsilon*}\zeta'\|_{0,p,\varepsilon} + \|\pi_u\zeta'\|_{L^p}\right).\end{aligned}$$

Here c_1 is the constant of Lemma 4.5 and c_2 is the constant of Proposition 4.2. With $C_0c_1c_2\varepsilon \leq 1/2$ we obtain

$$\|\pi_u\zeta'\|_{L^p} \leq c_3\big(\varepsilon\|\mathcal{D}^{\varepsilon*}\zeta'\|_{0,p,\varepsilon} + \|\pi_u\mathcal{D}^{\varepsilon*}\zeta'\|_{L^p}\big), \tag{23}$$

where $c_3 := 2C_0c_1c_2$. The inequality (20) for $k = 0$ now follows from (23) and Proposition 4.2. To prove (21) for $k = 0$ we use Proposition 4.2 and (23) again to obtain

$$\begin{aligned} \|\zeta' - \pi_u\zeta'\|_{1,p,\varepsilon} &\leq c_2\varepsilon\left(\|\mathcal{D}^{\varepsilon*}\zeta'\|_{0,p,\varepsilon} + \|\pi_u\zeta'\|_{L^p}\right)\\ &\leq c_2\varepsilon(1 + c_3\varepsilon)\|\mathcal{D}^{\varepsilon*}\zeta'\|_{0,p,\varepsilon} + c_2c_3\varepsilon\|\pi_u\mathcal{D}^{\varepsilon*}\zeta'\|_{L^p}\\ &\leq c_4\varepsilon\|\mathcal{D}^{\varepsilon*}\zeta'\|_{0,p,\varepsilon},\end{aligned}$$

where $c_4 := c_2(1+2c_3)$.

Now let $v \in \mathrm{Vect}(\Sigma)$. By definition of the $(1,p,\varepsilon)$-norm and (15), there exists a constant $c_5 = c_5(v,p,c_0)$ such that

$$\|\tilde{\nabla}_{A,v}\zeta\|_{0,p,\varepsilon} \le c_5\varepsilon^{-1}\|\zeta\|_{1,p,\varepsilon}, \qquad \|\pi_u\tilde{\nabla}_{A,v}\zeta - \tilde{\nabla}_{A,v}\pi_u\zeta\|_{L^p} \le c_5\|\zeta\|_{0,p,\varepsilon},$$

for $(u,A) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$, $\zeta \in \mathcal{X}_u$ and $\varepsilon \in (0,1]$. Let $c_6 = c_6(v,p,c_0)$ be the constant of Lemma 4.6. Then, by (20) with $k=0$ and Lemma 4.6, we have

$$\begin{aligned}
\varepsilon\|\tilde{\nabla}_{A,v}\zeta'\|_{1,p,\varepsilon} &\le c\varepsilon\left(\varepsilon\|\mathcal{D}^{\varepsilon*}\tilde{\nabla}_{A,v}\zeta'\|_{0,p,\varepsilon} + \|\pi_u\mathcal{D}^{\varepsilon*}\tilde{\nabla}_{A,v}\zeta'\|_{L^p}\right) \\
&\le c\varepsilon^2\|\mathcal{D}^{\varepsilon*}\tilde{\nabla}_{A,v}\zeta' - \tilde{\nabla}_{A,v}\mathcal{D}^{\varepsilon*}\zeta'\|_{0,p,\varepsilon} + c\varepsilon^2\|\tilde{\nabla}_{A,v}\mathcal{D}^{\varepsilon*}\zeta'\|_{0,p,\varepsilon} \\
&\quad + c\varepsilon\|\pi_u(\mathcal{D}^{\varepsilon*}\tilde{\nabla}_{A,v}\zeta' - \tilde{\nabla}_{A,v}\mathcal{D}^{\varepsilon*}\zeta')\|_{L^p} \\
&\quad + c\varepsilon\|\pi_u\tilde{\nabla}_{A,v}\mathcal{D}^{\varepsilon*}\zeta' - \tilde{\nabla}_{A,v}\pi_u\mathcal{D}^{\varepsilon*}\zeta'\|_{L^p} + c\varepsilon\|\tilde{\nabla}_{A,v}\pi_u\mathcal{D}^{\varepsilon*}\zeta'\|_{L^p} \\
&\le 2cc_5\left(\varepsilon\|\mathcal{D}^{\varepsilon*}\zeta'\|_{1,p,\varepsilon} + \|\pi_u\mathcal{D}^{\varepsilon*}\zeta'\|_{1,p,\varepsilon}\right) + 2cc_6\|\zeta'\|_{1,p,\varepsilon} \\
&\le 2c(c_5+cc_6)\left(\varepsilon\|\mathcal{D}^{\varepsilon*}\zeta'\|_{1,p,\varepsilon} + \|\pi_u\mathcal{D}^{\varepsilon*}\zeta'\|_{1,p,\varepsilon}\right).
\end{aligned}$$

The last inequality follows again from (20) with $k=0$. The estimate (20) for $k=1$ now follows by taking the sum over finitely many suitably chosen vector fields v.

To prove (21) for $k=1$ we observe that $\pi_u\mathcal{D}^{\varepsilon*}\pi_u\zeta' = \mathcal{D}^{0*}\pi_u\zeta'$ and choose c_7 such that $\|\pi_u\zeta\|_{1,p,\varepsilon} \le c_7\|\zeta\|_{1,p,\varepsilon}$ for every $\zeta \in \mathcal{X}_u$. Let c_8 be the constant of Lemma 4.5. Then, by (20) with $k=1$ and Lemma 4.5, we have

$$\begin{aligned}
\|\zeta' - \pi_u\zeta'\|_{2,p,\varepsilon} &\le c\left(\varepsilon\|\mathcal{D}^{\varepsilon*}(\zeta' - \pi_u\zeta')\|_{1,p,\varepsilon} + \|\pi_u\mathcal{D}^{\varepsilon*}(\zeta' - \pi_u\zeta')\|_{1,p,\varepsilon}\right) \\
&\le c\varepsilon\left(\|\mathcal{D}^{\varepsilon*}\zeta'\|_{1,p,\varepsilon} + \|\pi_u\mathcal{D}^{\varepsilon*}\zeta'\|_{1,p,\varepsilon}\right) \\
&\quad + c\varepsilon\|\mathcal{D}^{\varepsilon*}\pi_u\zeta' - \pi_u\mathcal{D}^{\varepsilon*}\zeta'\|_{1,p,\varepsilon} \\
&\quad + c\|\pi_u\mathcal{D}^{\varepsilon*}\zeta' - \mathcal{D}^{0*}\pi_u\zeta'\|_{1,p,\varepsilon} \\
&\le c(1+c_7)\varepsilon\|\mathcal{D}^{\varepsilon*}\zeta'\|_{1,p,\varepsilon} + cc_8\varepsilon\|\zeta'\|_{1,p,\varepsilon} + cc_8\|\zeta' - \pi_u\zeta'\|_{1,p,\varepsilon} \\
&\le c_9\varepsilon\|\mathcal{D}^{\varepsilon*}\zeta'\|_{1,p,\varepsilon}.
\end{aligned}$$

The last inequality follows from (20) and (21) with $k=0$. □

Lemma 4.8. *Let $p \ge 2$ and $c_0 > 0$. Suppose that $\mathcal{D}^0 := \mathcal{D}^0_{(u,A)}$ is onto for every $(u,A) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$. Then there exist positive constants c and ε_0 such that*

$$\|\mathcal{D}^{\varepsilon*}\zeta'\|_{k+1,p,\varepsilon} \le c\Big(\varepsilon\,\|\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon*}\zeta'\|_{k,p,\varepsilon} + \|\pi_u\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon*}\zeta'\|_{k,p,\varepsilon}\Big), \tag{24}$$

$$\|\mathcal{D}^{\varepsilon*}\zeta' - \pi_u\mathcal{D}^{\varepsilon*}\zeta'\|_{k+1,p,\varepsilon} \le c\varepsilon\,\|\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon*}\zeta'\|_{k,p,\varepsilon} \tag{25}$$

for every $(u,A)\in\widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$, $\zeta'\in\mathcal{X}'_u$, $\varepsilon\in(0,\varepsilon_0]$, *and* $k=0,1$.

Proof. The proof has nine steps.

Step 1. *Let* $q>1$ *such that* $1/p+1/q=1$. *Then there exists a constant* $c_0>0$ *such that*

$$\|\xi_0\|_{L^p}+\|\xi_0\|_{L^q}\le c_0\,\|\xi_0\|_{L^2}\,,\qquad \|\xi_0'\|_{L^q}\le c_0\left\|\mathcal{D}^{0*}\xi_0'\right\|_{L^q}$$

for every $(u,A)\in\widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$, *every* $\xi_0\in\ker\mathcal{D}^0$, *and every* $\xi_0'\in\Omega^{0,1}(\Sigma,H_u)$.

These are standard estimates for elliptic pdes. The first estimate uses L^2 regularity for the operator $\mathcal{D}^0$, the Sobolev embedding $W^{1,2}\hookrightarrow L^p$, and the Hölder inequality. The second estimate uses L^q regularity for $\mathcal{D}^{0*}$ and the fact that $\mathcal{D}^{0*}$ is injective.

Step 2. *There exists a constant* $c_1>0$ *such that*

$$\left\|\mathcal{D}^{0*}\xi_0'\right\|_{L^p}\le c_1\sup_{\xi_0''\ne0}\frac{\langle\mathcal{D}^{0*}\xi_0',\mathcal{D}^{0*}\xi_0''\rangle}{\|\mathcal{D}^{0*}\xi_0''\|_{L^q}}$$

for every $(u,A)\in\widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$ *and every* $\xi_0'\in\Omega^{0,1}(\Sigma,H_u)$.

Let $\xi_1,\dots,\xi_m$ be an L^2-orthonormal basis of $\ker\mathcal{D}^0$. Given ξ_0' choose $\xi\in L^q(\Sigma,H_u)$ such that

$$\langle\xi,\mathcal{D}^{0*}\xi_0'\rangle=\left\|\mathcal{D}^{0*}\xi_0'\right\|_{L^p}\,,\qquad \|\xi\|_{L^q}=1.$$

Let $\xi_0''\in W^{1,q}(\Sigma,\Lambda^{0,1}T^*\Sigma\otimes H_u)$ be the unique section such that

$$\xi=\mathcal{D}^{0*}\xi_0''+\sum_{j=1}^m\langle\xi,\xi_j\rangle\xi_j.$$

Then

$$\begin{aligned}
\left\|\mathcal{D}^{0*}\xi_0'\right\|_{L^p}&=\langle\mathcal{D}^{0*}\xi_0'',\mathcal{D}^{0*}\xi_0'\rangle\\
&=\left\|\xi-\sum_{j=1}^m\langle\xi,\xi_j\rangle\xi_j\right\|_{L^q}\frac{\langle\mathcal{D}^{0*}\xi_0'',\mathcal{D}^{0*}\xi_0'\rangle}{\|\mathcal{D}^{0*}\xi_0''\|_{L^q}}\\
&\le\left(1+\sum_{j=1}^m\|\xi_j\|_{L^p}\|\xi_j\|_{L^q}\right)\frac{\langle\mathcal{D}^{0*}\xi_0'',\mathcal{D}^{0*}\xi_0'\rangle}{\|\mathcal{D}^{0*}\xi_0''\|_{L^q}}\\
&\le(1+m{c_0}^2)\frac{\langle\mathcal{D}^{0*}\xi_0'',\mathcal{D}^{0*}\xi_0'\rangle}{\|\mathcal{D}^{0*}\xi_0''\|_{L^q}}.
\end{aligned}$$

Step 3. *There exists a constant* $c_2 > 0$ *such that*

$$\|\pi_u \mathcal{D}^{\varepsilon *}\zeta'\|_{L^p} \le c_2 \Big(\left\|\pi_u \mathcal{D}^{\varepsilon *}\zeta' - \mathcal{D}^{0*}\pi_u\zeta'\right\|_{L^p} + \left\|\mathcal{D}^0 \pi_u \mathcal{D}^{\varepsilon *}\zeta'\right\|_{L^p} \Big)$$

for every $(u,A) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$*, every* $\zeta' \in \mathcal{X}'_u$*, and every* $\varepsilon \in (0,1]$*.*

For every $\xi'_0 \in \Omega^{0,1}(\Sigma, H_u)$ we have

$$\begin{aligned}
\frac{\langle \mathcal{D}^{0*}\xi'_0, \mathcal{D}^{0*}\pi_u\zeta'\rangle}{\|\mathcal{D}^{0*}\xi'_0\|_{L^q}} &= \frac{\langle \mathcal{D}^{0*}\xi'_0, \mathcal{D}^{0*}\pi_u\zeta' - \pi_u\mathcal{D}^{\varepsilon *}\zeta'\rangle}{\|\mathcal{D}^{0*}\xi'_0\|_{L^q}} + \frac{\langle \xi'_0, \mathcal{D}^0\pi_u\mathcal{D}^{\varepsilon *}\zeta'\rangle}{\|\mathcal{D}^{0*}\xi'_0\|_{L^q}} \\
&\le \left\|\mathcal{D}^{0*}\pi_u\zeta' - \pi_u\mathcal{D}^{\varepsilon *}\zeta'\right\|_{L^p} + \left\|\mathcal{D}^0\pi_u\mathcal{D}^{\varepsilon *}\zeta'\right\|_{L^p} \frac{\|\xi'_0\|_{L^q}}{\|\mathcal{D}^{0*}\xi'_0\|_{L^q}} \\
&\le \left\|\mathcal{D}^{0*}\pi_u\zeta' - \pi_u\mathcal{D}^{\varepsilon *}\zeta'\right\|_{L^p} + c_0 \left\|\mathcal{D}^0\pi_u\mathcal{D}^{\varepsilon *}\zeta'\right\|_{L^p}.
\end{aligned}$$

Here the last inequality follows from Step 1. Now, by Step 2,

$$\begin{aligned}
\|\pi_u\mathcal{D}^{\varepsilon *}\zeta'\|_{L^p} &\le \left\|\mathcal{D}^{0*}\pi_u\zeta' - \pi_u\mathcal{D}^{\varepsilon *}\zeta'\right\|_{L^p} + \left\|\mathcal{D}^{0*}\pi_u\zeta'\right\|_{L^p} \\
&\le \left\|\mathcal{D}^{0*}\pi_u\zeta' - \pi_u\mathcal{D}^{\varepsilon *}\zeta'\right\|_{L^p} + c_1 \sup_{\xi'_0\neq 0} \frac{\langle \mathcal{D}^{0*}\pi_u\zeta', \mathcal{D}^{0*}\xi'_0\rangle}{\|\mathcal{D}^{0*}\xi'_0\|_{L^q}} \\
&\le (1+c_1)\left\|\mathcal{D}^{0*}\pi_u\zeta' - \pi_u\mathcal{D}^{\varepsilon *}\zeta'\right\|_{L^p} + c_0c_1\left\|\mathcal{D}^0\pi_u\mathcal{D}^{\varepsilon *}\zeta'\right\|_{L^p}.
\end{aligned}$$

Step 4. *There exist positive constants* ε_0 *and* c_3 *such that*

$$\|\pi_u\mathcal{D}^{\varepsilon *}\zeta'\|_{L^p} \le c_3\Big(\varepsilon\,\|\mathcal{D}^{\varepsilon *}\zeta'\|_{0,p,\varepsilon} + \varepsilon\,\|\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon *}\zeta'\|_{0,p,\varepsilon} + \|\pi_u\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon *}\zeta'\|_{L^p}\Big)$$

for every $(u,A) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$*, every* $\zeta' \in \mathcal{X}'_u$*, and every* $\varepsilon \in (0,\varepsilon_0]$*.*

We apply Lemma 4.5 to both operators $\mathcal{D}^\varepsilon$ and $\mathcal{D}^{\varepsilon *}$. Then, by Step 3,

$$\begin{aligned}
\|\pi_u\mathcal{D}^{\varepsilon *}\zeta'\|_{L^p} &\le c_2\Big(\left\|\pi_u\mathcal{D}^{\varepsilon *}\zeta' - \mathcal{D}^{0*}\pi_u\zeta'\right\|_{L^p} \\
&\qquad + \left\|\mathcal{D}^0\pi_u\mathcal{D}^{\varepsilon *}\zeta' - \pi_u\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon *}\zeta'\right\|_{0,p,\varepsilon} + \|\pi_u\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon *}\zeta'\|_{L^p}\Big) \\
&\le c_2\Big(c\,\|\zeta' - \pi_u\zeta'\|_{0,p,\varepsilon} + c\,\|\mathcal{D}^{\varepsilon *}\zeta' - \pi_u\mathcal{D}^{\varepsilon *}\zeta'\|_{0,p,\varepsilon} \\
&\qquad + \|\pi_u\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon *}\zeta'\|_{L^p}\Big) \\
&\le c_4\Big(\varepsilon\,\|\mathcal{D}^{\varepsilon *}\zeta'\|_{0,p,\varepsilon} + \varepsilon\,\|\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon *}\zeta'\|_{0,p,\varepsilon} + \|\pi_u\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon *}\zeta'\|_{L^p}\Big).
\end{aligned}$$

The last inequality follows from Lemma 4.7 and Proposition 4.2.

Step 5. *We prove (24) for $k = 0$.*

By Proposition 4.2 and Step 4,

$$\begin{aligned}\|\mathcal{D}^{\varepsilon *}\zeta'\|_{1,p,\varepsilon} &\le c\Big(\varepsilon \|\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon *}\zeta'\|_{1,p,\varepsilon} + \|\pi_u\mathcal{D}^{\varepsilon *}\zeta'\|_{L^p}\Big)\\ &\le c\Big(c_3\varepsilon \|\mathcal{D}^{\varepsilon *}\zeta'\|_{0,p,\varepsilon} + (1+c_3)\varepsilon \|\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon *}\zeta'\|_{0,p,\varepsilon}\\ &\qquad + c_3 \|\pi_u\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon *}\zeta'\|_{L^p}\Big)\end{aligned}$$

for all $(u, A) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$ and $\zeta' \in \mathcal{X}'_u$ and $\varepsilon \in (0, \varepsilon_0]$. With $cc_3\varepsilon \le 1/2$ we obtain (24) for $k = 0$.

Step 6. *We prove (25) for $k = 0$.*

By Proposition 4.2 and Step 4,

$$\begin{aligned}\|\mathcal{D}^{\varepsilon *}\zeta' - \pi_u\mathcal{D}^{\varepsilon *}\zeta'\|_{1,p,\varepsilon} &\le c\varepsilon\Big(\|\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon *}\zeta'\|_{1,p,\varepsilon} + \|\pi_u\mathcal{D}^{\varepsilon *}\zeta'\|_{L^p}\Big)\\ &\le c\varepsilon\Big(c_3\varepsilon \|\mathcal{D}^{\varepsilon *}\zeta'\|_{0,p,\varepsilon} + (1+c_3)\varepsilon \|\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon *}\zeta'\|_{0,p,\varepsilon}\\ &\qquad + c_3 \|\pi_u\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon *}\zeta'\|_{L^p}\Big)\\ &\le c_5\varepsilon \|\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon *}\zeta'\|_{0,p,\varepsilon}.\end{aligned}$$

Here the last inequality follows from Step 5.

Step 7. *There exist positive constants ε_0 and c_6 such that*

$$\begin{aligned}\varepsilon^2\|[\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon *}, \tilde{\nabla}_{A,v}]\zeta'\|_{L^p} &\le c_6\|\zeta'\|_{2,p,\varepsilon},\\ \varepsilon\|[\pi_u\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon *}, \tilde{\nabla}_{A,v}]\zeta'\|_{L^p} &\le c_6\left(\varepsilon\|\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon *}\zeta'\|_{0,p,\varepsilon} + \|\pi_u\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon *}\zeta'\|_{L^p} + \|\zeta'\|_{1,p,\varepsilon}\right)\end{aligned}$$

for every $(u, A) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$, every $\zeta' \in \mathcal{X}'_u$, and every $\varepsilon \in (0, \varepsilon_0]$.

The first estimate follows immediately from Lemma 4.6. To prove the second estimate, recall from Lemma 4.3 that

$$\pi_u\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon *}\zeta' = \pi_u 2DD^*\xi' + \pi_u(DJ - JD)L_u\psi'$$

where $D : \Omega^0(\Sigma, u^*TM/\mathrm{G}) \to \Omega^{0,1}(\Sigma, u^*TM/\mathrm{G})$ is the Cauchy–Riemann operator defined by (109) and $R := (DJ - JD)L_u$ is a zeroth order operator (Lemma B.5). Hence

$$2DD^*\pi_u\xi' = -2[\pi_u, DD^*]\xi' + \pi_u\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon *}\zeta' - \pi_u R\psi'.$$

By (16) in the proof of Lemma 4.5, the commutator $[\pi_u, DD^*]$ is a first order operator in ξ'. Hence there exists a constant $c_7 = c_7(p, c_0)$ such that

$$\|\tilde{\nabla}_A{}^*\tilde{\nabla}_A\pi_u\xi'\|_{L^p} \le c_7\left(\|\pi_u\mathcal{D}^\varepsilon\mathcal{D}^{\varepsilon*}\zeta'\|_{L^p} + \varepsilon^{-1}\|\zeta'\|_{1,p,\varepsilon}\right)$$

Moreover, by Lemma 4.7,

$$\begin{aligned}\|\tilde{\nabla}_A{}^*\tilde{\nabla}_A(\xi' - \pi_u\xi')\|_{L^p} &\le \varepsilon^{-2}\|\xi' - \pi_u\xi'\|_{2,p,\varepsilon}\\ &\le c_8\varepsilon^{-1}\|\mathcal{D}^{\varepsilon*}\zeta'\|_{1,p,\varepsilon}\\ &\le c_9\left(\|\mathcal{D}^\varepsilon\mathcal{D}^{\varepsilon*}\zeta'\|_{0,p,\varepsilon} + \varepsilon^{-1}\|\pi_u\mathcal{D}^\varepsilon\mathcal{D}^{\varepsilon*}\zeta'\|_{0,p,\varepsilon}\right).\end{aligned}$$

The last inequality follows from Step 5. Now the commutator

$$[\tilde{\nabla}_{A,v}, \pi_u\mathcal{D}^\varepsilon\mathcal{D}^{\varepsilon*}]\zeta' = 2[\tilde{\nabla}_{A,v}, \pi_u DD^*]\xi' + [\tilde{\nabla}_{A,v}, \pi_u R]\psi'$$

is a second order operator in ξ' and a zeroth order operator in ψ'. Hence the assertion follows from the last two inequalities.

Step 8. *We prove (24) for $k = 1$.*

Let c_{10} be the constant in (24) for $k = 0$ and c_{11} be the constant of Lemma 4.6 Then, for every $v \in \mathrm{Vect}(\Sigma)$, we have

$$\begin{aligned}&\varepsilon\|\tilde{\nabla}_{A,v}\mathcal{D}^{\varepsilon*}\zeta'\|_{1,p,\varepsilon}\\ &\le \varepsilon\|\mathcal{D}^{\varepsilon*}\tilde{\nabla}_{A,v}\zeta'\|_{1,p,\varepsilon} + \varepsilon\|\tilde{\nabla}_{A,v}\mathcal{D}^{\varepsilon*}\zeta' - \mathcal{D}^{\varepsilon*}\tilde{\nabla}_{A,v}\zeta'\|_{1,p,\varepsilon}\\ &\le c_{10}\varepsilon^2\|\mathcal{D}^\varepsilon\mathcal{D}^{\varepsilon*}\tilde{\nabla}_{A,v}\zeta'\|_{0,p,\varepsilon} + c_{10}\varepsilon\|\pi_u\mathcal{D}^\varepsilon\mathcal{D}^{\varepsilon*}\tilde{\nabla}_{A,v}\zeta'\|_{L^p} + c_{11}\|\zeta'\|_{2,p,\varepsilon}\\ &\le c_{10}\varepsilon^2\|\tilde{\nabla}_{A,v}\mathcal{D}^\varepsilon\mathcal{D}^{\varepsilon*}\zeta'\|_{0,p,\varepsilon} + c_{10}\varepsilon\|\tilde{\nabla}_{A,v}\pi_u\mathcal{D}^\varepsilon\mathcal{D}^{\varepsilon*}\zeta'\|_{L^p}\\ &\quad + c_{10}\varepsilon^2\|[\mathcal{D}^\varepsilon\mathcal{D}^{\varepsilon*}, \tilde{\nabla}_{A,v}]\zeta'\|_{0,p,\varepsilon} + c_{10}\varepsilon\|[\pi_u\mathcal{D}^\varepsilon\mathcal{D}^{\varepsilon*}, \tilde{\nabla}_{A,v}]\zeta'\|_{L^p} + c_{11}\|\zeta'\|_{2,p,\varepsilon}\\ &\le c_{10}\varepsilon^2\|\tilde{\nabla}_{A,v}\mathcal{D}^\varepsilon\mathcal{D}^{\varepsilon*}\zeta'\|_{0,p,\varepsilon} + c_{10}\varepsilon\|\tilde{\nabla}_{A,v}\pi_u\mathcal{D}^\varepsilon\mathcal{D}^{\varepsilon*}\zeta'\|_{L^p}\\ &\quad + c_6c_{10}\left(\varepsilon\|\mathcal{D}^\varepsilon\mathcal{D}^{\varepsilon*}\zeta'\|_{0,p,\varepsilon} + \|\pi_u\mathcal{D}^\varepsilon\mathcal{D}^{\varepsilon*}\zeta'\|_{L^p}\right) + (2c_6c_{10} + c_{11})\|\zeta'\|_{2,p,\varepsilon}\\ &\le c_{12}\left(\varepsilon\|\mathcal{D}^\varepsilon\mathcal{D}^{\varepsilon*}\zeta'\|_{1,p,\varepsilon} + \|\pi_u\mathcal{D}^\varepsilon\mathcal{D}^{\varepsilon*}\zeta'\|_{1,p,\varepsilon}\right).\end{aligned}$$

The penultimate inequality follows from Step 7, and the last step from (20) and (24) and the definition of the $(1, p, \varepsilon)$-norm. Now (24) for $k = 1$ follows by taking the sum over finitely many vector fields $v \in \mathrm{Vect}(\Sigma)$.

Step 9. *We prove (25) for $k = 1$.*

By Step 8, suppose that (24) holds with $k = 1$ and $c = c_{13}$, choose c_{14} such that $\|\pi_u\zeta\|_{1,p,\varepsilon} \le c_{14}\|\zeta\|_{1,p,\varepsilon}$ for every $\zeta \in \mathcal{X}_u$, and let c_{15} be the constant of

Lemma 4.5. Then

$$\begin{aligned}
\|\mathcal{D}^{\varepsilon*}\zeta' - \pi_u\mathcal{D}^{\varepsilon*}\zeta'\|_{2,p,\varepsilon} &\leq c_{13}\varepsilon\|\mathcal{D}^{\varepsilon}(\mathcal{D}^{\varepsilon*}\zeta' - \pi_u\mathcal{D}^{\varepsilon*}\zeta')\|_{1,p,\varepsilon} \\
&\quad + c_{13}\|\pi_u\mathcal{D}^{\varepsilon}(\mathcal{D}^{\varepsilon*}\zeta' - \pi_u\mathcal{D}^{\varepsilon*}\zeta')\|_{1,p,\varepsilon} \\
&\leq c_{13}\varepsilon\left(\|\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon*}\zeta'\|_{1,p,\varepsilon} + \|\pi_u\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon*}\zeta'\|_{1,p,\varepsilon}\right) \\
&\quad + c_{13}\varepsilon\|\mathcal{D}^{\varepsilon}\pi_u\mathcal{D}^{\varepsilon*}\zeta' - \pi_u\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon*}\zeta'\|_{1,p,\varepsilon} \\
&\quad + c_{13}\|(\pi_u\mathcal{D}^{\varepsilon} - \mathcal{D}^{0}\pi_u)\mathcal{D}^{\varepsilon*}\zeta'\|_{1,p,\varepsilon} \\
&\leq (1+c_{14})c_{13}\varepsilon\|\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon*}\zeta'\|_{1,p,\varepsilon} \\
&\quad + c_{15}c_{13}\left(\varepsilon\|\mathcal{D}^{\varepsilon*}\zeta'\|_{1,p,\varepsilon} + \|\mathcal{D}^{\varepsilon*}\zeta' - \pi_u\mathcal{D}^{\varepsilon*}\zeta'\|_{1,p,\varepsilon}\right) \\
&\leq c_{16}\varepsilon\|\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon*}\zeta'\|_{1,p,\varepsilon}.
\end{aligned}$$

The last inequality follows from Steps 5 and 6. □

5 Quadratic estimates

Fix $p > 2$, $c_0 > 0$ and $(u, A) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$, and consider the map

$$\mathcal{F}^{\varepsilon} = \mathcal{F}^{\varepsilon}_{(u,A)} : \mathcal{X}_u \to \mathcal{X}'_u$$

given by

$$\mathcal{F}^{\varepsilon}(\xi, \alpha) = \begin{pmatrix} \rho(\xi)\left(\bar{\partial}_{J,A+\alpha}(\exp_u(\xi))\right) \\ \varepsilon^{-2}L_u^*\xi - d_A^*\alpha \\ \varepsilon^{-2}\mu(\exp_u(\xi)) + *F_{A+\alpha} \end{pmatrix}. \tag{26}$$

Here $\rho(\xi) : T_{\exp_u(\xi)}M \to T_uM$ denotes parallel transport along the geodesic $r \mapsto \exp_u(r\xi)$ with respect to the Hermitian connection $\tilde{\nabla} := \nabla - \frac{1}{2}J\nabla J$ on TM. The differential of $\mathcal{F}^{\varepsilon}$ at zero is given by

$$d\mathcal{F}^{\varepsilon}(0) = \mathcal{D}^{\varepsilon}_{(u,A)}.$$

Let $\mathcal{F}^{\varepsilon}_i$ denote the ith component of $\mathcal{F}^{\varepsilon}$. Since $\mathcal{F}^{\varepsilon}_2$ is a linear map, the following proposition only deals with the first and third components of $\mathcal{F}^{\varepsilon}$.

Proposition 5.1. *For every $p > 2$ and every $c_0 > 0$ there exists a constant $c > 0$ such that the following holds for every $(u, A) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$, any two pairs $\zeta = (\xi, \alpha), \hat{\zeta} = (\hat{\xi}, \hat{\alpha}) \in \mathcal{X}_u$, and every $\varepsilon \in (0, 1]$.*

(i) *If* $\|\xi\|_{L^\infty} + \|\hat{\xi}\|_{L^\infty} \le 1$ *then*

$$\begin{aligned}
&\|\mathcal{F}_1^\varepsilon(\zeta+\hat{\zeta}) - \mathcal{F}_1^\varepsilon(\zeta) - d\mathcal{F}_1^\varepsilon(\zeta)\hat{\zeta}\|_{L^p} \\
&\le c\|\hat{\xi}\|_{L^\infty}\Big(\|\hat{\xi}\|_{L^p} + \|\tilde{\nabla}_A\hat{\xi}\|_{L^p} + \|\hat{\alpha}\|_{L^p}\Big) \\
&\quad + c\|\hat{\xi}\|_{L^\infty}^2\Big(\|\tilde{\nabla}_A\xi\|_{L^p} + \|\alpha\|_{L^p} + \|\tilde{\nabla}_A\hat{\xi}\|_{L^p} + \|\hat{\alpha}\|_{L^p}\Big).
\end{aligned}$$

If, in addition, $\|\tilde{\nabla}_A\xi\|_{L^p} + \|\tilde{\nabla}_A\hat{\xi}\|_{L^p} + \|\alpha\|_{L^p} + \|\hat{\alpha}\|_{L^p} \le \varepsilon^{2/p-1}$ *then*

$$\begin{aligned}
&\|\mathcal{F}_1^\varepsilon(\zeta+\hat{\zeta}) - \mathcal{F}_1^\varepsilon(\zeta) - d\mathcal{F}_1^\varepsilon(\zeta)\hat{\zeta}\|_{L^p} \\
&\le c\|\hat{\xi}\|_{L^\infty}\Big(\varepsilon^{-1}\|\hat{\xi}\|_{L^p} + \|\tilde{\nabla}_A\hat{\xi}\|_{L^p} + \|\hat{\alpha}\|_{L^p}\Big).
\end{aligned}$$

(ii) *If* $\|\xi\|_{L^\infty} + \|\hat{\xi}\|_{L^\infty} \le 1$ *then*

$$\|\mathcal{F}_3^\varepsilon(\zeta+\hat{\zeta}) - \mathcal{F}_3^\varepsilon(\zeta) - d\mathcal{F}_3^\varepsilon(\zeta)\hat{\zeta}\|_{L^p} \le c\Big(\|\hat{\alpha}\|_{L^\infty}\|\hat{\alpha}\|_{L^p} + \varepsilon^{-2}\|\hat{\xi}\|_{L^\infty}\|\hat{\xi}\|_{L^p}\Big).$$

(iii) *If* $\|\xi\|_{L^\infty} \le 1$ *then*

$$\begin{aligned}
\|d\mathcal{F}_1^\varepsilon(\zeta)\hat{\zeta} - d\mathcal{F}_1^\varepsilon(0)\hat{\zeta}\|_{L^p} \;\le\;& c\|\hat{\xi}\|_{L^\infty}\Big(\|\xi\|_{L^p} + \|\tilde{\nabla}_A\xi\|_{L^p} + \|\alpha\|_{L^p}\Big) \\
&+ c\|\xi\|_{L^\infty}\Big(\|\hat{\xi}\|_{L^p} + \|\tilde{\nabla}_A\hat{\xi}\|_{L^p} + \|\hat{\alpha}\|_{L^p}\Big).
\end{aligned}$$

(iv) *If* $\|\xi\|_{L^\infty} \le 1$ *then*

$$\|d\mathcal{F}_3^\varepsilon(\zeta)\hat{\zeta} - d\mathcal{F}_3^\varepsilon(0)\hat{\zeta}\|_{L^p} \le c\Big(\varepsilon^{-2}\|\xi\|_{L^p}\|\hat{\xi}\|_{L^\infty} + \|\alpha\|_{L^p}\|\hat{\alpha}\|_{L^\infty}\Big).$$

The estimates in Proposition 5.1 differ from the ones in [DS2] in that the first derivatives of ξ appear on the right hand sides of the inequalities. This is because the nonlinearities in the Cauchy-Riemann equations appear in the first order terms whereas the nonlinearities in the anti-self-duality equations only appear in the zeroth order terms. In our equations the nonlinear terms involving α are of zeroth order. Hence no derivatives of α appear in the quadratic estimates. This is crucial for our adiabatic limit argument.

Proof of Proposition 5.1. In local holomorphic coordinates $s + it$ on Σ the map $\mathcal{F}^\varepsilon$ is given by

$$\mathcal{F}^\varepsilon(\zeta) = \begin{pmatrix} \rho(\xi)\Big(\partial_s u_\xi + X_{\Phi+\varphi}(u_\xi) + J\big(\partial_t u_\xi + X_{\Psi+\psi}(u_\xi)\big)\Big) \\ \varepsilon^{-2}L_u^*\xi + \lambda^{-2}\left(\nabla_{A,s}\varphi + \nabla_{A,t}\psi\right) \\ \varepsilon^{-2}\mu(u_\xi) + \lambda^{-2}\left(\partial_s(\Psi+\psi) - \partial_t(\Phi+\varphi) + [\Phi+\varphi, \Psi+\psi]\right) \end{pmatrix},$$

where $u_\xi := \exp_u(\xi)$, $\zeta := (\xi, \varphi, \psi)$ and $\alpha := \varphi\, ds + \psi\, dt$. Suppose that $\|\xi\|_{L^\infty} \le 1$. The second derivatives of $\mathcal{F}_1^\varepsilon$ and $\mathcal{F}_3^\varepsilon$ satisfy the following *pointwise* estimates for suitable constants $c_1 = c_1(u, A, v_s, v_t)$ and $c_2 = c_2(u)$ (i.e. c_2 does not depend on the derivatives of u):

$$\begin{aligned}
|d^2\mathcal{F}_1^\varepsilon(\zeta)(\zeta_1,\zeta_2)| &\le c_1\Big(\big(1+|\alpha|+|\tilde\nabla_A\xi|\big)\,|\xi_1|\,|\xi_2| \\
&\qquad + |\xi_1|\,\big(|\tilde\nabla_A\xi_2|+|\alpha_2|\big) + |\xi_2|\,\big(|\tilde\nabla_A\xi_1|+|\alpha_1|\big)\Big) \qquad (27)\\
|d^2\mathcal{F}_3^\varepsilon(\zeta)(\zeta_1,\zeta_2)| &\le c_2\big(|\alpha_1||\alpha_2| + \varepsilon^{-2}|\xi_1||\xi_2|\big). \qquad (28)
\end{aligned}$$

The estimate (28) is obvious and (27) will be proved below. Now consider the identities

$$\begin{aligned}
\mathcal{F}^\varepsilon(\zeta+\hat\zeta) - \mathcal{F}^\varepsilon(\zeta) - d\mathcal{F}^\varepsilon(\zeta)\hat\zeta &= \int_0^1 (1-r)d^2\mathcal{F}^\varepsilon(\zeta+r\hat\zeta)(\hat\zeta,\hat\zeta)\,dr \qquad (29)\\
d\mathcal{F}^\varepsilon(\zeta)\hat\zeta - d\mathcal{F}^\varepsilon(0)\hat\zeta &= \int_0^1 d^2\mathcal{F}^\varepsilon(r\zeta)(\zeta,\hat\zeta)\,dr. \qquad (30)
\end{aligned}$$

To prove assertions (i) and (ii) replace (ζ,ζ_1,ζ_2) by $(\zeta + r\hat\zeta, \hat\zeta, \hat\zeta)$ in (27) and (28), insert the resulting inequalities in (29), and integrate over Σ. Moreover, to derive the second assertion in (i) from the first we use the inequality $\varepsilon^{2/p-1}\|\hat\xi\|_{L^\infty} \le c(\varepsilon^{-1}\|\hat\xi\|_{L^p} + \|\tilde\nabla_A\hat\xi\|_{L^p})$ of Lemma 4.4. To prove assertions (iii) and (iv) replace (ζ,ζ_1,ζ_2) by $(r\zeta,\zeta,\hat\zeta)$ in (27) and (28), insert the resulting inequalities in (30), and integrate over Σ.

To prove (27) we give an explicit formula for the second derivative of $\mathcal{F}_1^\varepsilon$ in local coordinates on M:

$$d^2\mathcal{F}_1^\varepsilon(\zeta)(\zeta_1,\zeta_2) = d^2\rho(\xi)(\xi_1,\xi_2)H + \big(d\rho(\xi)\xi_2\big)H_1 + \big(d\rho(\xi)\xi_1\big)H_2 + \rho(\xi)H_{12}$$

Here $E_j := E_j(u,\xi)$ for $j = 1, 2$ (see Appendix C), and H, H_1, H_2 and H_{12} are defined as follows. The section $H = H(\zeta)$ is given by

$$\begin{aligned}
H &= E_1(v_s + X_\varphi(u)) + E_2(\nabla_{A,s}\xi + \nabla_\xi X_\varphi(u)) \\
&\quad + J(u_\xi)\Big(E_1(v_t + X_\psi(u)) + E_2(\nabla_{A,t}\xi + \nabla_\xi X_\psi(u))\Big).
\end{aligned}$$

Now we use the following notation for $i = 1, 2$:

$$\begin{aligned}
\partial E_i(v,w) &:= \big(\partial_2 E_i(u,\xi)v\big)w, \\
\partial^2 E_i(v_1,v_2,w) &:= \big(\partial_2\partial_2 E_i(u,\xi)(v_1,v_2)\big)w.
\end{aligned}$$

The section $H_j = H_j(\zeta;\zeta_j)$ is linear in ζ_j and is defined by

$$\begin{aligned}
H_j \;=\; & \partial E_1(\xi_j, v_s + X_\varphi(u)) + E_1 X_{\varphi_j}(u) + \partial E_2\left(\xi_j, \nabla_{A,s}\xi + \nabla_\xi X_\varphi(u)\right) \\
& + E_2\left(\nabla_{A,s}\xi_j + \nabla_{\xi_j} X_\varphi(u) + \nabla_\xi X_{\varphi_j}(u)\right) \\
& + \Big(dJ(u_\xi)E_2\xi_j\Big)\Big(E_1(v_t + X_\psi(u)) + E_2(\nabla_{A,t}\xi + \nabla_\xi X_\psi(u))\Big) \\
& + J(u_\xi)\Big(\partial E_1(\xi_j, v_t + X_\psi(u)) + \partial E_2\left(\xi_j, \nabla_{A,t}\xi + \nabla_\xi X_\psi(u)\right) \\
& \qquad + E_1 X_{\psi_j}(u) + E_2\left(\nabla_{A,t}\xi_j + \nabla_{\xi_j} X_\psi(u) + \nabla_\xi X_{\psi_j}(u)\right)\Big).
\end{aligned}$$

The section $H_{12} = H_{12}(\zeta;\zeta_1,\zeta_2)$ is bilinear in ζ_1 and ζ_2, and is defined by

$$\begin{aligned}
H_{12} \;=\; & \partial^2 E_1(\xi_1,\xi_2, v_s + X_\varphi(u)) + \partial E_1(\xi_2, X_{\varphi_1}(u)) + \partial E_1(\xi_1, X_{\varphi_2}(u)) \\
& + E_2\left(\nabla_{\xi_2} X_{\varphi_1}(u) + \nabla_{\xi_1} X_{\varphi_2}(u)\right) + \partial^2 E_2\left(\xi_1,\xi_2, \nabla_{A,s}\xi + \nabla_\xi X_\varphi(u)\right) \\
& + \partial E_2\left(\xi_2, \nabla_{A,s}\xi_1 + \nabla_{\xi_1} X_\varphi(u) + \nabla_\xi X_{\varphi_1}(u)\right) \\
& + \partial E_2\left(\xi_1, \nabla_{A,s}\xi_2 + \nabla_{\xi_2} X_\varphi(u) + \nabla_\xi X_{\varphi_2}(u)\right) \\
& + \left(d^2 J(u_\xi)(E_2\xi_1, E_2\xi_2)\right)\Big(E_1\left(v_t + X_\psi(u)\right) + E_2\left(\nabla_{A,t}\xi + \nabla_\xi X_\psi(u)\right)\Big) \\
& + \left(dJ(u_\xi)\partial E_2(\xi_1,\xi_2)\right)\Big(E_1\left(v_t + X_\psi(u)\right) + E_2\left(\nabla_{A,t}\xi + \nabla_\xi X_\psi(u)\right)\Big) \\
& + \left(dJ(u_\xi)E_2\xi_1\right)\Big(\partial E_1(\xi_2, v_t + X_\psi(u)) + \partial E_2\left(\xi_2, \nabla_{A,t}\xi + \nabla_\xi X_\psi(u)\right) \\
& \qquad + E_1 X_{\psi_2}(u) + E_2\left(\nabla_{A,t}\xi_2 + \nabla_{\xi_2} X_\psi(u) + \nabla_\xi X_{\psi_2}(u)\right)\Big) \\
& + \left(dJ(u_\xi)E_2\xi_2\right)\Big(\partial E_1(\xi_1, v_t + X_\psi(u)) + \partial E_2(\xi_1, \nabla_{A,t}\xi + \nabla_\xi X_\psi(u)) \\
& \qquad + E_1 X_{\psi_1}(u) + E_2\left(\nabla_{A,t}\xi_1 + \nabla_{\xi_1} X_\psi(u) + \nabla_\xi X_{\psi_1}(u)\right)\Big) \\
& + J(u_\xi)\Big(\partial^2 E_1\left(\xi_1,\xi_2, v_t + X_\psi(u)\right) \\
& \qquad + \partial E_1(\xi_2, X_{\psi_1}(u)) + \partial E_1(\xi_1, X_{\psi_2}(u)) \\
& \qquad + \partial^2 E_2\left(\xi_1,\xi_2, \nabla_{A,t}\xi + \nabla_\xi X_\psi(u)\right) \\
& \qquad + \partial E_2\left(\xi_2, \nabla_{A,t}\xi_1 + \nabla_{\xi_1} X_\psi(u) + \nabla_\xi X_{\psi_1}(u)\right) \\
& \qquad + \partial E_2\left(\xi_1, \nabla_{A,t}\xi_2 + \nabla_{\xi_2} X_\psi(u) + \nabla_\xi X_{\psi_2}(u)\right) \\
& \qquad + E_2\left(\nabla_{\xi_2} X_{\psi_1}(u) + \nabla_{\xi_1} X_{\psi_2}(u)\right)\Big).
\end{aligned}$$

The inequalities (27) and (28) now follow by a term by term inspection of H, H_1, H_2, and H_{12}, assuming $\|\xi\|_{L^\infty} \le 1$. $\square$

Proposition 5.2. *For every $p > 2$ and every $c_0 > 0$ there exists a constants $c > 0$ such that the following holds for every $(u, A) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$, any two pairs $\zeta = (\xi, \alpha), \hat{\zeta} = (\hat{\xi}, \hat{\alpha}) \in \mathcal{X}_u$, and every $\varepsilon \in (0, 1]$.*
(i) *If $\|\xi\|_{L^\infty} + \|\hat{\xi}\|_{L^\infty} \le 1$ and $\|\zeta\|_{1,p,\varepsilon} + \|\hat{\zeta}\|_{1,p,\varepsilon} \le \varepsilon^{2/p}$ then*

$$\|\mathcal{F}^\varepsilon(\zeta + \hat{\zeta}) - \mathcal{F}^\varepsilon(\zeta) - d\mathcal{F}^\varepsilon(\zeta)\hat{\zeta}\|_{0,p,\varepsilon} \le c\varepsilon^{-1-2/p}\|\hat{\zeta}\|^2_{1,p,\varepsilon}.$$

(ii) *If $\|\xi\|_{L^\infty} + \|\hat{\xi}\|_{L^\infty} \le 1$ and $\|\zeta\|_{2,p,\varepsilon} + \|\hat{\zeta}\|_{2,p,\varepsilon} \le \varepsilon^{2/p}$ then*

$$\|\mathcal{F}^\varepsilon(\zeta + \hat{\zeta}) - \mathcal{F}^\varepsilon(\zeta) - d\mathcal{F}^\varepsilon(\zeta)\hat{\zeta}\|_{1,p,\varepsilon} \le c\varepsilon^{-1-2/p}\|\hat{\zeta}\|^2_{2,p,\varepsilon}.$$

(iii) *If $\|\xi\|_{L^\infty} \le 1$ then*

$$\|d\mathcal{F}^\varepsilon(\zeta)\hat{\zeta} - d\mathcal{F}^\varepsilon(0)\hat{\zeta}\|_{0,p,\varepsilon} \le c\varepsilon^{-1-2/p}\|\zeta\|_{1,p,\varepsilon}\|\hat{\zeta}\|_{1,p,\varepsilon}.$$

(iv) *If $\|\xi\|_{L^\infty} \le 1$ and $\|\zeta\|_{1,p,\varepsilon} \le \varepsilon^{2/p}$ then*

$$\|d\mathcal{F}^\varepsilon(\zeta)\hat{\zeta} - d\mathcal{F}^\varepsilon(0)\hat{\zeta}\|_{1,p,\varepsilon} \le c\varepsilon^{-1-2/p}\|\zeta\|_{2,p,\varepsilon}\|\hat{\zeta}\|_{2,p,\varepsilon}.$$

Proof. Assertions (i) and (iii) follow immediately from Proposition 5.1. To prove (ii) we observe that in estimating the quadratic terms in $\partial\mathcal{F}^\varepsilon_1$ we encounter products of the following forms

- $\partial^2\xi \cdot \hat{\xi} \cdot \hat{\xi}$ and $\partial\varphi \cdot \hat{\xi} \cdot \hat{\xi}$. Here the L^p-norms of $\partial^2\xi$ and $\partial\varphi$ can be estimated by $\varepsilon^{2/p-2}$ and the L^∞-norm of $\hat{\xi} \cdot \hat{\xi}$ by $\varepsilon^{-4/p}\|\hat{\zeta}\|^2_{1,p,\varepsilon}$.

- $\xi \cdot \partial\hat{\xi} \cdot \partial\hat{\xi}$, $\xi \cdot \hat{\xi} \cdot \partial^2\hat{\xi}$, $\xi \cdot \partial\hat{\xi} \cdot \hat{\varphi}$, and $\xi \cdot \partial\hat{\varphi} \cdot \hat{\xi}$. The L^p-norms of these products can be estimated by $\varepsilon^{-2-2/p}\|\hat{\zeta}\|_{1,p,\varepsilon}\|\hat{\zeta}\|_{2,p,\varepsilon}$.

- $\partial\xi \cdot \partial\hat{\xi} \cdot \hat{\xi}$, $\varphi \cdot \partial\hat{\xi} \cdot \hat{\xi}$, and $\partial\xi \cdot \hat{\varphi} \cdot \hat{\xi}$. In these cases the L^p-norm of $\partial\xi$ is bounded by $\varepsilon^{2/p-1}$ and the L^∞-norms of $\partial\hat{\xi} \cdot \hat{\xi}$ and $\hat{\varphi} \cdot \hat{\xi}$ are bounded by $\varepsilon^{-1-4/p}\|\hat{\zeta}\|_{1,p,\varepsilon}\|\hat{\zeta}\|_{2,p,\varepsilon}$.

Similarly, in estimating the quadratic terms in $\partial\mathcal{F}^\varepsilon_3$ we encounter products of the following forms

- $\varepsilon^{-2}\hat{\xi} \cdot \partial\hat{\xi}$ and $\partial\hat{\varphi} \cdot \hat{\psi}$. The L^p-norms of these products can be estimated by $\varepsilon^{-3-2/p}\|\hat{\zeta}\|^2_{1,p,\varepsilon}$.

- $\varepsilon^{-2}\partial\xi \cdot \hat{\xi} \cdot \hat{\xi}$. Here the L^p-norm of $\partial\xi$ bounded by $\varepsilon^{2/p-1}$ and the L^∞-norm of $\varepsilon^{-2}\hat{\xi} \cdot \hat{\xi}$ is bounded by $\varepsilon^{-2-4/p}\|\hat{\zeta}\|^2_{1,p,\varepsilon}$.

This proves (ii). The proof of (iv) is similar. □

Assertions (i) and (iii) in Proposition 5.2 are weaker than Proposition 5.1; in the former the first derivatives of α appear on the right hand sides of the estimates. The full strength of Proposition 5.1 will be required in the proof of Theorem 6.2 below.

6 Proof of Theorem B

¿From now on we assume (H1) and (H4). In this section we establish the existence of a $\mathcal{G}(P)$-equivariant map $\widetilde{\mathcal{T}}^\varepsilon : \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0) \to \widetilde{\mathcal{M}}^\varepsilon_{B,\Sigma}$ that satisfies the requirements of Theorem B.

Theorem 6.1. *For every $c_0 > 0$ and every $p > 2$ there exist positive constants ε_0 and c such that for every $\varepsilon \in (0, \varepsilon_0]$ the following holds. For every $(u_0, A_0) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$ there exists a unique pair $\zeta_\varepsilon = (\xi_\varepsilon, \alpha_\varepsilon) \in \mathcal{X}_{u_0}$ such that*

$$(u_\varepsilon, A_\varepsilon) := (\exp_{u_0}(\xi_\varepsilon), A_0 + \alpha_\varepsilon) \in \widetilde{\mathcal{M}}^\varepsilon_{B,\Sigma}, \tag{31}$$

$$-d^*_{A_0}\alpha_\varepsilon + \varepsilon^{-2} L^*_{u_0}\xi_\varepsilon = 0, \qquad \zeta_\varepsilon \in \mathrm{im}\,(\mathcal{D}^\varepsilon_{(u,A)})^*, \tag{32}$$

$$\|\zeta_\varepsilon\|_{2,p,\varepsilon;(u_0,A_0)} \le c\varepsilon^2. \tag{33}$$

The map $(u_0, A_0) \mapsto (u_\varepsilon, A_\varepsilon)$ is $\mathcal{G}(P)$-equivariant and will be denoted by $\widetilde{\mathcal{T}}^\varepsilon$.

The next theorem shows that uniqueness holds under a slightly weaker hypothesis, namely in a larger neighbourhood of (u, A).

Theorem 6.2. *For every $c_0 > 0$ and every $p > 2$ there exist positive constants δ and ε_0 such that for every $\varepsilon \in (0, \varepsilon_0]$ the following holds. Suppose that $(u_0, A_0) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$ and $(\xi, \alpha) \in \mathcal{X}_{u_0}$ satisfy (31), (32), and*

$$\|\xi\|_{1,p,\varepsilon} + \varepsilon \|\alpha\|_{L^p} + \varepsilon^{2/p} \|(\xi, \alpha)\|_{\infty,\varepsilon} \le \delta\varepsilon^{2/p+1/2}. \tag{34}$$

Then $(\exp_{u_0}(\xi), A_0 + \alpha) = \widetilde{\mathcal{T}}^\varepsilon(u, A)$.

Corollary 6.3. *For every* $c_0 > 0$ *and every* $p > 2$ *there exist positive constants* δ *and* ε_0 *such that for every* $\varepsilon \in (0, \varepsilon_0]$ *the following holds. Suppose that* $(u_0, A_0) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$ *and* $\zeta = (\xi, \alpha) \in \mathcal{X}_{u_0}$ *satisfy (31), (32), and*

$$\|\zeta\|_{1,p,\varepsilon;(u_0,A_0)} \leq \delta\varepsilon^{2/p+1/2}. \tag{35}$$

Then $(\exp_{u_0}(\xi), A_0 + \alpha) = \widetilde{\mathcal{T}}^\varepsilon(u, A)$.

Proof. Theorem 6.2 and Lemma 4.4. □

Proof of Theorem 6.1. The proof is similar to that of Theorem 5.1 in [DS2]. However, in the present case the nonlinearities (in the quadratic estimates) appear in the highest order terms, and we establish estimates for the $(2, p, \varepsilon)$-norms and not just the $(1, p, \varepsilon)$-norms (as in [DS2]). We assume throughout that the exponential map at each point in $\mu^{-1}(0)$ is defined in a ball of radius one.

Abbreviate $\mathcal{D}^\varepsilon := \mathcal{D}^\varepsilon_{(u_0,A_0)}$ and let $\mathcal{F}^\varepsilon : \mathcal{X}_{u_0} \to \mathcal{X}'_{u_0}$ be defined by (26). Then

$$d\mathcal{F}^\varepsilon(0) = \mathcal{D}^\varepsilon, \qquad \mathcal{F}^\varepsilon(0) = (0, 0, *F_{A_0}).$$

Hence, by Lemma 2.2, there exists a constant $C_0 > 0$, depending only on c_0 and p, such that

$$\|\mathcal{F}^\varepsilon(0)\|_{1,p,\varepsilon} = \varepsilon \|F_{A_0}\|_{L^p} + \varepsilon^2 \|d_{A_0} * F_{A_0}\|_{L^p} \leq C_0\varepsilon.$$

We use Newton iteration to obtain a zero of $\mathcal{F}^\varepsilon$, and hence a solution of (7). Let $\zeta_\nu = (\xi_\nu, \alpha_\nu) \in \mathcal{X}_{u_0}$ be the sequence defined recursively by $\zeta_0 := 0$ and

$$\zeta_{\nu+1} := \zeta_\nu + \hat{\zeta}_\nu, \qquad \hat{\zeta}_\nu \in \operatorname{im}\mathcal{D}^{\varepsilon*}, \qquad \mathcal{D}^\varepsilon\hat{\zeta}_\nu = -\mathcal{F}^\varepsilon(\zeta_\nu). \tag{36}$$

We prove by induction over ν that there exist positive constants ε_0, c_1, C, depending only on c_0 and p, such that

$$\begin{aligned} \|\hat{\zeta}_\nu\|_{2,p,\varepsilon} &\leq c_1\|\mathcal{F}^\varepsilon(\zeta_\nu)\|_{1,p,\varepsilon}, && (37)\\ \|\hat{\zeta}_\nu\|_{2,p,\varepsilon} &\leq 2^{-\nu}C_0c_1\varepsilon^2, && (38)\\ \|\mathcal{F}^\varepsilon(\zeta_{\nu+1})\|_{1,p,\varepsilon} &\leq C\varepsilon^{1-2/p}\|\hat{\zeta}_\nu\|_{2,p,\varepsilon} && (39) \end{aligned}$$

for $\varepsilon \in (0, \varepsilon_0]$ and $\nu \geq 0$. The constants are chosen such that the linear estimates of Lemma 4.8 hold for $0 < \varepsilon \leq \varepsilon_0$ with $c = c_1 \geq 1$, the quadratic estimates of Proposition 5.2 hold for $0 < \varepsilon \leq 1$ with $c = c_2 \geq 1$, the L^∞ estimates of Lemma 4.4 hold for $0 < \varepsilon \leq 1$ with $c = c_\infty \geq 1$, and

$$C = 3C_0c_1c_2, \qquad Cc_1\varepsilon_0^{1-2/p} \leq 1/2, \qquad 3C_0c_1c_\infty\varepsilon_0^{2-2/p} \leq 1.$$

For $\nu = 0$ the estimates (37) and (38) follow from Lemma 4.8. Namely, by (24) with $k = 1$, we have

$$\|\hat{\zeta}_0\|_{2,p,\varepsilon} \leq c_1\varepsilon\|\mathcal{F}^\varepsilon(0)\|_{1,p,\varepsilon} \leq C_0c_1\varepsilon^2.$$

The estimate (39) for $\nu = 0$ follows from the identity $d\mathcal{F}^\varepsilon(0)\hat{\zeta}_0 = -\mathcal{F}^\varepsilon(0)$ and Proposition 5.2 (ii). Namely, since $C_0c_1c_\infty\varepsilon^{2-2/p} \leq 1$ we have

$$\|\hat{\zeta}_0\|_{2,p,\varepsilon} \leq C_0c_1\varepsilon^2 \leq \varepsilon^{2/p}, \qquad \|\hat{\zeta}_0\|_{\infty,\varepsilon} \leq C_0c_1c_\infty\varepsilon^{2-2/p} \leq 1.$$

Hence the hypotheses of Proposition 5.2 (ii) are satisfied with $\zeta = 0$ and $\hat{\zeta} = \hat{\zeta}_0$, and hence

$$\begin{aligned}
\|\mathcal{F}^\varepsilon(\zeta_1)\|_{1,p,\varepsilon} &= \|\mathcal{F}^\varepsilon(\hat{\zeta}_0) - \mathcal{F}^\varepsilon(0) - d\mathcal{F}^\varepsilon(0)\hat{\zeta}_0\|_{1,p,\varepsilon} \\
&\leq c_2\varepsilon^{-1-2/p}\|\hat{\zeta}_0\|_{2,p,\varepsilon}^2 \\
&\leq C_0c_1c_2\varepsilon^{1-2/p}\|\hat{\zeta}_0\|_{2,p,\varepsilon}.
\end{aligned}$$

Since $C \geq C_0c_1c_2$ this proves (39) for $\nu = 0$. Now assume that the sequences $\zeta_0, \ldots, \zeta_\nu$ and $\hat{\zeta}_0, \ldots, \hat{\zeta}_{\nu-1}$ have been constructed up to some integer $\nu \geq 1$ and that the estimates (37-39) have been established for all integers up to $\nu - 1$. Then, by (38),

$$\|\zeta_\nu\|_{2,p,\varepsilon} \leq \sum_{j=0}^{\nu-1}\|\zeta_{j+1} - \zeta_j\|_{2,p,\varepsilon} = \sum_{j=0}^{\nu-1}\|\hat{\zeta}_j\|_{2,p,\varepsilon} \leq 2C_0c_1\varepsilon^2, \tag{40}$$

and hence

$$\|\zeta_\nu\|_{\infty,\varepsilon} \leq 2C_0c_1c_\infty\varepsilon^{2-2/p} \leq 1.$$

This shows that $\xi_\nu(p)$ lies in the domain of the exponential map at $u_0(p)$ for every $p \in P$ and so ζ_ν lies in the domain of $\mathcal{F}^\varepsilon$. Let $\hat{\zeta}_\nu$ and $\zeta_{\nu+1}$ be defined by (36). Then, by Lemma 4.8, $\hat{\zeta}_\nu$ satisfies the estimate (37). To prove (38) we observe that, by the induction hypothesis,

$$\begin{aligned}
\|\hat{\zeta}_\nu\|_{2,p,\varepsilon} &\leq c_1\|\mathcal{F}^\varepsilon(\zeta_\nu)\|_{1,p,\varepsilon} \\
&\leq Cc_1\varepsilon^{1-2/p}\|\hat{\zeta}_{\nu-1}\|_{2,p,\varepsilon} \\
&\leq 2^{-1}\|\hat{\zeta}_{\nu-1}\|_{2,p,\varepsilon} \\
&\leq 2^{-\nu}C_0c_1\varepsilon^2.
\end{aligned}$$

To prove (39) we observe that, by (40),

$$\begin{array}{rcccl} \|\zeta_\nu\|_{2,p,\varepsilon} + \|\hat{\zeta}_\nu\|_{2,p,\varepsilon} & \leq & 3C_0c_1\varepsilon^2 & \leq & \varepsilon^{2/p}, \\ \|\zeta_\nu\|_{\infty,\varepsilon} + \|\hat{\zeta}_\nu\|_{\infty,\varepsilon} & \leq & 3C_0c_1c_\infty\varepsilon^{2-2/p} & \leq & 1. \end{array}$$

Thus the hypotheses of Proposition 5.2 (ii) and (iv) are satisfied with $\zeta = \zeta_\nu$ and $\hat{\zeta} = \hat{\zeta}_\nu$. Hence

$$\begin{aligned} \|\mathcal{F}^\varepsilon(\zeta_{\nu+1})\|_{1,p,\varepsilon} &\leq \|\mathcal{F}^\varepsilon(\zeta_\nu + \hat{\zeta}_\nu) - \mathcal{F}^\varepsilon(\zeta_\nu) - d\mathcal{F}^\varepsilon(\zeta_\nu)\hat{\zeta}_\nu\|_{1,p,\varepsilon} \\ &\quad + \|d\mathcal{F}^\varepsilon(\zeta_\nu)\hat{\zeta}_\nu - d\mathcal{F}^\varepsilon(0)\hat{\zeta}_\nu\|_{1,p,\varepsilon} \\ &\leq c_2\varepsilon^{-1-2/p}\big(\|\hat{\zeta}_\nu\|_{2,p,\varepsilon} + \|\zeta_\nu\|_{2,p,\varepsilon}\big)\|\hat{\zeta}_\nu\|_{2,p,\varepsilon} \\ &\leq 3C_0c_1c_2\varepsilon^{1-2/p}\|\hat{\zeta}_\nu\|_{2,p,\varepsilon} \\ &= C\varepsilon^{1-2/p}\|\hat{\zeta}_\nu\|_{2,p,\varepsilon}. \end{aligned}$$

This completes the induction.

By (38), the sequence ζ_ν is Cauchy in the $W^{2,p}$-norm. Moreover, by examining the second component of $\mathcal{F}^\varepsilon$ we find that ζ_ν satisfies (32) for every ν and hence so does its limit

$$\zeta_\varepsilon := (\xi_\varepsilon, \alpha_\varepsilon) := \lim_{\nu\to\infty}(\xi_\nu, \alpha_\nu).$$

By (40), this limit also satisfies (33) with $c := 2C_0c_1$. Moreover, by (38) and (39), the sequence $\mathcal{F}^\varepsilon(\zeta_\nu)$ converges to zero in the $(1, p, \varepsilon)$-norm and hence $\mathcal{F}^\varepsilon(\zeta_\varepsilon) = 0$. Hence ζ_ε satisfies (31) and it follows from elliptic regularity that ζ_ε is smooth. This proves existence.

We prove uniqueness. Suppose $\zeta = (\xi, \alpha) \in \mathcal{X}_{u_0}$ satisfies (31), (32), and $\|\zeta\|_{1,p,\varepsilon} \leq c\varepsilon^2$. Then, by Lemma 4.8 and Proposition 5.2 (i) and (iii),

$$\begin{aligned} \|\zeta - \zeta_\varepsilon\|_{1,p,\varepsilon} &\leq c_1\|\mathcal{D}^\varepsilon(\zeta - \zeta_\varepsilon)\|_{0,p,\varepsilon} \\ &\leq c_1\|\mathcal{F}^\varepsilon(\zeta) - \mathcal{F}^\varepsilon(\zeta_\varepsilon) - d\mathcal{F}^\varepsilon(\zeta_\varepsilon)(\zeta - \zeta_\varepsilon)\|_{0,p,\varepsilon} \\ &\quad + c_1\|d\mathcal{F}^\varepsilon(0)(\zeta - \zeta_\varepsilon) - d\mathcal{F}^\varepsilon(\zeta_\varepsilon)(\zeta - \zeta_\varepsilon)\|_{0,p,\varepsilon} \\ &\leq c_1c_2\varepsilon^{-1-2/p}\big(\|\zeta - \zeta_\varepsilon\|_{1,p,\varepsilon} + \|\zeta_\varepsilon\|_{1,p,\varepsilon}\big)\|\zeta - \zeta_\varepsilon\|_{1,p,\varepsilon} \\ &\leq 3cc_1c_2\varepsilon^{1-2/p}\|\zeta - \zeta_\varepsilon\|_{1,p,\varepsilon}. \end{aligned}$$

If $3cc_1c_2\varepsilon^{1-2/p} < 1$ then $\zeta = \zeta_\varepsilon$. This proves uniqueness.

Since the conditions (31), (32), and (33) are gauge invariant, it follows that the map $(u, A) \mapsto (u_\varepsilon, A_\varepsilon)$ is $\mathcal{G}(P)$-equivariant. □

Proof of Theorem 6.2. In this proof we drop the subscript 0. Fix two pairs $(u, A) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$ and $\zeta = (\xi, \alpha) \in \mathcal{X}_u$ that satisfy (32), (34), and

$$(\exp_u(\xi), A + \alpha) \in \widetilde{\mathcal{M}}^\varepsilon_{B,\Sigma}.$$

We prove that ζ satisfies (33), provided that δ and ε are sufficiently small. By ellipticity of the operator $\mathcal{D}^0 := \mathcal{D}^0_{(u,A)}$, there exists a constant $c_1 = c_1(p, c_0) > 0$ such that

$$\|\tilde{\nabla}_A \pi_u \xi\|_{L^p} \le c_1 \left(\|\mathcal{D}^0 \pi_u \xi\|_{L^p} + \|\pi_u \xi\|_{L^p} \right).$$

Now let $c_2 = c_2(p, c_0)$ be the constant of Lemma 4.5 and $c_3 = c_3(p, c_0)$ be the constant of Proposition 4.2. Then

$$\begin{aligned}
\|\tilde{\nabla}_A \xi\|_{L^p} &\le \varepsilon^{-1} \|\xi - \pi_u \xi\|_{1,p,\varepsilon} + \|\tilde{\nabla}_A \pi_u \xi\|_{L^p} \\
&\le \varepsilon^{-1} \|\xi - \pi_u \xi\|_{1,p,\varepsilon} + c_1 \left(\|\mathcal{D}^0 \pi_u \xi\|_{L^p} + \|\pi_u \xi\|_{L^p} \right) \\
&\le \varepsilon^{-1} \|\xi - \pi_u \xi\|_{1,p,\varepsilon} \\
&\quad + c_1 \left(\|(\mathcal{D}^0 \pi_u - \pi_u \mathcal{D}^\varepsilon)\xi\|_{L^p} + \|\pi_u \mathcal{D}^\varepsilon \zeta\|_{L^p} + \|\pi_u \xi\|_{L^p} \right) \\
&\le (\varepsilon^{-1} + c_1 c_2) \|\xi - \pi_u \xi\|_{1,p,\varepsilon} + c_1 \left(\|\pi_u \mathcal{D}^\varepsilon \zeta\|_{L^p} + \|\pi_u \xi\|_{L^p} \right) \\
&\le c_4 \left(\|\mathcal{D}^\varepsilon \zeta\|_{0,p,\varepsilon} + \|\xi\|_{L^p} \right),
\end{aligned}$$

where $c_4 := c_3(1 + c_1 c_2) + c_1$. Hence

$$\begin{aligned}
\|\tilde{\nabla}_A \xi\|_{L^p} + \|\alpha\|_{L^p} &\le c_4 \|\mathcal{D}^\varepsilon \zeta\|_{0,p,\varepsilon} + c_4 \|\xi\|_{L^p} + \varepsilon^{-1} \|\zeta - \pi_u \zeta\|_{1,p,\varepsilon} \\
&\le (c_4 + c_5) \|\mathcal{D}^\varepsilon \zeta\|_{0,p,\varepsilon} + c_4 \|\xi\|_{L^p},
\end{aligned}$$

where $c_5 = c_5(p, c_0)$ is the constant of Lemma 4.8. Since

$$\mathcal{F}^\varepsilon(\zeta) = 0, \qquad \mathcal{F}^\varepsilon(0) = (0, 0, *F_A), \qquad \mathcal{D}^\varepsilon = d\mathcal{F}^\varepsilon(0)$$

we obtain

$$\begin{aligned}
\|\tilde{\nabla}_A \xi\|_{L^p} + \|\alpha\|_{L^p} &\le c_6 \|\mathcal{F}^\varepsilon(\zeta) - \mathcal{F}^\varepsilon(0) - d\mathcal{F}^\varepsilon(0)\zeta\|_{0,p,\varepsilon} \\
&\quad + c_6 \varepsilon \|F_A\|_{L^p} + c_4 \|\xi\|_{L^p}, \qquad (41)
\end{aligned}$$

where $c_6 := c_4 + c_5$. Now we use the refined quadratic estimate of Proposition 5.1 with $c = c_7$. By (34), we have

$$\|\tilde{\nabla}_A \xi\|_{L^p} + \|\alpha\|_{L^p} \le \delta \varepsilon^{2/p - 1/2} \le \varepsilon^{2/p - 1}$$

(provided that $\delta \le 1$). Thus the hypotheses of Proposition 5.1 (i) and (ii) are satisfied with $\zeta = 0$ and $\hat{\zeta}$ replaced by ζ. Hence, by (41),

$$\begin{aligned}
\|\tilde{\nabla}_A\xi\|_{L^p} + \|\alpha\|_{L^p} &\le c_6c_7\|\xi\|_{L^\infty}\Big(\varepsilon^{-1}\|\xi\|_{L^p} + \|\tilde{\nabla}_A\xi\|_{L^p} + \|\alpha\|_{L^p}\Big) \\
&\quad + c_6c_7\varepsilon\left(\|\alpha\|_{L^\infty}\|\alpha\|_{L^p} + \varepsilon^{-2}\|\xi\|_{L^\infty}\|\xi\|_{L^p}\right) \\
&\quad + c_6\varepsilon\|F_A\|_{L^p} + c_4\|\xi\|_{L^p} \\
&\le 3c_6c_7\|\zeta\|_{\infty,\varepsilon}\Big(\varepsilon^{-1}\|\xi\|_{L^p} + \|\tilde{\nabla}_A\xi\|_{L^p} + \|\alpha\|_{L^p}\Big) \\
&\quad + c_6\varepsilon\|F_A\|_{L^p} + c_4\|\xi\|_{L^p} \\
&\le 3c_6c_7\delta\varepsilon^{1/2}\Big(\varepsilon^{-1}\|\xi\|_{L^p} + \|\tilde{\nabla}_A\xi\|_{L^p} + \|\alpha\|_{L^p}\Big) \\
&\quad + c_0c_6\varepsilon + c_4\|\xi\|_{L^p} \\
&\le 3c_6c_7\delta\varepsilon^{1/2}\Big(\|\tilde{\nabla}_A\xi\|_{L^p} + \|\alpha\|_{L^p}\Big) \\
&\quad + c_0c_6\varepsilon + c_4\delta\varepsilon^{2/p+1/2} + 3c_6c_7\delta^2\varepsilon^{2/p}.
\end{aligned}$$

Here the last two inequalities follow from (34). With $3c_6c_7\delta\varepsilon^{1/2} \le 1/2$ we have

$$\|\tilde{\nabla}_A\xi\|_{L^p} + \|\alpha\|_{L^p} \le c_8(\delta + \varepsilon^{1-2/p})\varepsilon^{2/p}, \tag{42}$$

where $c_8 := 2c_0c_6 + 2c_4 + 6c_6c_7$. Since ζ satisfies (32) we can apply Lemma 4.8 (with $c = c_5$) to obtain

$$\begin{aligned}
\|\zeta\|_{1,p,\varepsilon} &\le c_5\left(\varepsilon\|\mathcal{D}^\varepsilon\zeta\|_{L^p} + \|\pi_u\mathcal{D}^\varepsilon\zeta\|_{L^p}\right) \\
&\le c_5\left(2\|d\mathcal{F}_1^\varepsilon(0)\zeta\|_{L^p} + \varepsilon^2\|d\mathcal{F}_3^\varepsilon(0)\zeta\|_{L^p}\right).
\end{aligned} \tag{43}$$

By Proposition 5.1 (i) with $c = c_7$ and (42), we now have

$$\begin{aligned}
\|d\mathcal{F}_1^\varepsilon(0)\zeta\|_{L^p} &= \|\mathcal{F}_1^\varepsilon(\zeta) - \mathcal{F}_1^\varepsilon(0) - d\mathcal{F}_1^\varepsilon(0)\zeta\|_{L^p} \\
&\le c_7\|\xi\|_{L^\infty}\Big(\|\xi\|_{L^p} + \|\tilde{\nabla}_A\xi\|_{L^p} + \|\alpha\|_{L^p}\Big) \\
&\quad + c_7\|\xi\|_{L^\infty}^2\Big(\|\tilde{\nabla}_A\xi\|_{L^p} + \|\alpha\|_{L^p}\Big) \\
&\le 2c_7\|\xi\|_{L^\infty}\Big(\|\xi\|_{L^p} + \|\tilde{\nabla}_A\xi\|_{L^p} + \|\alpha\|_{L^p}\Big) \\
&\le 2c_7\|\zeta\|_{\infty,\varepsilon}\left(\delta\varepsilon^{2/p+1/2} + c_8(\delta + \varepsilon^{1-2/p})\varepsilon^{2/p}\right) \\
&\le c_9(\delta + \varepsilon^{1-2/p})\|\zeta\|_{1,p,\varepsilon}.
\end{aligned}$$

Here we have used the fact that $\|\xi\|_{L^\infty} \le \delta\varepsilon^{1/2} \le 1$. Moreover, the penultimate inequality follows from (34) and (42) and the last inequality, with a suitable constant $c_9 = c_9(p, c_0)$, follows from Lemma 4.4. By Proposition 5.1 (ii) with $c = c_7$, we have

$$\begin{array}{rcl}\|d\mathcal{F}_3^\varepsilon(0)\zeta\|_{L^p} & \le & \|\mathcal{F}_3^\varepsilon(0) - \mathcal{F}_3^\varepsilon(\zeta) - d\mathcal{F}_3^\varepsilon(0)\zeta\|_{L^p} + \|F_A\|_{L^p} \\ & \le & c_7\left(\|\alpha\|_{L^\infty}\|\alpha\|_{L^p} + \varepsilon^{-2}\|\xi\|_{L^\infty}\|\xi\|_{L^p}\right) + \|F_A\|_{L^p} \\ & \le & 2c_7\left(\varepsilon^{-1}\|\alpha\|_{L^p} + \varepsilon^{-2}\|\xi\|_{L^p}\right)\|\zeta\|_{\infty,\varepsilon} + \|F_A\|_{L^p} \\ & \le & 2c_7\delta\varepsilon^{2/p}\varepsilon^{-3/2}\|\zeta\|_{\infty,\varepsilon} + \|F_A\|_{L^p} \\ & \le & c_{10}\delta\varepsilon^{-3/2}\|\zeta\|_{1,p,\varepsilon} + \|F_A\|_{L^p}.\end{array}$$

Here the penultimate inequality follows from (34) and the last follows from Lemma 4.4. Combining these two estimates with (43) we obtain

$$\|\zeta\|_{1,p,\varepsilon} \le c_5\left(2c_9(\delta + \varepsilon^{1-2/p}) + c_{10}\delta\varepsilon^{1/2}\right)\|\zeta\|_{1,p,\varepsilon} + c_5\varepsilon^2\|F_A\|_{L^p}.$$

If δ and ε are sufficiently small, we obtain

$$\|\zeta\|_{1,p,\varepsilon} \le 2c_5\varepsilon^2\|F_A\|_{L^p} \le 2c_0c_5\varepsilon^2.$$

Hence the result follows from the uniqueness argument at the end of the proof of Theorem 6.1. □

Corollary 6.3 has a slightly stronger hypothesis than Theorem 6.2, however, it does not seem to have a simpler proof. In order to significantly simplify the proof we would have to further strengthen the hypthesis and assume

$$\|\zeta\|_{1,p,\varepsilon} \le \delta\varepsilon^{2/p+1}$$

with a small constant δ (instead of $\|\zeta\|_{1,p,\varepsilon} \le \delta\varepsilon^{2/p+1/2}$ as in Corollary 6.3). Under this hypothesis uniqueness can be established with the same straight forward argument that is used at the end of the proof of Theorem 6.1. However, such a weaker result just fails to suffice for the proof of Theorem D. Namely, in Section 10 we shall establish an inequality of the form

$$\|\zeta\|_{1,p,\varepsilon} \le c\varepsilon^{2/p+1}$$

under the hypotheses of Theorem D. In this inequality the constant c is not small and so the argument in the proof of Theorem 6.1 does not suffice to give uniqueness. However, if ε is chosen so small that $c\varepsilon^{1/2} \le \delta$ then we can use Corollary 6.3 to obtain uniqueness.

7 Relative Coulomb gauge

This section is of preparatory nature. We prove a local slice theorem for the action of the gauge group $\mathcal{G} = \mathcal{G}(P)$ on $\mathcal{B} = C^\infty_{\mathrm{G}}(P, M) \times \mathcal{A}(P)$. The infinitesimal action is the operator $d_{(u,A)} : \Omega^0(\Sigma, \mathfrak{g}_P) \to T_{(u,A)}\mathcal{B}$ given by

$$d_{(u,A)}\eta := (-L_u\eta, d_A\eta).$$

Denote by $d^{*_\varepsilon}_{(u,A)}$ its formal adjoint with respect to the ε-inner product, i.e.

$$d^{*_\varepsilon}_{(u,A)}(\xi, \alpha) := \varepsilon^2 d_A^*\alpha - L_u^*\xi$$

for $(\xi, \alpha) \in T_{(u,A)}\mathcal{B}$. The next proposition restates the local slice theorem for the $\mathcal{G}$-action on $\mathcal{B}$ with ε-dependent norms for elements (u_0, A_0) of the moduli space $\widetilde{\mathcal{M}}^0_{B,\Sigma}$. The result continues to hold for every element $(u_0, A_0) \in \mathcal{B}$ with $\mu(u_0) = 0$. However, in this generality, more care must be taken in determining the norm on $\mathcal{B}$ with respect to which the constants c and δ depend continuously on (u_0, A_0). In the case of J-holomorphic curves the $W^{1,p}$-norm controls all higher derivatives and therefore the choice of the norm is immaterial.

Proposition 7.1. *Assume (H1). For every $p > 2$ and every $c_0 > 0$ there exist positive constants δ and c such that for every $\varepsilon \in (0, 1]$ the following holds. Let $(u_0, A_0) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$ and $\zeta = (\xi, \alpha) \in T_{(u_0,A_0)}\mathcal{B}$ such that*

$$\|\zeta\|_{1,p,\varepsilon;(u_0,A_0)} \le \delta\varepsilon^{2/p}. \tag{44}$$

Denote $(u, A) := (\exp_{u_0}(\xi), A_0 + \alpha)$. Then there exist a unique pair $\zeta_0 = (\xi_0, \alpha_0) \in T_{(u_0,A_0)}\mathcal{B}$ and a unique section $\eta_0 \in \Omega^0(\Sigma, \mathfrak{g}_P)$ such that

$$d^{*_\varepsilon}_{(u_0,A_0)}\zeta_0 = 0, \qquad g^*(u, A) = (\exp_{u_0}(\xi_0), A_0 + \alpha_0), \qquad g := e^{\eta_0},$$

and

$$\|\eta_0\|_{2,p,\varepsilon;A_0} + \|\zeta_0\|_{1,p,\varepsilon;(u_0,A_0)} \le c\,\|\zeta\|_{1,p,\varepsilon;(u_0,A_0)}\,. \tag{45}$$

Proposition 7.1 can be understood as a quantitative version of the implicit function theorem with ε-dependent norms and constants independent of ε. As in the case of Theorem 6.1 we shall prove it with a Newton type iteration. The relevant linear estimates are established in Lemma 7.2 below, and the quadratic estimate in Lemma 7.3.

Lemma 7.2. *For every $p \geq 2$ and every $c_0 > 0$ there exists a constant $c > 0$ such that the following holds for every $(u_0, A_0) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$ and every $\varepsilon \in (0, 1]$. For every $\zeta = (\xi, \alpha) \in W^{1,p}(\Sigma, u_0^*TM/\mathrm{G}) \oplus T^*\Sigma \otimes \mathfrak{g}_P)$ there exists a unique $\eta \in W^{2,p}(\Sigma, \mathfrak{g}_P)$ such that*

$$d^{*_\varepsilon}_{(u_0,A_0)} d_{(u_0,A_0)}\eta = d^{*_\varepsilon}_{(u_0,A_0)}\zeta. \tag{46}$$

Moreover, η satisfies the estimates

$$\|\eta\|_{1,p,\varepsilon} \leq c\,\|\zeta\|_{0,p,\varepsilon}\,, \qquad \|\eta\|_{2,p,\varepsilon} \leq c\,\left\|d^{*_\varepsilon}_{(u_0,A_0)}\zeta\right\|_{L^p}. \tag{47}$$

Lemma 7.3. *For every $p > 2$ and every $c_0 > 0$ there exist positive constants δ and c such that the following holds for every $(u_0, A_0) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$ and every $\varepsilon \in (0, 1]$. Assume that $\zeta_0 = (\xi_0, \alpha_0) \in T_{(u_0,A_0)}\mathcal{B}$ and $\eta \in \Omega^0(\Sigma, \mathfrak{g}_P)$ satisfy the inequality*

$$\|\eta\|_{2,p,\varepsilon} + \|\zeta_0\|_{1,p,\varepsilon} \leq \delta\varepsilon^{2/p}. \tag{48}$$

Then there exists a unique pair $\zeta_1 = (\xi_1, \alpha_1) \in T_{(u_0,A_0)}\mathcal{B}$ such that

$$(\exp_{u_0}(\xi_1), A_0 + \alpha_1) = g^*(\exp_{u_0}(\xi_0), A_0 + \alpha_0), \tag{49}$$

where $g := e^\eta$ and

$$\|\zeta_1 - \zeta_0\|_{0,p,\varepsilon} \leq c\,\|\eta\|_{1,p,\varepsilon}\,, \qquad \|\zeta_1 - \zeta_0\|_{1,p,\varepsilon} \leq c\,\|\eta\|_{2,p,\varepsilon}\,. \tag{50}$$

Moreover,

$$\left\|d^{*_\varepsilon}_{(u_0,A_0)}\left(\zeta_1 - \zeta_0 - d_{(u_0,A_0)}\eta\right)\right\|_{L^p} \leq c\varepsilon^{-2/p}\left(\|\zeta_0\|_{1,p,\varepsilon} + \|\eta\|_{2,p,\varepsilon}\right)\|\eta\|_{1,p,\varepsilon}\,. \tag{51}$$

Lemma 7.4. *For every $p \geq 2$ and every $c_0 > 0$ there exist positive constant δ and c such that the following holds for every $(u_0, A_0) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$ and every $\varepsilon \in (0, 1]$. If $\eta_1, \eta_2 \in \Omega^0(\Sigma, \mathfrak{g}_P)$ satisfy $\|\eta_1\|_{L^\infty} \leq \delta$ and $\|\eta_2\|_{L^\infty} \leq \delta$ then there exists a unique element $\eta \in \Omega^0(\Sigma, \mathfrak{g}_P)$ such that*

$$e^\eta = e^{\eta_1}e^{\eta_2}, \qquad 2^{-1}\|\eta\|_{L^\infty} \leq \|\eta_1 + \eta_2\|_{L^\infty} \leq 2\,\|\eta\|_{L^\infty}\,.$$

Moreover, η satisfies the estimate

$$c^{-1}\|\eta\|_{2,p,\varepsilon;A_0} \leq \|\eta_1 + \eta_2\|_{2,p,\varepsilon;A_0} \leq c\,\|\eta\|_{2,p,\varepsilon;A_0}\,.$$

Proof. For a fixed connection A_0 and $\varepsilon = 1$ the result is obvious. Choose c_1, c_2, c_3 such that

$$c_k^{-1} \|\eta\|_{W^{k,p}} \le \|\eta_1 + \eta_2\|_{W^{k,p}} \le c_k \|\eta\|_{W^{k,p}}$$

for $k = 1, 2, 3$, whenever η_1, η_2, η are sufficiently small in the C^0-norm and satisfy $e^\eta = e^{\eta_1} e^{\eta_2}$. Here the $W^{k,p}$ norms are understood with respect to the connection A_0. It follows that

$$\varepsilon \|d_{A_0}(\eta_1 + \eta_2)\|_{L^p} \le \varepsilon c_1 \left(\|d_{A_0}\eta\|_{L^p} + \|\eta\|_{L^p} \right) \le c_1 \|\eta\|_{1,p,\varepsilon;A_0}$$

and hence

$$\|\eta_1 + \eta_2\|_{1,p,\varepsilon;A_0} \le (c_0 + c_1) \|\eta\|_{1,p,\varepsilon;A_0}$$

for $0 \le \varepsilon \le 1$. The other three inequalities follow by similar arguments. This proves the lemma for a fixed connection A_0. Moreover, the constant c depends continuously on A_0 with respect to the C^1-norm, and is gauge invariant (with respect to the action of $\mathcal{G}$ on $\Omega^0(\Sigma, \mathfrak{g}_P)$ by conjugation). Hence, by Lemma 2.2, it can be chosen independent of A_0 as long as $(u_0, A_0) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$ for some u_0. □

Proof of Lemma 7.2. The operator $d^{*_\varepsilon}_{(u_0,A_0)} d_{(u_0,A_0)} : W^{2,p}(\Sigma, \mathfrak{g}_P) \to L^p(\Sigma, \mathfrak{g}_P)$ is given by

$$d^{*_\varepsilon}_{(u_0,A_0)} d_{(u_0,A_0)}\eta = \varepsilon^2 d^*_{A_0} d_{A_0}\eta + L^*_{u_0} L_{u_0}\eta.$$

By our standing hypotheses, $\mu^{-1}(0)$ is compact and $L_x : \mathfrak{g} \to T_x M$ is injective for every $x \in \mu^{-1}(0)$. Hence there exists a constant $c_1 > 0$ such that

$$c_1^{-1} |\eta| \le |L_x \eta|_z \le c_1 |\eta| \tag{52}$$

for every $x \in \mu^{-1}(0)$, every $\eta \in \mathfrak{g}$, and every $z \in \Sigma$. (Here $|\cdot|_z$ denotes the metric on M induced by J_z and ω.) Hence the operator $d^{*_\varepsilon}_{(u_0,A_0)} d_{(u_0,A_0)}$ is injective and hence, by elliptic regularity, it is bijective.

Next we prove that there exists a constant $c_2 = c_2(p, c_0) > 0$ such that, for every pair $(u_0, A_0) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$, every $\eta \in \Omega^0(\Sigma, \mathfrak{g}_P)$, and every $\varepsilon \in (0, 1]$, we have

$$\|d_{A_0}\eta\|_{L^p} \le \varepsilon \left\| d^*_{A_0} d_{A_0}\eta \right\|_{L^p} + c_2 \varepsilon^{-1} \|\eta\|_{L^p}. \tag{53}$$

For a fixed connection $A_0 \in \mathcal{A}(P)$ this follows directly from the interpolation inequality in [GT, Theorem 7.27] and the L^p-estimate for the operator $d_{A_0}^* d_{A_0}$. Now the identity

$$\begin{aligned} d_A^* d_A \eta - d_{A_0}^* d_{A_0} \eta &= [A - A_0 \wedge d_{A_0} \eta] + *[*(A - A_0) \wedge d_{A_0} \eta] \\ &\quad - *[d_{A_0} * (A - A_0), \eta] + *[*(A - A_0) \wedge [A - A_0, \eta]] \end{aligned}$$

shows that the constant in (53) depends continuously on A with respect to the C^1-norm. Moreover, the inequality (53) is gauge invariant. Hence it follows from Lemma 2.2 (with $\ell = 2$) and the Arzéla-Ascoli theorem that the estimate (53) holds with a uniform constant c_2 for all A_0 such that $(u_0, A_0) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$ for some u_0.

Using the identity

$$d\,|\eta|^{p-2} = (p-2)\,|\eta|^{p-4}\,\langle \eta, d_{A_0}\eta\rangle \in \Omega^1(\Sigma)$$

for $\eta \in \Omega^0(\Sigma, \mathfrak{g}_P)$ and integration by parts we obtain

$$\int_\Sigma |\eta|^{p-2}\,|d_{A_0}\eta|^2 = \int_\Sigma |\eta|^{p-2}\,\langle \eta, d_{A_0}^* d_{A_0}\eta\rangle - (p-2)\int_\Sigma |\eta|^{p-4}\,|\langle \eta, d_{A_0}\eta\rangle|^2.$$

The last term on the right is negative. Now (46) is equivalent to

$$\varepsilon^2 d_{A_0}^*\alpha - L_{u_0}^*\xi = \varepsilon^2 d_{A_0}^* d_{A_0}\eta + L_{u_0}^* L_{u_0}\eta.$$

Hence, by the previous identity and (52), we have

$$\begin{aligned} &\int_\Sigma \left(c_1^{-2}\,|\eta|^p + \varepsilon^2\,|\eta|^{p-2}\,|d_{A_0}\eta|^2\right) \\ &\le \int_\Sigma |\eta|^{p-2}\left(|L_{u_0}\eta|^2 + \varepsilon^2\,|d_{A_0}\eta|^2\right) \\ &\le \int_\Sigma |\eta|^{p-2}\,\langle \eta, \varepsilon^2 d_{A_0}^* d_{A_0}\eta + L_{u_0}^* L_{u_0}\eta\rangle \\ &= \int_\Sigma |\eta|^{p-2}\,\langle \eta, \varepsilon^2 d_{A_0}^*\alpha - L_{u_0}^*\xi\rangle \\ &= \int_\Sigma |\eta|^{p-2}\left(\varepsilon^2\langle d_{A_0}\eta, \alpha\rangle + \varepsilon^2(p-2)|\eta|^{-2}\langle\langle \eta, d_{A_0}\eta\rangle, \langle \eta, \alpha\rangle\rangle - \langle L_{u_0}\eta, \xi\rangle\right) \\ &\le c_1 \int_\Sigma |\eta|^{p-1}\,|\xi| + \varepsilon^2(p-1)\int_\Sigma |\eta|^{p-2}\,|d_{A_0}\eta|\,|\alpha| \\ &\le c_1 \int_\Sigma |\eta|^{p-1}\,|\xi| + \frac{\varepsilon^2(p-1)^2}{2}\int_\Sigma |\eta|^{p-2}\,|\alpha|^2 + \frac{\varepsilon^2}{2}\int_\Sigma |\eta|^{p-2}\,|d_{A_0}\eta|^2. \end{aligned}$$

Therefore, by Hölder's inequality,

$$c_1^{-2}\left\|\eta\right\|_{L^p}^p \le c_1\left\|\eta\right\|_{L^p}^{p-1}\left\|\xi\right\|_{L^p} + \frac{\varepsilon^2(p-1)^2}{2}\left\|\eta\right\|_{L^p}^{p-2}\left\|\alpha\right\|_{L^p}^2,$$

$$c_1^{-2}\left\|\eta\right\|_{L^p}^p \le \left\|\eta\right\|_{L^p}^{p-1}\left\|\tilde{\eta}\right\|_{L^p}, \qquad \tilde{\eta} := \varepsilon^2 d_{A_0}^* d_{A_0}\eta + L_{u_0}^* L_{u_0}\eta.$$

Hence

$$c_1^{-2}\left\|\eta\right\|_{L^p}^2 \le c_1\left\|\eta\right\|_{L^p}\left\|\xi\right\|_{L^p} + \frac{\varepsilon^2(p-1)^2}{2}\left\|\alpha\right\|_{L^p}^2,$$

and hence

$$\left\|\eta\right\|_{L^p}^2 \le c_1^6\left\|\xi\right\|_{L^p}^2 + c_1^2\varepsilon^2(p-1)^2\left\|\alpha\right\|_{L^p}^2.$$

Thus we have proved the inequalities

$$\left\|\eta\right\|_{L^p} \le c_1\max\{p-1, c_1^2\}\left\|\zeta\right\|_{0,p,\varepsilon}, \qquad \left\|\eta\right\|_{L^p} \le c_1^2\left\|\tilde{\eta}\right\|_{L^p}. \tag{54}$$

By (52), (53), and (54),

$$\begin{array}{rcl}\left\|\eta\right\|_{2,p,\varepsilon;A_0} & = & \left\|\eta\right\|_{L^p} + \varepsilon\left\|d_{A_0}\eta\right\|_{L^p} + \varepsilon^2\left\|d_{A_0}^* d_{A_0}\eta\right\|_{L^p} \\ & \le & (1+c_2)\left\|\eta\right\|_{L^p} + 2\varepsilon^2\left\|d_{A_0}^* d_{A_0}\eta\right\|_{L^p} \\ & \le & (1+c_2+2c_1^2)\left\|\eta\right\|_{L^p} + 2\left\|\tilde{\eta}\right\|_{L^p} \\ & \le & \left(2 + c_1^2(1+c_2+2c_1^2)\right)\left\|\tilde{\eta}\right\|_{L^p}.\end{array}$$

This proves the second estimate in (47).

To prove the first estimate in (47) we use a rescaling argument in local holomorphic coordinates on Σ. Cover Σ by finitely many open sets, each of which is holomorphically diffeomorphic to the unit square in $\mathbb{C}$, suppose that the coordinate charts extend to a closed square of side length two, and choose trivializations of the bundle P over each of these (extended) open sets. In these coordinates we write the metric in the form $\lambda^2(ds^2+dt^2)$, and we write $A := A_0 = \Phi\,ds + \Psi\,dt$, $\alpha = \varphi\,ds + \psi\,dt$. Moreover, $u := u_0 : [0,2]^2 \to M$, $\xi : [0,2]^2 \to TM$ is a vector field along u, and $\eta : [0,2]^2 \to \mathfrak{g}$. In this notation equation (46) has the form

$$\nabla_s\nabla_s\eta + \nabla_t\nabla_t\eta = \nabla_s\varphi + \nabla_t\psi + (\lambda/\varepsilon)^2 L_u^*(L_u\eta + \xi), \tag{55}$$

where $\nabla_s\eta := \nabla_{A,s}^{\mathrm{G}}\eta = \partial_s\eta + [\Phi,\eta]$ and $\nabla_t\eta := \nabla_{A,t}^{\mathrm{G}}\eta = \partial_t\eta + [\Psi,\eta]$. Now we introduce new functions, defined on the square $[0,2/\varepsilon]^2$, by

$$\begin{array}{rclrcl}\tilde{\eta}(s,t) & := & \eta(\varepsilon s,\varepsilon t), & \tilde{\lambda}(s,t) & := & \lambda(\varepsilon s,\varepsilon t), \\ \tilde{\xi}(s,t) & := & \xi(\varepsilon s,\varepsilon t), & \tilde{u}(s,t) & := & u(\varepsilon s,\varepsilon t), \\ \tilde{\varphi}(s,t) & := & \varepsilon\varphi(\varepsilon s,\varepsilon t), & \tilde{\Phi}(s,t) & := & \varepsilon\Phi(\varepsilon s,\varepsilon t), \\ \tilde{\psi}(s,t) & := & \varepsilon\psi(\varepsilon s,\varepsilon t), & \tilde{\Psi}(s,t) & := & \varepsilon\Psi(\varepsilon s,\varepsilon t).\end{array}$$

Then (55) is equivalent to

$$\tilde{\nabla}_s\tilde{\nabla}_s\tilde{\eta} + \tilde{\nabla}_t\tilde{\nabla}_t\tilde{\eta} = \tilde{\nabla}_s\tilde{\varphi} + \tilde{\nabla}_t\tilde{\psi} + \tilde{\lambda}^2 L_{\tilde{u}}^*(L_{\tilde{u}}\tilde{\eta} + \tilde{\xi}),$$

where $\tilde{\nabla}_s\tilde{\eta} := \partial_s\tilde{\eta} + [\tilde{\Phi}, \tilde{\eta}]$ and $\tilde{\nabla}_t\tilde{\eta} := \partial_t\tilde{\eta} + [\tilde{\Psi}, \tilde{\eta}]$. This equation can be written in the form

$$\Delta\tilde{\eta} = \partial_s\tilde{f} + \partial_t\tilde{g} + \tilde{h}$$

where $\Delta := \partial_s\partial_s + \partial_t\partial_t$ and $\tilde{f}, \tilde{g}, \tilde{h} : [0, 2/\varepsilon]^2 \to \mathfrak{g}$ are given by

$$\tilde{f} := \tilde{\varphi} - 2[\tilde{\Phi}, \tilde{\eta}], \qquad \tilde{g} := \tilde{\psi} - 2[\tilde{\Psi}, \tilde{\eta}],$$

$$\tilde{h} := \tilde{\lambda}^2 L_{\tilde{u}}^*(L_{\tilde{u}}\tilde{\eta} + \tilde{\xi}) + [\tilde{\Phi}, \tilde{\varphi} - [\tilde{\Phi}, \tilde{\eta}]] + [\tilde{\Psi}, \tilde{\psi} - [\tilde{\Psi}, \tilde{\eta}]] + [\partial_s\tilde{\Phi} + \partial_t\tilde{\Psi}, \tilde{\eta}].$$

Hence there exists a constant $c_3 > 0$ such that, for all real numbers a, b such that $1/2 \le a < b \le 2/\varepsilon - 1/2$, we have

$$\int_{[a,b]^2} \left(|\tilde{\nabla}_s\tilde{\eta}|^p + |\tilde{\nabla}_t\tilde{\eta}|^p\right) \le c_3 \int_{[a-1/2,b+1/2]^2} \left(|\tilde{f}|^p + |\tilde{g}|^p + |\tilde{h}|^p + |\tilde{\eta}|^p\right).$$

Here the constant c_3 is independent of a and b. It follows that

$$\begin{aligned}\int_{[a,b]^2} \left(|\tilde{\nabla}_s\tilde{\eta}|^p + |\tilde{\nabla}_t\tilde{\eta}|^p\right)\tilde{\lambda}^{2-p} &\le c_4 \int_{[a-1/2,b+1/2]^2} \left(|\tilde{\varphi}|^p + |\tilde{\psi}|^p\right)\tilde{\lambda}^{2-p} \\ &\quad + c_4 \int_{[a-1/2,b+1/2]^2} \left(|\tilde{\xi}|^p + |\tilde{\eta}|^p\right)\tilde{\lambda}^2,\end{aligned}$$

where the constant c_4 depends on the metric and on the C^1-norms of $\tilde{\Phi}$ and $\tilde{\Psi}$. With $a = 1/2\varepsilon$, $b = 3/2\varepsilon$, and $0 < \varepsilon \le 1$ we obtain

$$\begin{aligned}\varepsilon^p \int_{[1/2,3/2]^2} \left(|\nabla_s\eta|^p + |\nabla_t\eta|^p\right)\lambda^{2-p} &\le c_4\varepsilon^p \int_{[0,2]^2} \left(|\varphi|^p + |\psi|^p\right)\lambda^{2-p} \\ &\quad + c_4 \int_{[0,2]^2} \left(|\xi|^p + |\eta|^p\right)\lambda^2.\end{aligned}$$

Hence, by taking the sum over the coordinate charts,

$$\varepsilon \left\|d_{A_0}\eta\right\|_{L^p} \le N^{1/p} c_4 \left(\varepsilon \left\|\alpha\right\|_{L^p} + \left\|\xi\right\|_{L^p} + \left\|\eta\right\|_{L^p}\right).$$

Here N is the number of open sets in the cover and the constant c_4 depends continuously on A_0 with respect to the C^1-norm. Hence, by Lemma 2.2, c_4 can be chosen independent of the pair $(u_0, A_0) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$. Combining the last inequality with (54) we obtain the first estimate in (47) as claimed. □

In the following proof we use the identity

$$(e^{\eta})^*A - A - d_A\eta = \sum_{k=1}^{\infty} \frac{(-1)^k}{(k+1)!}\mathrm{ad}(\eta)^k d_A\eta \tag{56}$$

for $A \in \mathcal{A}(P)$ and $\eta \in \Omega^0(\Sigma, \mathfrak{g}_P)$, where $\mathrm{ad}(\eta)\alpha := [\eta, \alpha]$ for $\alpha \in \Omega^1(\Sigma, \mathfrak{g}_P)$.

Proof of Lemma 7.3. Throughout the proof we denote by $c_1, c_2, c_3, \ldots$ positive constants depending only on p and c_0 (and not on the pair (u_0, A_0)). Fix a pair $(u_0, A_0) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$ and choose a positive constant δ_0 that is smaller than the injectivity radius of M on the compact set $u_0(P)$. Suppose that $\eta \in \Omega^0(\Sigma, \mathfrak{g}_P)$ and $\zeta_0 = (\xi_0, \alpha_0) \in T_{(u_0,A_0)}\mathcal{B}$ satisfy the hypotheses of Lemma 7.3 with a sufficiently small constant $\delta > 0$. Let c_1 be the constant of Lemma 4.4. Then, by (48),

$$\|\xi_0\|_{L^\infty} + \|\eta\|_{L^\infty} \le c_1\varepsilon^{-2/p}\left(\|\xi_0\|_{1,p,\varepsilon} + \|\eta\|_{2,p,\varepsilon}\right) \le c_1\delta.$$

Hence, if δ is sufficiently small, it follows that the C^0-distance between u_0 and $e^{-r\eta}\exp_{u_0}(\xi_0)$ is smaller than δ_0 for every $r \in [0,1]$. Hence there exists a unique smooth path $[0,1] \to T_{(u_0,A_0)}\mathcal{B} : r \mapsto \zeta_r = (\xi_r, \alpha_r)$ starting at ζ_0 such that

$$(\exp_{u_0}(\xi_r), A_0 + \alpha_r) = g^*(u, A),$$

where

$$(u, A) := (\exp_{u_0}(\xi_0), A_0 + \alpha_0), \qquad g := e^{r\eta}. \tag{57}$$

The endpoint ζ_1 of this path obviously satisfies (49). We prove the inequalities

$$\|\partial_r\zeta_r\|_{0,p,\varepsilon} \le c\,\|\eta\|_{1,p,\varepsilon}\,, \qquad \|\partial_r\zeta_r\|_{1,p,\varepsilon} \le c\,\|\eta\|_{2,p,\varepsilon}\,, \tag{58}$$

$$\left\| d^{*_\varepsilon}_{(u_0,A_0)}\left(\partial_r\zeta_r - d_{(u_0,A_0)}\eta\right)\right\|_{L^p} \le c\varepsilon^{-2/p}\left(\|\zeta_0\|_{1,p,\varepsilon} + \|\eta\|_{2,p,\varepsilon}\right)\|\eta\|_{1,p,\varepsilon} \tag{59}$$

for $0 \le r \le 1$, where the constant c depends only on c_0 and p. Then the inequalities (50) and (51) follow by integrating the function $r \mapsto \partial_r\zeta_r$ over the interval $0 \le r \le 1$.

For every $u \in C^\infty_{\mathrm{G}}(P,M)$ whose C^0-distance to u_0 is less than δ_0 we define the linear operator $Z(u) : \Omega^0(\Sigma, \mathfrak{g}_P) \to \Omega^0(\Sigma, u_0^*TM/\mathrm{G})$ by

$$Z(u)\hat{\eta} := d\exp_{u_0}^{-1}(u)L_u\hat{\eta}$$

for $\hat{\eta} \in \Omega^0(\Sigma, \mathfrak{g}_P)$. Then

$$\partial_r\zeta_r = (-Z(g^{-1}u)\eta, d_{g^*A}\eta_0), \tag{60}$$

where (u,A) and g are as in (57). We prove the first inequality in (58). Since $A = A_0 + \alpha_0$ we have

$$d_{g^*A}\eta = d_{A_0}\eta + [g^*A_0 - A_0, \eta] + [g^{-1}\alpha_0 g, \eta] \tag{61}$$

and we must estimate the three terms on the right with $g := e^{r\eta}$. Since $\|\eta\|_{L^\infty} \le c_1\delta$ it follows from (56), with η replaced by $r\eta$, that

$$\|[g^*A_0 - A_0, \eta]\|_{L^p} \le c_2 \|d_{A_0}\eta\|_{L^p} \|\eta\|_{L^\infty} \le c_3\varepsilon^{-1-2/p} \|\eta\|_{1,p,\varepsilon}^2 .$$

Moreover,

$$\left\|[g^{-1}\alpha_0 g, \eta]\right\|_{L^p} \le \varepsilon^{-1-2/p} \|\zeta_0\|_{0,p,\varepsilon} \|\eta\|_{1,p,\varepsilon} .$$

Hence, by (48) and (61), $\|d_{g^*A}\eta\|_{L^p} \le c_4\varepsilon^{-1}\|\eta\|_{1,p,\varepsilon}$ and hence the first inequality in (58) follows from (60).

Next we prove the second inequality in (58) and (59). Using the identity

$$d_{A_0}[g^{-1}\alpha_0 g, \eta] = [g^{-1}(d_{A_0}\alpha_0)g, \eta] + [[(A_0 - g^*A_0)\wedge g^{-1}\alpha_0 g], \eta] - [g^{-1}\alpha_0 g \wedge d_{A_0}\eta]$$

and (56) we obtain

$$\begin{aligned} \left\|[g^{-1}\alpha_0 g, \eta]\right\|_{1,p,\varepsilon} &= \left\|[g^{-1}\alpha_0 g, \eta]\right\|_{L^p} \\ &\quad + \varepsilon \left\|d_{A_0}[g^{-1}\alpha_0 g, \eta]\right\|_{L^p} + \varepsilon \left\|d_{A_0}[*g^{-1}\alpha_0 g, \eta]\right\|_{L^p} \\ &\le c_5\varepsilon^{-1-2/p} \|\zeta_0\|_{1,p,\varepsilon} \|\eta\|_{1,p,\varepsilon} . \end{aligned}$$

Similarly, using the identity

$$d_{A_0}[(A_0 - g^*A_0), \eta] = [d_{A_0}(A_0 - g^*A_0), \eta] - [(A_0 - g^*A_0) \wedge d_{A_0}\eta]$$

and (56) we obtain

$$\|[(A_0 - g^*A_0), \eta]\|_{1,p,\varepsilon} \le c_6\varepsilon^{-1-2/p} \|\eta\|_{2,p,\varepsilon} \|\eta\|_{1,p,\varepsilon} .$$

Hence, by (61),

$$\left\|d_{g^*A}\eta - d_{A_0}\eta\right\|_{1,p,\varepsilon} \le c_7\varepsilon^{-1-2/p}\left(\|\zeta_0\|_{1,p,\varepsilon} + \|\eta\|_{2,p,\varepsilon}\right)\|\eta\|_{1,p,\varepsilon}.$$

Moreover, since $Z(u_0) = L_{u_0}$, we have

$$\begin{aligned}
&\left\|Z(g^{-1}u)\eta - L_{u_0}\eta\right\|_{1,p,\varepsilon} \\
&\le \left\|Z(g^{-1}u)\eta - Z(u)\eta\right\|_{1,p,\varepsilon} + \|Z(u)\eta - Z(u_0)\eta\|_{1,p,\varepsilon} \\
&\le c_8\left(\|\eta\|_{1,p,\varepsilon}\|\eta\|_{L^\infty} + \|\xi_0\|_{1,p,\varepsilon}\|\eta\|_{L^\infty} + \|\xi_0\|_{L^\infty}\|\eta\|_{1,p,\varepsilon}\right) \\
&\le c_9\varepsilon^{-2/p}\left(\|\zeta_0\|_{1,p,\varepsilon} + \|\eta\|_{2,p,\varepsilon}\right)\|\eta\|_{1,p,\varepsilon}.
\end{aligned}$$

Here we have used the inequality $\|\eta\|_{L^\infty} \le c_1\delta$ from (48) and Lemma 4.4. The constants c_7 and c_9 in the last two estimates depend continuously on the pair (u_0, A_0) with respect to the C^1-norm and are gauge invariant. Hence, by Lemma 2.2, they can be chosen independent of $(u_0, A_0) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$. Hence the second inequality in (58) follows from (60). To prove (59) we observe that

$$\begin{aligned}
&d^{*_\varepsilon}_{(u_0,A_0)}\partial_r\zeta_r - d^{*_\varepsilon}_{(u_0,A_0)}d_{(u_0,A_0)}\eta \\
&= \varepsilon^2 d^*_{A_0}\left(d_{g^*A}\eta - d_{A_0}\eta\right) + L^*_{u_0}\left(Z(g^{-1}u)\eta - L_{u_0}\eta\right),
\end{aligned}$$

where (u, A) and $g := e^{r\eta}$ are as in (57). The terms on the right have been estimated above and this proves (59). Thus we have proved the existence of ζ_1. The inequality (50) with δ sufficiently small guarantees that the C^0 distance between u_0 and $\exp_{u_0}(\xi_1)$ is smaller than the injectivity radius. This proves uniqueness. □

Proof of Proposition 7.1. The proof is based on a Newton type iteration. Let

$$(u_1, A_1) := (u, A) = (\exp_{u_0}(\xi), A_0 + \alpha), \qquad \zeta_1 := \zeta.$$

For $\nu \ge 2$ we define $\zeta_\nu = (\xi_\nu, \alpha_\nu) \in T_{(u_0,A_0)}\mathcal{B}$ inductively by

$$(\exp_{u_0}(\xi_{\nu+1}), A_0 + \alpha_{\nu+1}) := (u_{\nu+1}, A_{\nu+1}) := g_\nu^*(u_\nu, A_\nu),$$

where $(u_\nu, A_\nu) := (\exp_{u_0}(\xi_\nu), A_0 + \alpha_\nu)$, $g_\nu := e^{\hat\eta_\nu}$, and $\hat\eta_\nu \in \Omega^0(\Sigma, \mathfrak{g}_P)$ is the unique solution of the equation

$$d^{*_\varepsilon}_{(u_0,A_0)}d_{(u_0,A_0)}\hat\eta_\nu + d^{*_\varepsilon}_{(u_0,A_0)}\zeta_\nu = 0.$$

To construct these sequences we must ensure that in each step ζ_ν and $\hat{\eta}_\nu$ satisfy the hypotheses of Lemma 7.3 so that $\zeta_{\nu+1}$ can be chosen as in the assertion of Lemma 7.3. We shall prove this below. And we shall also prove that these sequences satisfy the following estimates.

$$\begin{aligned}
\left\|\zeta_\nu\right\|_{1,p,\varepsilon} &\leq C\left\|\zeta\right\|_{1,p,\varepsilon}, && (62)\\
\left\|d^{*_\varepsilon}_{(u_0,A_0)}\zeta_\nu\right\|_{L^p} &\leq C\varepsilon^{-2/p}\left\|\zeta_{\nu-1}\right\|_{1,p,\varepsilon}\left\|d^{*_\varepsilon}_{(u_0,A_0)}\zeta_{\nu-1}\right\|_{L^p}, && (63)\\
\left\|d^{*_\varepsilon}_{(u_0,A_0)}\zeta_\nu\right\|_{L^p} &\leq 2^{1-\nu}\left\|d^{*_\varepsilon}_{(u_0,A_0)}\zeta\right\|_{L^p}, && (64)\\
\left\|\hat{\eta}_\nu\right\|_{2,p,\varepsilon} &\leq C2^{-\nu}\left\|\zeta\right\|_{1,p,\varepsilon}, && (65)
\end{aligned}$$

The constants C and δ are chosen as follows. Suppose that the constants $c_1, c_2, c_3, c_4, c_5 \geq 1$ and $\delta_0, \delta_3, \delta_4 \in (0,1]$ satisfy the following conditions.

- The injectivity radius of M on $u_0(P)$ is bigger than δ_0.
- The inequality (52) holds with c_1 for every $x \in \mu^{-1}(0)$.
- The assertion of Lemma 7.2 holds for $0 < \varepsilon \leq 1$ with c replaced by c_2.
- The assertion of Lemma 7.3 holds for $0 < \varepsilon \leq 1$ with c replaced by c_3 and δ replaced by δ_3.
- The assertion of Lemma 7.4 holds for $0 < \varepsilon \leq 1$ with c replaced by c_4 and δ replaced by δ_4.
- The assertion of Lemma 4.4 holds for $0 < \varepsilon \leq 1$ with c replaced by c_5.

Now choose positive constants C and δ such that

$$c_1c_2(1+2c_2)c_3 \leq C, \quad 2c_2(1+c_2)c_3C\delta \leq 1, \quad 2C\delta \leq \delta_3, \quad 4c_4c_5C\delta \leq \delta_4.$$

We prove that the estimates (62-65) hold for $\nu = 1$. Since $C/2 \geq c_1c_2$, the inequality (65) with $\nu = 1$ follows from Lemma 7.2. Since $C \geq 1$, the inequality (62) holds for $\nu = 1$. The inequality (63) is vacuous for $\nu = 1$ and (64) is obvious.

Now suppose that the sequences have been constructed and the inequalities (62-65) have been established up to some integer $\nu \geq 1$. Then

$$\left\|\hat{\eta}_\nu\right\|_{2,p,\varepsilon} + \left\|\zeta_\nu\right\|_{1,p,\varepsilon} \leq C(1+2^{-\nu})\left\|\zeta\right\|_{1,p,\varepsilon} \leq 2C\delta\varepsilon^{2/p} \leq \delta_3\varepsilon^{2/p}.$$

Hence the hypotheses of Lemma 7.3 are satisfied with ζ_0 replaced by ζ_ν and η replaced by $\hat{\eta}_\nu$. Choose $\zeta_{\nu+1} = (\xi_{\nu+1}, \alpha_{\nu+1})$ as in the assertion of Lemma 7.3. By Lemma 7.2, we have

$$\|\hat{\eta}_\nu\|_{1,p,\varepsilon} \le c_2 \|\zeta_\nu\|_{0,p,\varepsilon}, \qquad \|\hat{\eta}_\nu\|_{2,p,\varepsilon} \le c_2 \left\| d^{*_\varepsilon}_{(u_0,A_0)} \zeta_\nu \right\|_{L^p}.$$

Moreover, $d^{*_\varepsilon}_{(u_0,A_0)}(\zeta_\nu + d_{(u_0,A_0)}\eta_\nu) = 0$, and hence, by (51),

$$\begin{aligned} \left\| d^{*_\varepsilon}_{(u_0,A_0)} \zeta_{\nu+1} \right\|_{L^p} &\le c_3 \varepsilon^{-2/p} \left(\|\zeta_\nu\|_{1,p,\varepsilon} + \|\hat{\eta}_\nu\|_{2,p,\varepsilon} \right) \|\hat{\eta}_\nu\|_{1,p,\varepsilon} \\ &\le c_2(1+c_2)c_3 \varepsilon^{-2/p} \|\zeta_\nu\|_{1,p,\varepsilon} \left\| d^{*_\varepsilon}_{(u_0,A_0)} \zeta_\nu \right\|_{L^p}. \end{aligned}$$

Since $c_2(1+c_2)c_3 \le C$, this proves (63) with ν replaced by $\nu+1$. Moreover, by (62),

$$\begin{aligned} \left\| d^{*_\varepsilon}_{(u_0,A_0)} \zeta_{\nu+1} \right\|_{L^p} &\le c_2(1+c_2)c_3 C \varepsilon^{-2/p} \|\zeta\|_{1,p,\varepsilon} \left\| d^{*_\varepsilon}_{(u_0,A_0)} \zeta_\nu \right\|_{L^p} \\ &\le c_2(1+c_2)c_3 C\delta \left\| d^{*_\varepsilon}_{(u_0,A_0)} \zeta_\nu \right\|_{L^p}. \end{aligned}$$

Since $2c_2(1+c_2)c_3C\delta \le 1$, this proves (64) with ν replaced by $\nu+1$. Now let $\hat{\eta}_{\nu+1}$ be the unique solution of $d^*_{(u_0,A_0)} d_{(u_0,A_0)} \hat{\eta}_{\nu+1} + d^*_{(u_0,A_0)} \zeta_{\nu+1} = 0$. Then, by Lemma 7.2 and (64),

$$\|\hat{\eta}_{\nu+1}\|_{2,p,\varepsilon} \le c_2 \left\| d^{*_\varepsilon}_{(u_0,A_0)} \zeta_{\nu+1} \right\|_{L^p} \le c_2 2^{-\nu} \left\| d^{*_\varepsilon}_{(u_0,A_0)} \zeta \right\|_{L^p} \le c_1 c_2 2^{-\nu} \|\zeta\|_{1,p,\varepsilon}.$$

Since $2c_1c_2 \le C$ this implies (65) with ν replaced by $\nu+1$. It remains to prove (62) with ν replaced by $\nu+1$. By (50) and (64), we have

$$\|\zeta_{j+1} - \zeta_j\|_{1,p,\varepsilon} \le c_3 \|\hat{\eta}_j\|_{2,p,\varepsilon} \le c_2 c_3 2^{1-j} \left\| d^{*_\varepsilon}_{(u_0,A_0)} \zeta \right\|_{L^p} \le c_1 c_2 c_3 2^{1-j} \|\zeta\|_{1,p,\varepsilon} \tag{66}$$

for $j = 1, \ldots, \nu$. Hence

$$\|\zeta_{\nu+1}\|_{1,p,\varepsilon} \le \|\zeta\|_{1,p,\varepsilon} + \sum_{j=1}^{\nu} \|\zeta_{j+1} - \zeta_j\|_{1,p,\varepsilon} \le (1 + 2c_1c_2c_3) \|\zeta\|_{1,p,\varepsilon}.$$

Since $1 + 2c_1c_2c_3 \le C$ this proves (62) with ν replaced by $\nu+1$. This completes the induction.

By (66), ζ_ν is a Cauchy sequence in the $W^{1,p}$-norm. Moreover,

$$(u_\nu, A_\nu) = \bigl(\exp_{u_0}(\xi_\nu), A_0 + \alpha_\nu\bigr) = h_\nu^*(u, A),$$

where $h_\nu := g_1 g_2 \cdots g_{\nu-1}$. We prove by induction that there exists a sequence $\eta_\nu \in \Omega^0(\Sigma, \mathfrak{g}_P)$ such that

$$h_\nu = e^{\eta_\nu}, \qquad \|\eta_{\nu+1} - \eta_\nu\|_{2,p,\varepsilon} \le c_4 C 2^{-\nu} \|\zeta\|_{1,p,\varepsilon}. \tag{67}$$

For $\nu = 1$ we set $h_1 := \mathbb{1}$ and $\eta_1 := 0$. Suppose that the sequence has been constructed for all integers up to $\nu \ge 1$. Then

$$\|\eta_\nu\|_{2,p,\varepsilon} \le \sum_{j=1}^{\nu-1} \|\eta_{j+1} - \eta_j\|_{2,p,\varepsilon} \le c_4 C \|\zeta\|_{1,p,\varepsilon}. \tag{68}$$

Hence, by Lemma 4.4, (44), and (65),

$$\|\eta_\nu\|_{L^\infty} \le c_5 \varepsilon^{-2/p} \|\eta_\nu\|_{2,p,\varepsilon} \le c_4 c_5 C \varepsilon^{-2/p} \|\zeta\|_{1,p,\varepsilon} \le c_4 c_5 C \delta \le \delta_4/4,$$

$$\|\hat{\eta}_\nu\|_{L^\infty} \le c_5 \varepsilon^{-2/p} \|\hat{\eta}_\nu\|_{2,p,\varepsilon} \le c_5 C \varepsilon^{-2/p} \|\zeta\|_{1,p,\varepsilon} \le c_5 C \delta \le \delta_4/4.$$

By Lemma 7.4, there exists a section $\eta_{\nu+1} \in \Omega^0(\Sigma, \mathfrak{g}_P)$ such that

$$e^{\eta_{\nu+1}} = e^{\eta_\nu} e^{\hat{\eta}_\nu} = h_\nu g_\nu = h_{\nu+1}, \qquad \|\eta_{\nu+1}\|_{L^\infty} \le 2 \|\eta_\nu + \hat{\eta}_\nu\|_{L^\infty} \le \delta_4.$$

Applying Lemma 7.4 to $-\eta_\nu$ and $\eta_{\nu+1}$ we find

$$\|\eta_{\nu+1} - \eta_\nu\|_{2,p,\varepsilon} \le c_4 \|\hat{\eta}_\nu\|_{2,p,\varepsilon} \le c_4 C 2^{-\nu} \|\zeta\|_{1,p,\varepsilon}.$$

The last inequality follows from (65). This completes the induction. Thus we have proved that h_ν satisfies (67) and hence is a Cauchy sequence in $\mathcal{G}^{2,p}(P)$. Denote

$$\zeta := \lim_{\nu\to\infty} \zeta_\nu, \qquad h := \lim_{\nu\to\infty} h_\nu, \qquad \eta := \lim_{\nu\to\infty} \eta_\nu.$$

Then

$$e^\eta = h, \qquad h^*(u, A) = (\exp_{u_0}(\xi), A_0 + \alpha), \qquad d_{(u_0,A_0)}^{*_\varepsilon} \zeta = 0.$$

The last equation follows from (64). Moreover, by (62) and (68), we have $\|\eta\|_{2,p,\varepsilon} + \|\zeta\|_{1,p,\varepsilon} \le C(1 + c_4) \|\zeta\|_{1,p,\varepsilon}$. Hence (45) holds with $c := C(1 + c_4)$.

To complete the existence proof we must show that η and ζ are smooth. We shall prove that the sequence ζ_ν is bounded on $W^{k,p}$ for every k. Here it suffices to obtain rather crude estimates with constants which depend on ε and are allowed to diverge as ε tends to zero. We fix a constant $\varepsilon > 0$ and prove by induction that for every integer $k \geq 1$ there exists a constant $c_k = c_k(p, \varepsilon, u_0, A_0, u, A)$ such that, for every ν,

$$\|\zeta_\nu\|_{W^{k,p}} \leq c_k, \qquad \|\hat{\eta}_\nu\|_{W^{k+1,p}} \leq c_k 2^{-\nu}. \tag{69}$$

For $k = 1$ this follows from (62) and (65). Now let $k \geq 2$ and assume that these estimates have been established with k replaced by $k-1$. Observe that there exists a constant $C_k \geq 1$ such that, for every ν,

$$\begin{aligned} \|\zeta_{\nu+1} - \zeta_\nu\|_{W^{k,p}} &\leq C_k \|\hat{\eta}_\nu\|_{W^{k+1,p}}, \\ \|\hat{\eta}_\nu\|_{W^{k+1,p}} &\leq C_k \left\| d^{*\varepsilon}_{(u_0,A_0)} \zeta_\nu \right\|_{W^{k-1,p}}, \\ \left\| d^{*\varepsilon}_{(u_0,A_0)} \zeta_{\nu+1} \right\|_{W^{k-1,p}} &\leq C_k \left(\|\zeta_\nu\|_{W^{k,p}} + \|\hat{\eta}_\nu\|_{W^{k+1,p}} \right) \|\hat{\eta}_\nu\|_{W^{k,p}}. \end{aligned}$$

The first two inequalities are obvious, and the last follows by inspecting the formula (60) in the proof of Lemma 7.3. Combining these inequalities with the induction hypothesis, we obtain

$$\begin{aligned} \|\zeta_{\nu+1}\|_{W^{k,p}} &\leq \|\zeta_\nu\|_{W^{k,p}} + C_k \|\hat{\eta}_\nu\|_{W^{k+1,p}}, \\ \|\hat{\eta}_{\nu+1}\|_{W^{k+1,p}} &\leq C_k^2 c_{k-1} \left(\|\zeta_\nu\|_{W^{k,p}} + \|\hat{\eta}_\nu\|_{W^{k+1,p}} \right) 2^{-\nu}. \end{aligned}$$

Abbreviate

$$a_\nu := \|\zeta_{\nu+\nu_0}\|_{W^{k,p}} + C_k \|\hat{\eta}_{\nu+\nu_0}\|_{W^{k+1,p}}$$

and choose ν_0 so large that $C_k^3 c_{k-1} 2^{-\nu_0} \leq 1$. Then

$$\begin{aligned} a_{\nu+1} &\leq \|\zeta_{\nu+\nu_0}\|_{W^{k,p}} + C_k \|\hat{\eta}_{\nu+\nu_0}\|_{W^{k+1,p}} + C_k \|\hat{\eta}_{\nu+\nu_0+1}\|_{W^{k+1,p}} \\ &= a_\nu + C_k \|\hat{\eta}_{\nu+\nu_0+1}\|_{W^{k+1,p}} \\ &\leq a_\nu + C_k^3 c_{k-1} \left(\|\zeta_\nu\|_{W^{k,p}} + \|\hat{\eta}_\nu\|_{W^{k+1,p}} \right) 2^{-\nu-\nu_0} \\ &\leq (1 + 2^{-\nu}) a_\nu \end{aligned}$$

for all ν and hence the sequence a_ν is bounded. It follows that the sequences $\|\zeta_\nu\|_{W^{k,p}}$ and $2^\nu \|\hat{\eta}_\nu\|_{W^{k+1,p}}$ are bounded. Thus we have proved that $\hat{\eta}_\nu$ and ζ_ν satisfy (69). This completes the induction. It follows that ζ is smooth and hence, so is η. This proves existence.

We prove uniqueness. Choose $\delta > 0$ so small that

$$c_5 c\delta \le \delta_0, \qquad 2c_4 c\delta \le \delta_3, \qquad c_5 c\delta \le \delta_4, \qquad 2c_2c_3c_4c\delta < 1.$$

Assume that $\zeta_0, \zeta_1 \in T_{(u_0,A_0)}\mathcal{B}$ and $\eta_0, \eta_1 \in \Omega^0(\Sigma, \mathfrak{g}_P)$ satisfy the requirements of the proposition. Then

$$d^{*_\varepsilon}_{(u_0,A_0)}\zeta_i = 0, \qquad g_i^*(u, A) = (\exp_{u_0}(\xi_i), A_0 + \alpha_i),$$

for $i = 0, 1$, where $g_i := e^{\eta_i}$. By Lemma 4.4, we have

$$\|\eta_i\|_{L^\infty} \le c_5\varepsilon^{-2/p}\|\eta_i\|_{2,p,\varepsilon} \le c_5 c\varepsilon^{-2/p}\|\zeta\|_{1,p,\varepsilon} \le c_5 c\delta \le \delta_4$$

for $i = 0, 1$. Hence, by Lemma 7.4, there exists a unique element $\eta \in \Omega^0(\Sigma, \mathfrak{g}_P)$ such that

$$g := e^\eta = g_0^{-1}g_1, \qquad c_4^{-1}\|\eta_1 - \eta_0\|_{2,p,\varepsilon} \le \|\eta\|_{2,p,\varepsilon} \le c_4\|\eta_1 - \eta_0\|_{2,p,\varepsilon}.$$

The gauge transformation g satisfies

$$g^*(\exp_{u_0}(\xi_0), A_0 + \alpha_0) = (\exp_{u_0}(\xi_1), A_0 + \alpha_1).$$

Moreover,

$$\|\zeta_0\|_{1,p,\varepsilon} + \|\eta\|_{2,p,\varepsilon} \le 2c_4 c\|\zeta\|_{1,p,\varepsilon} \le 2c_4 c\delta\varepsilon^{2/p} \le \delta_3\varepsilon^{2/p}.$$

Hence ζ_0 and η satisfy the hypotheses of Lemma 7.3. We use Lemma 7.2 and the estimate (51) of Lemma 7.3 to obtain

$$\begin{aligned}
\|\eta\|_{2,p,\varepsilon} &\le c_2\left\|d^{*_\varepsilon}_{(u_0,A_0)}d_{(u_0,A_0)}\eta\right\|_{L^p} \\
&\le c_2c_3\varepsilon^{-2/p}\left(\|\zeta_1 - \zeta_0\|_{1,p,\varepsilon} + \|\eta\|_{2,p,\varepsilon}\right)\|\eta\|_{1,p,\varepsilon} \\
&\le c_2c_3c_4\varepsilon^{-2/p}\left(\|\zeta_1 - \zeta_0\|_{1,p,\varepsilon} + \|\eta_1 - \eta_0\|_{2,p,\varepsilon}\right)\|\eta\|_{1,p,\varepsilon} \\
&\le 2c_2c_3c_4c\varepsilon^{-2/p}\|\zeta\|_{1,p,\varepsilon}\|\eta\|_{1,p,\varepsilon} \\
&\le 2c_2c_3c_4c\delta\|\eta\|_{1,p,\varepsilon}.
\end{aligned}$$

Since $2c_2c_3c_4c\delta < 1$ we have $\eta = 0$ and hence $\eta_1 = \eta_0$. Hence $\alpha_0 = \alpha_1$ and $\exp_{u_0}(\xi_0) = \exp_{u_0}(\xi_1)$. By (44), (45), and Lemma 4.4, we have

$$\|\xi_i\|_{L^\infty} \le c_5\varepsilon^{-2/p}\|\zeta_i\|_{1,p,\varepsilon} \le c_5 c\varepsilon^{-2/p}\|\zeta\|_{1,p,\varepsilon} \le c_5 c\delta \le \delta_0$$

for $i = 0, 1$. Hence $\xi_0 = \xi_1$. □

8 Proof of Theorem C

In this section we prove that the map $\widetilde{\mathcal{T}}^\varepsilon : \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0) \to \widetilde{\mathcal{M}}^\varepsilon_{B,\Sigma}$ introduced in Theorem 6.1 is locally surjective. This is the content of Theorem C and is restated more precisely as follows.

Theorem 8.1. *Assume* $(H1)$ *and* $(H4)$ *and let* $\bar{B} \in H_2(\bar{M};\mathbb{Z})$ *be a nontorsion homology class. Then, for every* $c_0 > 0$ *and every* $p > 2$, *there exist positive constants* ε_0 *and* δ *such that the following holds for every* $\varepsilon \in (0, \varepsilon_0]$. *If* $(\bar{u}_0, \bar{A}_0) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0 - 1)$ *and* $(u, A) = (\exp_{\bar{u}_0}(\bar{\xi}), \bar{A}_0 + \bar{\alpha}) \in \widetilde{\mathcal{M}}^\varepsilon_{B,\Sigma}$ *where* $\bar{\zeta} = (\bar{\xi}, \bar{\alpha}) \in T_{(\bar{u}_0,\bar{A}_0)}\mathcal{B}$ *satisfies*

$$\left\|\bar{\zeta}\right\|_{1,p,\varepsilon;(\bar{u}_0,\bar{A}_0)} \le \delta\varepsilon^{2/p+1/2},$$

then there exist $\bar{\xi}_0 \in \ker \mathcal{D}^0_{(\bar{u}_0,\bar{A}_0)}$ *and* $\eta_0 \in \Omega^0(\Sigma, \mathfrak{g}_P)$ *such that*

$$g^*(u, A) = \widetilde{\mathcal{T}}^\varepsilon(u_0, A_0), \qquad g := e^{\eta_0}, \qquad (u_0, A_0) := \mathcal{F}^0_{(\bar{u}_0,\bar{A}_0)}(\bar{\xi}_0),$$

$$\left\|\bar{\zeta}_0\right\|_{W^{1,p}} + \left\|\eta_0\right\|_{2,p,\varepsilon;A_0} \le c\left\|\bar{\zeta}\right\|_{1,p,\varepsilon;(\bar{u}_0,\bar{A}_0)}.$$

Here $\mathcal{F}^0_{(\bar{u}_0,\bar{A}_0)}$ is the map of Theorem 2.1. The proof of Theorem 8.1 is based on Corollary 6.3 and on the construction of a tubular neighbourhood of the moduli space $\mathcal{M}^0_{B,\Sigma}(c_0)$ in the quotient $\mathcal{B}/\mathcal{G}$.

Proposition 8.2. *Assume* $(H1)$ *and* $(H4)$ *and let* $\bar{B} \in H_2(\bar{M};\mathbb{Z})$ *be a nontorsion homology class. For every* $p > 2$ *and every* $c_0 > 0$ *there exist positive constants* δ, ε_0, *and* c *such that, for every* $\varepsilon \in (0, \varepsilon_0]$, *the following holds. Let* $(\bar{u}_0, \bar{A}_0) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0 - 1)$ *and* $(u, A) = (\exp_{\bar{u}_0}(\bar{\xi}), \bar{A}_0 + \bar{\alpha})$, *where the pair* $\bar{\zeta} := (\bar{\xi}, \bar{\alpha}) \in T_{(\bar{u}_0,\bar{A}_0)}\mathcal{B}$ *satisfies*

$$\left\|\bar{\zeta}\right\|_{1,p,\varepsilon;(\bar{u}_0,\bar{A}_0)} \le \delta\varepsilon^{2/p}. \tag{70}$$

Then there exist $\bar{\xi}_0 \in \ker \mathcal{D}^0_{(\bar{u}_0,\bar{A}_0)}$, $\eta_0 \in \Omega^0(\Sigma, \mathfrak{g}_P)$, $(u_0, A_0) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$, *and* $\zeta_0 = (\xi_0, \alpha_0) \in T_{(u_0,A_0)}\mathcal{B}$, *such that*

$$g^*(u, A) = (\exp_{u_0}(\xi_0), A_0 + \alpha_0), \qquad (u_0, A_0) = \mathcal{F}^0_{(\bar{u}_0,\bar{A}_0)}(\bar{\xi}_0), \tag{71}$$

where $g := e^{\eta_0}$, *and*

$$d^{*_\varepsilon}_{(u_0,A_0)}\zeta_0 = 0, \qquad \zeta_0 \in \mathrm{im}\,(\mathcal{D}^\varepsilon_{(u_0,A_0)})^*, \tag{72}$$

$$\left\|\bar{\xi}_0\right\|_{W^{1,p}} + \left\|\eta_0\right\|_{2,p,\varepsilon;\bar{A}_0} + \left\|\zeta_0\right\|_{1,p,\varepsilon;(u_0,A_0)} \le c\left\|\bar{\zeta}\right\|_{1,p,\varepsilon;(\bar{u}_0,\bar{A}_0)}. \tag{73}$$



Figure 1: A tubular neighbourhood of $\mathcal{M}^0$.

The proof of Proposition 8.2 is based on Proposition 7.1. The latter can be restated as follows. Let $\mathcal{H}^\varepsilon \subset T\mathcal{B}$ denote the horizontal subbundle with fibres

$$\mathcal{H}^\varepsilon_{(u,A)} := \ker d^{*_\varepsilon}_{(u,A)} \subset T_{(u,A)}\mathcal{B}.$$

Given a pair $(u,A) \in \mathcal{B}$ and constants p, c_0, δ, ε denote by $\mathcal{U}^0 = \mathcal{U}^0(\delta,\varepsilon) \subset \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0)$ the open set

$$\mathcal{U}^0 := \left\{(u_0,A_0) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0) \middle| (u,A) = (\exp_{u_0}(\xi), A_0 + \alpha), \|\zeta\|_{1,p,\varepsilon} \le \delta\varepsilon^{2/p}\right\}.$$

If δ and ε are sufficiently small then Proposition 7.1 defines two maps

$$\mathcal{S}^\varepsilon : \mathcal{U}^0 \to \mathcal{H}^\varepsilon, \qquad \mathcal{N}^\varepsilon : \mathcal{U}^0 \to \Omega^0(\Sigma, \mathfrak{g}_P)$$

such that $\mathcal{S}^\varepsilon$ is a section of $\mathcal{H}^\varepsilon$ over $\mathcal{U}^0$ and, for every $(u_0,A_0) \in \mathcal{U}^0$, the pair $(\xi_0,\alpha_0) := \mathcal{S}^\varepsilon(u_0,A_0)$ and the gauge transformation $g := e^\eta$, where $\eta := \mathcal{N}^\varepsilon(u_0,A_0)$, satisfy $g^*(u,A) = (\exp_{u_0}(\xi_0), A_0 + \alpha_0)$ and (45). In particular,

$$\|\mathcal{S}^\varepsilon(u_0,A_0)\|_{1,p,\varepsilon} \le c\|\zeta\|_{1,p,\varepsilon},$$

where $(u,A) = (\exp_{u_0}(\xi), A_0 + \alpha)$ and $\|\zeta\|_{1,p,\varepsilon} \le \delta\varepsilon^{2/p}$. In this notation Proposition 8.2 asserts that for every $(u,A) \in \mathcal{B}$, whose distance to $\widetilde{\mathcal{M}}^0_{B,\Sigma}$ in the $(1,p,\varepsilon)$-norm is less than $\delta\varepsilon^{2/p}$ for a sufficiently small constant δ, there exists a pair $(u_0,A_0) \in \mathcal{U}^0$ such that $\mathcal{S}^\varepsilon(u_0,A_0)$ lies in the image of $(\mathcal{D}^\varepsilon_{(u_0,A_0)})^*$.

Lemma 8.3. *For every $p > 2$ and every $c_0 > 0$ there exist positive constants δ, ε_0, and c such that the following holds for every $\varepsilon \in (0, \varepsilon_0]$. Let $(u, A) \in \mathcal{B}$ and $I \subset \mathbb{R}$ be an interval. Suppose that $I \to \mathcal{U}^0(\delta, \varepsilon) : r \mapsto (u_0(r), A_0(r))$ is a smooth path, and let $\zeta(r) = (\xi(r), \alpha(r)) \in T_{(u_0(r), A_0(r))}\mathcal{B}$ be the corresponding vector field along this path that satisfies*

$$(u, A) = (\exp_{u_0}(\xi), A_0 + \alpha), \qquad \|\zeta\|_{1,p,\varepsilon} \leq \delta\varepsilon^{2/p}.$$

Then the function $r \mapsto \zeta_\varepsilon(r) := \mathcal{S}^\varepsilon(u_0(r), A_0(r))$ satisfies the estimate

$$\left\|(\partial_r u_0, \partial_r A_0) + \tilde{\nabla}_r \zeta_\varepsilon\right\|_{1,p,\varepsilon} \leq c\left(\|\zeta\|_{1,p,\varepsilon} + \left\|\varepsilon^2 d_{A_0}^* \partial_r A_0 - L_{u_0}^* \partial_r u_0\right\|_{L^p}\right), \quad (74)$$

where $\tilde{\nabla}_r \zeta_\varepsilon := (\tilde{\nabla}_r \xi_\varepsilon, \partial_r \alpha_\varepsilon)$ and $\tilde{\nabla}_r \xi_\varepsilon := \nabla_r \xi_\varepsilon - \frac{1}{2} J(\nabla_{\partial_r u_0} J)\xi_\varepsilon$.

Proof. Let $g(r) := e^{\eta_\varepsilon(r)}$, where $\eta_\varepsilon(r) := \mathcal{N}^\varepsilon(u_0(r), A_0(r)) \in \Omega^0(\Sigma, \mathfrak{g}_P)$, and denote $\dot{u}_0 := \partial_r u_0$, $\dot{A}_0 := \partial_r A_0$, $\dot{g} := \partial_r g$. Let δ_1 and c_1 be the constants of Proposition 7.1. Then

$$\|\eta_\varepsilon(r)\|_{2,p,\varepsilon} + \|\zeta_\varepsilon(r)\|_{1,p,\varepsilon} \leq c_1 \|\zeta(r)\|_{1,p,\varepsilon} \leq c_1\delta_1\varepsilon^{2/p}, \quad (75)$$

$$g^{-1}u = \exp_{u_0}(\xi_\varepsilon), \qquad g^*A = A_0 + \alpha_\varepsilon, \qquad \varepsilon^2 d_{A_0}^* \alpha_\varepsilon - L_{u_0}^* \xi_\varepsilon = 0.$$

Differentiating these identities we obtain

$$-L_{g^{-1}u}(g^{-1}\dot{g}) = E_1\dot{u}_0 + E_2\nabla_r\xi_\varepsilon, \qquad d_{g^*A}(g^{-1}\dot{g}) = \dot{A}_0 + \dot{\alpha}_\varepsilon, \quad (76)$$

where $E_1 := E_1(u_0, \xi_\varepsilon)$ and $E_2 := E_2(u_0, \xi_\varepsilon)$ (see Appendix C), and

$$\varepsilon^2 d_{A_0}^* \dot{\alpha}_\varepsilon - \varepsilon^2 * [\dot{A}_0 \wedge *\alpha_\varepsilon] - L_{u_0}^* \nabla_r \xi_\varepsilon - \rho(\dot{u}_0, \xi_\varepsilon) = 0, \quad (77)$$

where $\rho \in \Omega^2(M, \mathfrak{g})$ is given by $\langle \eta, \rho(\xi_1, \xi_2)\rangle := \langle \nabla_{\xi_1} X_\eta, \xi_2\rangle$ (see Lemma C.2). Inserting the expressions for $\nabla_r \xi_\varepsilon$ and $\dot{\alpha}_\varepsilon$ in (76) into (77) gives

$$\begin{aligned}
&\varepsilon^2 d_{A_0}^* d_{g^*A}(g^{-1}\dot{g}) + L_{u_0}^* E_2^{-1} L_{g^{-1}u}(g^{-1}\dot{g}) \\
&= \varepsilon^2 d_{A_0}^* \dot{A}_0 + \varepsilon^2 * [\dot{A}_0 \wedge *\alpha_\varepsilon] - L_{u_0}^* E_2^{-1} E_1 \dot{u}_0 + \rho(\dot{u}_0, \xi_\varepsilon).
\end{aligned}$$

Since $g^{-1}u = \exp_{u_0}(\xi_\varepsilon)$ and $g^*A = A_0 + \alpha_\varepsilon$ we have, by Lemma C.1,

$$\begin{aligned}
E_2^{-1} L_{g^{-1}u}(g^{-1}\dot{g}) &= L_{u_0}(g^{-1}\dot{g}) + \left(E_2^{-1}E_1 - \mathbb{1}\right) L_{u_0}(g^{-1}\dot{g}) + \nabla_{\xi_\varepsilon} X_{g^{-1}\dot{g}}(u_0), \\
d_{A_0}^* d_{g^*A}(g^{-1}\dot{g}) &= d_{A_0}^* d_{A_0}(g^{-1}\dot{g}) + [d_{A_0}^* \alpha_\varepsilon, g^{-1}\dot{g}] + *[*\alpha_\varepsilon \wedge d_{A_0}(g^{-1}\dot{g})].
\end{aligned}$$

Hence

$$\begin{aligned}&\varepsilon^2 d_{A_0}^* d_{A_0}(g^{-1}\dot{g}) + L_{u_0}^* L_{u_0}(g^{-1}\dot{g}) \\ &= -\,\varepsilon^2[d_{A_0}^*\alpha_\varepsilon, g^{-1}\dot{g}] - \varepsilon^2 * [*\alpha_\varepsilon \wedge d_{A_0}(g^{-1}\dot{g})] \\ &\quad - L_{u_0}^*\left(E_2^{-1}E_1 - \mathbb{1}\right) L_{u_0}(g^{-1}\dot{g}) - L_{u_0}^* \nabla_{\xi_\varepsilon} X_{g^{-1}\dot{g}}(u_0) \\ &\quad + \varepsilon^2 * [\dot{A}_0 \wedge *\alpha_\varepsilon] + L_{u_0}^*(\mathbb{1} - E_2^{-1}E_1)\dot{u}_0 + \rho(\dot{u}_0, \xi_\varepsilon) \\ &\quad + \varepsilon^2 d_{A_0}^* \dot{A}_0 - L_{u_0}^* \dot{u}_0.\end{aligned}$$

By Lemma 7.2 and Lemma 4.4, there exists a constant $c_2 > 0$ such that

$$\begin{aligned}\left\|g^{-1}\dot{g}\right\|_{2,p,\varepsilon} &\le c_2\varepsilon^{-2/p}\left\|\zeta_\varepsilon\right\|_{1,p,\varepsilon}\left\|g^{-1}\dot{g}\right\|_{2,p,\varepsilon} \\ &\quad + c_2\left(\left\|\zeta_\varepsilon\right\|_{0,p,\varepsilon} + \left\|\varepsilon^2 d_{A_0}^*\dot{A}_0 - L_{u_0}^*\dot{u}_0\right\|_{L^p}\right) \\ &\le c_1c_2\delta_1\left\|g^{-1}\dot{g}\right\|_{2,p,\varepsilon} \\ &\quad + c_2\left(\left\|\zeta_\varepsilon\right\|_{0,p,\varepsilon} + \left\|\varepsilon^2 d_{A_0}^*\dot{A}_0 - L_{u_0}^*\dot{u}_0\right\|_{L^p}\right).\end{aligned}$$

The last inequality follows from (75). With $c_1c_2\delta_1 \le 1/2$ it follows that

$$\left\|g^{-1}\dot{g}\right\|_{2,p,\varepsilon} \le 2c_2\left(\left\|\zeta_\varepsilon\right\|_{0,p,\varepsilon} + \left\|\varepsilon^2 d_{A_0}^*\dot{A}_0 - L_{u_0}^*\dot{u}_0\right\|_{L^p}\right).$$

Hence (74) follows from (76) and (45). □

Consider the vector bundle

$$\mathcal{V}^0 \longrightarrow \widetilde{\mathcal{M}}_{B,\Sigma}^0$$

whose fibre over $(u_0, A_0) \in \widetilde{\mathcal{M}}_{B,\Sigma}^0$ is the finite dimensional vector space $\mathcal{V}_{(u_0,A_0)}^0$ of all pairs

$$(\xi_0, \alpha_0) \in \Omega^0(\Sigma, H_{u_0}) \times \Omega^1(\Sigma, \mathfrak{g}_P)$$

that satisfy the equation

$$D\bar{\partial}_{J,A_0}(u_0)\xi_0 + X_{\alpha_0}(u_0)^{0,1} = 0. \tag{78}$$

This space can be identified with the kernel of the operator $\mathcal{D}_{(u_0,A_0)}^0$. Namely, the kernel of $\mathcal{D}_{(u_0,A_0)}^0$ consists of all sections $\xi_0 \in \Omega^0(\Sigma, H_{u_0})$ that satisfy (78) for some 1-form $\alpha_0 \in \Omega^1(\Sigma, \mathfrak{g}_P)$, and the 1-form α_0 is uniquely determined by ξ_0. Thus $\mathcal{V}^0 \longrightarrow \widetilde{\mathcal{M}}_{B,\Sigma}^0$ is a vector bundle of rank $m := \dim \mathcal{M}_{B,\Sigma}^0$.

Lemma 8.4. *For every $p > 2$ and every $c_0 > 0$ there exist positive constants ε_0 and c such that the following holds for every $\varepsilon \in (0, \varepsilon_0]$. Let $I \subset \mathbb{R}$ be an interval and*

$$I \to \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0) : r \mapsto (u_0(r), A_0(r))$$

be a smooth path such that, for every $r \in I$,

$$\|\partial_r u_0(r)\|_{C^2(\Sigma)} + \|\partial_r A_0(r)\|_{C^1(\Sigma)} \le c_0. \tag{79}$$

Then every smooth vector field $r \mapsto \zeta(r) = (\xi(r), \alpha(r)) \in T_{(u_0(r),A_0(r))}\mathcal{B}$ satisfies the inequality

$$\left\| \tilde{\nabla}_r \mathcal{D}^\varepsilon \zeta - \mathcal{D}^\varepsilon \tilde{\nabla}_r \zeta \right\|_{k,p,\varepsilon} \le c\varepsilon^{-1} \|\zeta\|_{k+1,p,\varepsilon}$$

for $k = 0, 1$, where $\mathcal{D}^\varepsilon := \mathcal{D}^\varepsilon_{(u_0(r),A_0(r))}$ and $\tilde{\nabla}_r \zeta := (\nabla_r \xi - \frac{1}{2} J(\nabla_{\partial_r u_0} J)\xi, \partial_r \alpha)$.

Proof. We denote

$$(\xi, \alpha) := \zeta, \qquad \zeta' := (\xi', \varphi', \psi') := \mathcal{D}^\varepsilon \zeta, \qquad \zeta'_r := (\xi'_r, \varphi'_r, \psi'_r) := \mathcal{D}^\varepsilon \tilde{\nabla}_r \zeta.$$

Moreover, we drop the subscript 0 and write $(u, A) := (u_0, A_0)$. Then, in local holomorphic coordinates on Σ and a local frame of P, ζ' is given by

$$\begin{aligned}
\xi' &= \tilde{\nabla}_{A,s}\xi + J\tilde{\nabla}_{A,t}\xi + \frac{1}{4}N(\xi, v_s - Jv_t) + \frac{1}{2}(J\partial_s J - \partial_t J)\xi + L_u\varphi + JL_u\psi, \\
\varphi' &= \lambda^{-2}\left(\nabla_{A,s}\varphi + \nabla_{A,t}\psi\right) + \varepsilon^{-2}L_u^*\xi, \\
\psi' &= \lambda^{-2}\left(\nabla_{A,s}\psi - \nabla_{A,t}\varphi\right) + \varepsilon^{-2}d\mu(u)\xi.
\end{aligned}$$

Here we use the notation of Remark 4.1. Differentiating these formulae with respect to r we obtain

$$\begin{aligned}
\tilde{\nabla}_r\xi' - \xi'_r &= \tilde{\nabla}_r\tilde{\nabla}_{A,s}\xi - \tilde{\nabla}_{A,s}\tilde{\nabla}_r\xi + J(\tilde{\nabla}_r\tilde{\nabla}_{A,t}\xi - \tilde{\nabla}_{A,t}\tilde{\nabla}_r\xi) \\
&\quad + \frac{1}{4}\tilde{\nabla}_r N(\xi, v_s - Jv_t) - \frac{1}{4}N(\tilde{\nabla}_r\xi, v_s - Jv_t) \\
&\quad + \frac{1}{2}\tilde{\nabla}_r((J\partial_s J - \partial_t J)\xi) - \frac{1}{2}(J\partial_s J - \partial_t J)\tilde{\nabla}_r\xi \\
&\quad + \nabla_{\partial_r u}X_\varphi(u) + J\nabla_{\partial_r u}X_\psi(u) - \frac{1}{2}J(\nabla_{\partial_r u}J)(L_u\varphi - JL_u\psi), \\
\partial_r\varphi' - \varphi'_r &= \lambda^{-2}\left([\partial_r\Phi, \varphi] + [\partial_r\Psi, \psi]\right) \\
&\quad + \varepsilon^{-2}\left(\rho(\partial_r u, \xi) - \frac{1}{2}d\mu(u)(\nabla_{\partial_r u}J)\xi\right), \\
\partial_r\psi' - \psi'_r &= \lambda^{-2}\left([\partial_r\Phi, \psi] - [\partial_r\Psi, \varphi]\right) \\
&\quad - \varepsilon^{-2}\left(\rho(\partial_r u, J\xi) + \frac{1}{2}L_u^*(\nabla_{\partial_r u}J)\xi\right).
\end{aligned}$$

Here $\rho = \rho_{s,t} \in \Omega^2(M, \mathfrak{g})$ is defined by Lemma C.2. The required estimates follow from these three identities via a term by term inspection. $\square$

Lemma 8.5. *For every* $p > 2$ *and every* $c_0 > 0$ *there exist positive constants* ε_0 *and* c *such that the following holds for every* $\varepsilon \in (0, \varepsilon_0]$. *Let* $r \mapsto (u_0(r), A_0(r))$ *be as in Lemma 8.4 and suppose that* $r \mapsto \zeta_0(r)$ *is a smooth section of* $\mathcal{V}^0$ *along this path. Abbreviate* $\mathcal{D}^\varepsilon := \mathcal{D}^\varepsilon_{(u_0(r),A_0(r))}$ *and let* $\zeta_\varepsilon(r) \in \ker \mathcal{D}^\varepsilon$ *be given by*

$$\zeta_\varepsilon(r) := \zeta_0(r) - \mathcal{D}^{\varepsilon *} \left(\mathcal{D}^\varepsilon \mathcal{D}^{\varepsilon *}\right)^{-1} \mathcal{D}^\varepsilon \zeta_0(r).$$

Then

$$\begin{aligned} \|\zeta_\varepsilon - \zeta_0\|_{0,2,\varepsilon} &\leq c\varepsilon^2 \|\xi_0\|_{L^2}, &(80)\\ \|\tilde{\nabla}_r \zeta_\varepsilon - \tilde{\nabla}_r \zeta_0\|_{0,2,\varepsilon} &\leq c\left(\|\xi_0\|_{L^2} + \|\tilde{\nabla}_r \xi_0\|_{L^2}\right). &(81)\end{aligned}$$

Proof. Let $(\xi_\varepsilon, \alpha_\varepsilon) := \zeta_\varepsilon$ for $\varepsilon \geq 0$ and $\zeta := (\xi, \alpha) := \zeta_\varepsilon - \zeta_0$. Then

$$\zeta \in \operatorname{im} \mathcal{D}^{\varepsilon *}, \qquad \mathcal{D}^\varepsilon \zeta = \begin{pmatrix} 0 \\ d_{A_0}^* \alpha_0 \\ - * d_{A_0} \alpha_0 \end{pmatrix}.$$

Hence, by Lemma 4.8, there exist constants $c_1, c_2 > 0$ (depending only on c_0) such that

$$\|\zeta\|_{1,2,\varepsilon} \leq c_1 \varepsilon \left\|\mathcal{D}^\varepsilon \zeta\right\|_{0,2,\varepsilon} = c_1 \varepsilon^2 \left(\|d_{A_0}\alpha_0\|_{L^2} + \left\|d_{A_0}^*\alpha_0\right\|_{L^2}\right) \leq c_2 \varepsilon^2 \left\|\xi_0\right\|_{L^2}.$$

The last inequality follows from (78) and the basic elliptic estimates for the operator $\mathcal{D}^0$. Thus we have proved (80). To prove (81) let

$$\zeta' := \zeta'(r) := -(\mathcal{D}^\varepsilon \mathcal{D}^{\varepsilon *})^{-1} \mathcal{D}^\varepsilon \zeta_0(r) \in \mathcal{X}'_{u_0(r)}$$

so that $\zeta = \mathcal{D}^{\varepsilon *}\zeta'$. Then, by Lemma 4.7 with $p = 2$,

$$\|\zeta'\|_{2,2,\varepsilon} \leq c_3 \left\|\mathcal{D}^{\varepsilon *}\zeta'\right\|_{1,2,\varepsilon} = c_3 \left\|\zeta\right\|_{1,2,\varepsilon} \leq c_2 c_3 \varepsilon^2 \left\|\xi_0\right\|_{L^2},$$

and hence, by Lemma 8.4 (with $c = c_4$ and $\mathcal{D}^\varepsilon$ replaced by $\mathcal{D}^{\varepsilon *}$),

$$\|\mathcal{D}^{\varepsilon *}\tilde{\nabla}_r \zeta' - \tilde{\nabla}_r \mathcal{D}^{\varepsilon *}\zeta'\|_{1,2,\varepsilon} \leq c_4 \varepsilon^{-1} \left\|\zeta'\right\|_{2,2,\varepsilon} \leq c_2 c_3 c_4 \varepsilon \left\|\xi_0\right\|_{L^2}.$$

Now it follows from Lemmata 4.8 (with $c = c_5$) and 8.4 (with $c = c_4$) that

$$\begin{aligned}
\|\mathcal{D}^{\varepsilon *}\tilde{\nabla}_r\zeta'\|_{1,2,\varepsilon} &\leq c_5\|\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon *}\tilde{\nabla}_r\zeta'\|_{0,2,\varepsilon} \\
&\leq c_5\|\mathcal{D}^{\varepsilon}(\mathcal{D}^{\varepsilon *}\tilde{\nabla}_r\zeta' - \tilde{\nabla}_r\mathcal{D}^{\varepsilon *}\zeta')\|_{0,2,\varepsilon} \\
&\quad + c_5\|\mathcal{D}^{\varepsilon}\tilde{\nabla}_r\zeta - \tilde{\nabla}_r\mathcal{D}^{\varepsilon}\zeta\|_{0,2,\varepsilon} + c_5\|\tilde{\nabla}_r\mathcal{D}^{\varepsilon}\zeta\|_{0,2,\varepsilon} \\
&\leq c_6\varepsilon^{-1}\|\mathcal{D}^{\varepsilon *}\tilde{\nabla}_r\zeta' - \tilde{\nabla}_r\mathcal{D}^{\varepsilon *}\zeta'\|_{1,2,\varepsilon} \\
&\quad + c_5\left(c_4\varepsilon^{-1}\|\zeta\|_{1,2,\varepsilon} + \|\tilde{\nabla}_r\mathcal{D}^{\varepsilon}\zeta\|_{0,2,\varepsilon}\right) \\
&\leq c_7\left(\|\xi_0\|_{L^2} + \|\tilde{\nabla}_r\xi_0\|_{L^2}\right).
\end{aligned}$$

Hence

$$\begin{aligned}
\|\tilde{\nabla}_r\zeta\|_{0,2,\varepsilon} &\leq \|\tilde{\nabla}_r\mathcal{D}^{\varepsilon *}\zeta' - \mathcal{D}^{\varepsilon *}\tilde{\nabla}_r\zeta'\|_{0,2,\varepsilon} + \|\mathcal{D}^{\varepsilon *}\tilde{\nabla}_r\zeta'\|_{0,2,\varepsilon} \\
&\leq (c_7 + c_2c_3c_4\varepsilon)\left(\|\xi_0\|_{L^2} + \|\tilde{\nabla}_r\xi_0\|_{L^2}\right).
\end{aligned}$$

This proves (81). □

The estimate (81) is fairly crude. More careful considerations give an additional factor ε. However, we shall not use this fact.

Lemma 8.6. *For every $p > 2$ and every $c_0 > 0$ there exist positive constants δ, ε_0, and c such that the following holds for every $\varepsilon \in (0, \varepsilon_0]$. Let $(u, A) \in \mathcal{B}$ and $I \subset \mathbb{R}$ be an interval. Suppose that the path $I \to \mathcal{U}^0(\delta, \varepsilon) : r \mapsto (u_0(r), A_0(r))$ and the vector field $r \mapsto \zeta(r) \in T_{(u_0(r),A_0(r))}\mathcal{B}$ are as in Lemma 8.3. Moreover, assume (79) and let $r \mapsto \zeta_0(r)$ and $r \mapsto \zeta_\varepsilon(r)$ be as in Lemma 8.5. Then*

$$\begin{aligned}
&\left|\frac{d}{dr}\left\langle\zeta_\varepsilon, \mathcal{S}^{\varepsilon}(u_0, A_0)\right\rangle_\varepsilon + \langle\xi_0, \dot{u}_0\rangle\right| \\
&\qquad \leq c\left(\|\xi_0\|_{L^2} + \|\tilde{\nabla}_r\xi_0\|_{L^2}\right)\left(\varepsilon^2 + \|\zeta\|_{1,p,\varepsilon} + \|L_{u_0}^*\partial_r u_0\|_{L^p}\right).
\end{aligned}$$

Proof. Abbreviate $\mathcal{S}^\varepsilon := \mathcal{S}^\varepsilon(u_0, A_0)$. Consider the identity

$$\begin{aligned}
\frac{d}{dr}\left\langle\zeta_\varepsilon, \mathcal{S}^\varepsilon\right\rangle_\varepsilon + \left\langle\xi_0, \partial_r u_0\right\rangle_\varepsilon &= \left\langle\tilde{\nabla}_r\zeta_\varepsilon, \mathcal{S}^\varepsilon\right\rangle_\varepsilon + \left\langle\zeta_\varepsilon, (\partial_r u_0, \partial_r A_0) + \tilde{\nabla}_r\mathcal{S}^\varepsilon\right\rangle_\varepsilon \\
&\quad + \left\langle\zeta_0 - \zeta_\varepsilon, (\partial_r u_0, \partial_r A_0)\right\rangle_\varepsilon - \varepsilon^2\langle\alpha_0, \partial_r A_0\rangle.
\end{aligned}$$

By Lemma 8.5, the $(0, 2, \varepsilon)$-norm of $\tilde{\nabla}_r\zeta_\varepsilon$ is bounded above by a constant times $\|\xi_0\|_{L^2} + \|\tilde{\nabla}_r\xi_0\|_{L^2}$. By Proposition 7.1, the $(0, 2, \varepsilon)$-norm of $\mathcal{S}^\varepsilon$ is bounded above by a constant times $\|\zeta\|_{1,p,\varepsilon}$. Hence the first term satisfies the required bound. For the second term the estimate follows from Lemma 8.3 and the fact that the $(0,2,\varepsilon)$-norm of ζ_ε is bounded above by a constant times $\|\xi_0\|_{L^2}$. For the third term we use (79) and (80) and for the last the estimate follows from (79). $\square$

Proof of Proposition 8.2. Let $U \subset \mathbb{R}^m$ be an open set containing zero and

$$U \longrightarrow \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0) : x \mapsto (u_0(x), A_0(x))$$

be the composition of the map $\mathcal{F}^0_{(\bar{u}_0,\bar{A}_0)}$ defined in Theorem 2.1 with a Hilbert space isomorphism $\mathbb{R}^m \to \ker \mathcal{D}^0_{(\bar{u}_0,\bar{A}_0)}$. Then

$$(u_0(0), A_0(0)) = (\bar{u}_0, \bar{A}_0), \qquad (\partial_i u_0(0), \partial_i A_0(0)) \in \mathcal{V}^0_{(\bar{u}_0,\bar{A}_0)}$$

for $i = 1, \ldots, m$; in particular,

$$L^*_{\bar{u}_0} \partial_i u_0(0) = 0, \qquad \langle \partial_i u_0(0), \partial_j u_0(0) \rangle_{L^2(\Sigma)} = \delta_{ij}.$$

Now choose m smooth sections $\zeta_{10}, \ldots, \zeta_{m0} : U \to \mathcal{V}^0$ so that

$$\zeta_{j0}(x) = (\xi_{j0}(x), \alpha_{j0}(x)) \in \mathcal{V}^0_{(u_0(x),A_0(x))}$$

and

$$\zeta_{j0}(0) = -(\partial_j u_0(0), \partial_j A_0(0)), \qquad \langle \xi_{i0}(x), \xi_{j0}(x) \rangle_{L^2(\Sigma)} = \delta_{ij}$$

for $x \in U$ and $j = 1, \ldots, m$. Given $x \in U$ we abbreviate $\mathcal{D}^\varepsilon := \mathcal{D}^\varepsilon_{(u_0(x),A_0(x))}$. If ε is sufficiently small then, by Lemma 4.8, this operator is surjective for every $x \in U$. In this case we define $\zeta_{j\varepsilon}(x) \in \ker \mathcal{D}^\varepsilon$ by

$$\zeta_{j\varepsilon}(x) := \zeta_{j0}(x) - \mathcal{D}^{\varepsilon *} \left(\mathcal{D}^\varepsilon \mathcal{D}^{\varepsilon *} \right)^{-1} \mathcal{D}^\varepsilon \zeta_{j0}(x)$$

for $j = 1, \ldots, m$. By Lemma 8.5, these vectors form a basis of ker $\mathcal{D}^\varepsilon$ for ε sufficiently small. Now let δ_1 and c_1 be the constants of Proposition 7.1. Choose $\delta_0 > 0$ so small that

$$|x| < \delta_0 \varepsilon^{2/p} \qquad \Longrightarrow \qquad (u_0(x), A_0(x)) \in \mathcal{U}^0(\delta_1, \varepsilon)$$

for $x \in U$ and $0 < \varepsilon \le 1$. Let $\mathcal{S}^\varepsilon : \mathcal{U}^0 \to \mathcal{H}^\varepsilon$ be the map of Proposition 7.1 as introduced above. Define $\theta = (\theta_1, \ldots, \theta_m) : U \to \mathbb{R}^m$ by

$$\theta_j(x) := \langle \zeta_{j\varepsilon}(x), \mathcal{S}^\varepsilon(u_0(x), A_0(x)) \rangle_\varepsilon ,$$

where $\langle\cdot,\cdot\rangle_\varepsilon$ denotes the $(0,2,\varepsilon)$-inner product on $T_{(u_0(x),A_0(x))}\mathcal{B}$. Then

$$\theta(x) = 0 \qquad \iff \qquad \mathcal{S}^\varepsilon(u_0(x), A_0(x)) \in \operatorname{im}\mathcal{D}^{\varepsilon *}.$$

We shall establish the existence of a zero of θ with the inverse function theorem. We must prove that $|d\theta(x) - \mathbb{1}| \le 1/2$ on a ball of radius r and that $\theta(0)$ is less than $r/2$.

To see this, we first observe that

$$|\theta_j(0)| \le \|\zeta_{j0}(0)\|_{0,2,\varepsilon} \left\|\mathcal{S}^\varepsilon(\bar{u}_0, \bar{A}_0)\right\|_{0,2,\varepsilon} \le c_2 \left\|\bar{\zeta}\right\|_{1,p,\varepsilon} \le c_2\delta\varepsilon^{2/p}.$$

Here we have used the fact that the $(0,2,\varepsilon)$-norm of $\zeta_{j\varepsilon}(0)$ is less than or equal to the $(0,2,\varepsilon)$-norm of $\zeta_{j0}(0) = -(\partial_j u_0(0), \partial_j A_0(0))$, that the L^2-norm of $\partial_j A_0(0)$ is controlled by $\|\partial_j u_0(0)\|_{L^2} = 1$, that the $(0,2,\varepsilon)$-norm of $\mathcal{S}^\varepsilon(\bar{u}_0, \bar{A}_0)$ is controlled by its $(1,p,\varepsilon)$-norm, and that, by Proposition 7.1 and (70), the latter is bounded above by $c_1\|\bar{\zeta}\|_{1,p,\varepsilon} \le c_1\delta\varepsilon^{2/p}$. Thus we have proved that

$$|\theta(0)| \le \sqrt{m}c_2 \left\|\bar{\zeta}\right\|_{1,p,\varepsilon} \le \sqrt{m}c_2\delta\varepsilon^{2/p}. \tag{82}$$

Now let $\zeta(x) = (\xi(x), \alpha(x)) \in T_{(u_0(x),A_0(x))}\mathcal{B}$ be the unique smooth section defined by

$$(u, A) = (\exp_{u_0(x)}(\xi(x)), A_0(x) + \alpha(x)), \qquad \zeta(0) = \bar{\zeta},$$

for x sufficiently small. Then there exists a constant $c_3 > 0$ such that

$$\|\zeta(x)\|_{1,p,\varepsilon} \le \left\|\bar{\zeta}\right\|_{1,p,\varepsilon} + c_3|x|$$

for x sufficiently small and $0 < \varepsilon \le 1$. Hence, by Proposition 7.1, we have that, for $|x| < \delta_0\varepsilon^{2/p}$ and $0 < \varepsilon \le 1$,

$$\begin{aligned} &\|\mathcal{N}^\varepsilon(u_0(x), A_0(x))\|_{2,p,\varepsilon} + \|\mathcal{S}^\varepsilon(u_0(x), A_0(x))\|_{1,p,\varepsilon} \\ &\le c_1 \|\zeta(x)\|_{1,p,\varepsilon} \le c_1 \left(\left\|\bar{\zeta}\right\|_{1,p,\varepsilon} + c_3|x|\right). \end{aligned} \tag{83}$$

Moreover, there exists a constant $c_4 > 0$ such that

$$|\delta_{ij} + \langle\xi_{j0}(x), \partial_i u_0(x)\rangle_{L^2}| + \|L^*_{u_0(x)}\partial_i u_0(x)\|_{L^p} \le c_4|x|,$$

$$\|\xi_{j0}(x)\|_{L^2} + \|\tilde{\nabla}_i\xi_{j0}(x)\|_{L^2} \le c_4$$

for x sufficiently small. Now suppose that δ_1 and ε_0 have been chosen so small that the assertion of Lemma 8.6 holds, with c replaced by c_5, for the paths $x_i \mapsto (u_0(x), A_0(x))$, $x_i \mapsto \zeta(x)$, $x_i \mapsto \zeta_{j0}(x)$, and $x_i \mapsto \zeta_{j\varepsilon}(x)$. Then

$$\begin{aligned}
|\partial_i\theta_j(x) - \delta_{ij}| &\le |\delta_{ij} + \langle \xi_{j0}(x), \partial_i u_0(x)\rangle| + |\partial_i\theta_j(x) + \langle \xi_{j0}(x), \partial_i u_0(x)\rangle| \\
&\le c_4|x| + c_5\big(\|\xi_{j0}(x)\|_{L^2} + \|\tilde\nabla_i \xi_{j0}(x)\|_{L^2}\big)\cdot \\
&\qquad \cdot\big(\varepsilon^2 + \|\zeta(x)\|_{1,p,\varepsilon} + \|L^*_{u_0(x)}\partial_i u_0(x)\|_{L^p}\big) \\
&\le c_4|x| + c_4c_5\big(\varepsilon^2 + \|\bar\zeta\|_{1,p,\varepsilon} + (c_3+c_4)|x|\big) \\
&\le c_6\big(\varepsilon^2 + \|\bar\zeta\|_{1,p,\varepsilon} + |x|\big)
\end{aligned}$$

for $|x| < \delta_0\varepsilon^{2/p}$ and $0 < \varepsilon \le \varepsilon_0$. Thus the Jacobian $d\theta(x) \in \mathbb{R}^{m\times m}$ satisfies

$$|d\theta(x) - \mathbb{1}| \le c_7\left(\varepsilon^2 + \delta\varepsilon^{2/p} + |x|\right).$$

Choose δ_0 and ε_0 so small that $c_7(\varepsilon_0^2 + 2\delta_0\varepsilon_0^{2/p}) \le 1/2$. Then

$$|x| \le \delta_0\varepsilon^{2/p}, \quad 0 < \varepsilon \le \varepsilon_0, \quad 0 < \delta \le \delta_0 \qquad \Longrightarrow \qquad |d\theta(x) - \mathbb{1}| \le 1/2.$$

Hence the inverse function theorem asserts that $\theta(B_r(0)) \supset B_{r/2}(\theta(0))$ whenever $r < \delta_0\varepsilon^{2/p}$. Now suppose that $\sqrt{m}c_2\delta < \delta_0/2$. Then, by (82), we have $2\,|\theta(0)| < \delta_0\varepsilon^{2/p}$ and hence we can apply the inverse function theorem with $r = 2\,|\theta(0)|$. Then $B_{r/2}(\theta(0))$ contains zero and, by the inverse function theorem, there exists a point $x_0 \in \mathbb{R}^m$ such that

$$\theta(x_0) = 0, \qquad |x_0| \le 2\,|\theta(0)| \le 2\sqrt{m}c_2\left\|\bar\zeta\right\|_{1,p,\varepsilon}.$$

The last inequality follows from (82). Now define

$$(u_0, A_0) := (u_0(x_0), A_0(x_0)), \qquad \zeta_0 := \mathcal{S}^\varepsilon(u_0, A_0), \qquad \eta_0 := \mathcal{N}^\varepsilon(u_0, A_0).$$

Then (71) and (72) are satisfied by definition. The estimate (73) follows from Proposition 7.1:

$$\|\eta_0\|_{2,p,\varepsilon} + \|\zeta_0\|_{1,p,\varepsilon} \le c_1\,\|\zeta(x_0)\|_{1,p,\varepsilon} \le c_1\left(\left\|\bar\zeta\right\|_{1,p,\varepsilon} + c_3|x_0|\right) \le c_8\left\|\bar\zeta\right\|_{1,p,\varepsilon}.$$

Moreover, the vector $\bar\xi_0$ in the assertion of Proposition 8.2 is the image of x_0 under our Hilbert space isomorphism $\mathbb{R}^m \to \ker\mathcal{D}^0_{(\bar u_0,\bar A_0)}$. Hence, by elliptic regularity for the Cauchy–Riemann operator, its $W^{1,p}$-norm is bounded by $|x_0|$ and hence by the $(1,p,\varepsilon)$-norm of $\bar\zeta$. $\square$

Proof of Theorem 8.1. Let $p > 2$ and $c_0 > 0$ be given. Choose positive constants ε_0, δ_1, δ_2, and c_1 such that Proposition 8.2 holds with δ replaced by δ_1 and c replaced by c_1, Corollary 6.3 holds with δ replaced by δ_2, and both results hold for $0 < \varepsilon \le \varepsilon_0$. Now choose $\delta > 0$ so small that

$$\delta {\varepsilon_0}^{1/2} \le \delta_1, \qquad c_1\delta \le \delta_2.$$

Let $\varepsilon \in (0, \varepsilon_0]$ and suppose that (u, A) and $(\bar{u}_0, \bar{A}_0)$ satisfy the hypotheses of Theorem B, namely

$$(\bar{u}_0, \bar{A}_0) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0 - 1)$$

and

$$(u, A) = (\exp_{\bar{u}_0}(\bar{\xi}), \bar{A}_0 + \bar{\alpha}) \in \widetilde{\mathcal{M}}^\varepsilon_{B,\Sigma},$$

where $\bar{\zeta} := (\bar{\xi}, \bar{\alpha}) \in T_{(\bar{u}_0,\bar{A}_0)}\mathcal{B}$ satisfies

$$\|\bar{\zeta}\|_{1,p,\varepsilon;(\bar{u}_0,\bar{A}_0)} \le \delta\varepsilon^{2/p+1/2} \le \delta_1\varepsilon^{2/p}.$$

By Proposition 8.2, there exist

$$\bar{\xi}_0 \in \ker \mathcal{D}^0_{(\bar{u}_0,\bar{A}_0)}, \qquad \eta_0 \in \Omega^0(\Sigma, \mathfrak{g}_P),$$

and

$$(u_0, A_0) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0), \qquad \zeta_0 = (\xi_0, \alpha_0) \in T_{(u_0,A_0)}\mathcal{B},$$

satisfying (71), (72), and (73), with c replaced by c_1. Hence

$$\|\zeta_0\|_{1,p,\varepsilon;(u_0,A_0)} \le c_1 \|\bar{\zeta}\|_{1,p,\varepsilon;(\bar{u}_0,\bar{A}_0)} \le c_1\delta\varepsilon^{2/p+1/2} \le \delta_2\varepsilon^{2/p+1/2}.$$

This estimate together with (72) shows that (u_0, A_0) and ζ_0 satisfy the hypotheses of Corollary 6.3. Hence, by (71),

$$g^*(u, A) = (\exp_{u_0}(\xi_0), A_0 + \alpha_0) = \widetilde{\mathcal{T}}^\varepsilon(u_0, A_0), \qquad g := e^{\eta_0}.$$

Moreover, again by (71),

$$(u_0, A_0) = \mathcal{F}^0_{(\bar{u}_0,\bar{A}_0)}(\bar{\xi}_0)$$

and, by (73),

$$\|\bar{\zeta}_0\|_{W^{1,p}} + \|\eta_0\|_{2,p,\varepsilon;\bar{A}_0} \le c_1 \|\bar{\zeta}\|_{1,p,\varepsilon;(\bar{u}_0,\bar{A}_0)}.$$

This proves the theorem. □

9 A priori estimates

In this section we assume that $J \in \mathcal{J}_{\mathrm{G}}(M,\omega)$ is independent of $z \in \Sigma$ (or in local coordinates is independent of s and t). Let $\Omega \subset \mathbb{C}$ be a bounded open set, $K \subset \Omega$ be a compact subset, and $\lambda : \bar{\Omega} \to (0,\infty)$ be a smooth function. Given $u : \Omega \to M$ and $\Phi, \Psi : \Omega \to \mathfrak{g}$ we define $v_s, v_t : \Omega \to u^*TM$ and $\kappa : \Omega \to \mathfrak{g}$ by

$$v_s := \partial_s u + X_\Phi(u), \quad v_t := \partial_t u + X_\Psi(u), \quad \kappa := \partial_s \Psi - \partial_t \Phi + [\Phi, \Psi].$$

Moreover, as in Remark 4.1, we use the notation

$$\begin{array}{lll} \nabla_{A,s}\xi & := & \nabla_s \xi + \nabla_\xi X_\Phi(u), \\ \nabla_{A,s}\eta & := & \partial_s \eta + [\Phi, \eta], \end{array} \qquad \begin{array}{lll} \nabla_{A,t}\xi & := & \nabla_t \xi + \nabla_\xi X_\Psi(u), \\ \nabla_{A,t}\eta & := & \partial_t \eta + [\Psi, \eta], \end{array}$$

for $\xi : \Omega \to u^*TM$ and $\eta : \Omega \to \mathfrak{g}$. Then

$$\nabla_{A,s}\mu(u) = d\mu(u)v_s = -L_u^* J v_s, \qquad \nabla_{A,t}\mu(u) = d\mu(u)v_t = -L_u^* J v_t.$$

Moreover,

$$\nabla_{A,s} v_t - \nabla_{A,t} v_s = L_u \kappa,$$

by Lemma C.3,

$$\nabla_{A,s} L_u \eta - L_u \nabla_{A,s}\eta = \nabla_{v_s} X_\eta(u), \qquad \nabla_{A,t} L_u \eta - L_u \nabla_{A,t}\eta = \nabla_{v_t} X_\eta(u),$$

and, by Lemma B.4,

$$\nabla_{A,s}\nabla_{A,t}\xi - \nabla_{A,t}\nabla_{A,s}\xi = R(v_s, v_t)\xi + \nabla_\xi X_\kappa(u).$$

In local coordinates equations (7) have the form

$$v_s + J v_t = 0, \qquad \lambda^{-2}\kappa + \varepsilon^{-2}\mu(u) = 0. \tag{84}$$

If (84) holds then

$$(\nabla_{v_s} J) v_t = (\nabla_{v_t} J) v_s, \qquad \nabla_{A,s} v_s + \nabla_{A,t} v_t = -J L_u \kappa.$$

Given a constant $C > 0$ we denote by $M^C \subset M$ the compact subset of all $x \in M$ that satisfy

$$|\mu(x)| \le C, \qquad |\eta| \le C|L_x \eta|$$

for every $\eta \in \mathfrak{g}$.

Lemma 9.1. *For every $C > 0$ and every triple (Ω, K, λ) as above there exist constants $\varepsilon_0 = \varepsilon_0(C, \Omega, K, \lambda) > 0$ and $c = c(C, \Omega, K, \lambda) > 0$, such that the following holds for every $\varepsilon \in (0, \varepsilon_0]$. If u, Φ, and Ψ satisfy (84) and*

$$\|v_s\|_{L^\infty(\Omega)} \le C, \qquad u(\Omega) \subset M^C$$

then

$$\sup_K |\mu(u)|^2 \le c\varepsilon^3, \qquad \int_K |\mu(u)|^2 \le c\varepsilon^4.$$

Proof. Let $\Delta = \partial^2/\partial s^2 + \partial^2/\partial t^2$ denote the standard Laplacian. For $r > 0$ denote $B_r(z_0) := \{z \in \mathbb{C} \,|\, |z - z_0| < r\}$, $B_r := B_r(0)$, and

$$\Omega_r := \{z \in \mathbb{C} \,|\, B_r(z) \subset \Omega\} \subset \Omega.$$

By (84), we have

$$d\mu(u)v_s = -L_u^* J v_s = -L_u^* v_t, \qquad d\mu(u)v_t = -L_u^* J v_t = L_u^* v_s,$$

and hence

$$\begin{aligned}
\nabla_{A,s}\nabla_{A,s}\mu(u) + \nabla_{A,t}\nabla_{A,t}\mu(u) &= \nabla_{A,s} d\mu(u)v_s + \nabla_{A,t}\mathrm{d}\mu(u)v_t \\
&= \nabla_{A,t} L_u^* v_s - \nabla_{A,s} L_u^* v_t \\
&= L_u^*(\nabla_{A,t} v_s - \nabla_{A,s} v_t) - 2\rho(v_s, v_t) \\
&= -L_u^* L_u \kappa - 2\rho(v_s, v_t) \\
&= (\lambda/\varepsilon)^2 L_u^* L_u \mu(u) - 2\rho(v_s, v_t).
\end{aligned}$$

Here $\rho \in \Omega^2(M, \mathfrak{g})$ is as in Lemma C.2. Thus

$$\begin{aligned}
\Delta|\mu(u)|^2 &= 2|\nabla_{A,s}\mu(u)|^2 + 2|\nabla_{A,t}\mu(u)|^2 \\
&\quad + 2\langle \mu(u), \nabla_{A,s}\nabla_{A,s}\mu(u) + \nabla_{A,t}\nabla_{A,t}\mu(u)\rangle \\
&= 2|\nabla_{A,s}\mu(u)|^2 + 2|\nabla_{A,t}\mu(u)|^2 \\
&\quad + 2(\lambda/\varepsilon)^2 |L_u\mu(u)|^2 - 4\langle \mu(u), \rho(v_s, v_t)\rangle.
\end{aligned}$$

Now choose a constant $c > 0$ such that

$$\min_\Omega \lambda^2 \ge c^{-1}, \qquad 2|\rho(\xi_1, \xi_2)| \le c\,|\xi_1|\,|\xi_2|\,,$$

for all $x \in M^C$ and $\xi_1, \xi_2 \in T_x M$. Then

$$\Delta\,|\mu(u)|^2 \ge \frac{2}{cC^2\varepsilon^2}\,|\mu(u)|^2 - 2cC^2\,|\mu(u)| \ge \frac{1}{cC^2\varepsilon^2}\,|\mu(u)|^2 - c^3C^6\varepsilon^2$$

and hence

$$\delta\,|\mu(u)|^2 \le a\varepsilon^4 + \varepsilon^2\Delta\,|\mu(u)|^2\,, \tag{85}$$

where $\delta := 1/cC^2$ and $a := c^3C^6$. Fix a constant $r > 0$ such that

$$z \in K \qquad \Longrightarrow \qquad B_{3r}(z) \subset \Omega.$$

Then, by (85) and Lemma 9.2 below, we have

$$\delta\int_{B_{2r}(z)} |\mu(u)|^2 \le 9\pi r^2 a\varepsilon^4 + \frac{4\varepsilon^2}{r^2}\int_{B_{3r}(z)} |\mu(u)|^2 \le 9\pi r^2 a\varepsilon^4 + 36\pi C^2\varepsilon^2$$

for every $z \in \Omega_{3r}$. Applying Lemma 9.2 again we obtain

$$\delta\int_{B_r(z)} |\mu(u)|^2 \le 4\pi r^2 a\varepsilon^4 + \frac{4\varepsilon^2}{r^2}\int_{B_{2r}(z)} |\mu(u)|^2 \le c'\varepsilon^4$$

for every $z \in \Omega_{3r}$, where $c' := 4\pi(ar^2 + 9a\varepsilon^2/\delta + 36C^2/\delta r^2)$. The L^2-estimate now follows by taking the sum over finitely many balls of radius r that cover the compact set Ω_{3r}. Moreover, by (85), the function $z \mapsto |\mu(u(z))|^2 + 4^{-1}a\varepsilon^2|z - z_0|^2$ is subharmonic in Ω for every $z_0 \in \mathbb{C}$. Hence, by the mean value inequality,

$$|\mu(u(z))|^2 \le \frac{a\varepsilon^2}{4\pi s^2}\int_{B_s} |z|^2 + \frac{1}{\pi s^2}\int_{B_s(z)} |\mu(u)|^2 \le \frac{a\varepsilon^2 s^2}{8} + \frac{c'\varepsilon^4}{\delta\pi s^2}$$

for $z \in \Omega_{3r}$ and $0 \le s \le r$. Assume $\varepsilon \le r^2$. Then we can choose $s := \sqrt{\varepsilon}$ and this proves L^∞ estimate. □

Lemma 9.2. *Let $u : B_{R+r} \to \mathbb{R}$ be a C^2-function and $f, g : B_{R+r} \to \mathbb{R}$ be continuous such that*

$$f \le g + \Delta u, \qquad u \ge 0, \qquad f \ge 0, \qquad g \ge 0.$$

Then

$$\int_{B_R} f \le \int_{B_{R+r}} g + \frac{4}{r^2}\int_{B_{R+r}\setminus B_R} u.$$

Moreover, if $g = cu$ then

$$\frac{\pi}{2}\sup_{B_R} u \le \left(c + \frac{4}{r^2}\right)\int_{B_{R+r}} u.$$

Proof. For $0 \le s \le r$ we have

$$\int_{B_R} f - \int_{B_{R+r}} g \le \int_{B_{R+s}} (f-g) \le \int_{B_{R+s}} \Delta u = \int_{\partial B_{R+s}} \frac{\partial u}{\partial \nu}$$

and hence

$$\frac{d}{ds}\int_{\partial B_{R+s}} u = \int_{\partial B_{R+s}} \frac{\partial u}{\partial \nu} + \frac{1}{R+s}\int_{\partial B_{R+s}} u \ge \int_{B_R} f - \int_{B_{R+r}} g.$$

Integrate this inequality over the interval $0 \le s \le t$ to obtain

$$\int_{B_R} f - \int_{B_{R+r}} g \le \frac{1}{t}\int_{\partial B_{R+t}} u \le \frac{2}{r}\int_{\partial B_{R+t}} u$$

for $r/2 \le t \le r$. The first inequality follows by integrating this inequality over the interval $r/2 \le t \le r$. The second inequality was proved in [DS2, Lemma 7.3]. □

Lemma 9.3. *For every triple (Ω, K, λ) as above and every $C_0 > 0$ there exist constants $\varepsilon_0 = \varepsilon_0(\Omega, K, \lambda) > 0$ and $c = c(\Omega, K, \lambda, C_0) > 0$ such that the following holds for every $\varepsilon \in (0, \varepsilon_0]$. If u, Φ, and Ψ satisfy (84) and*

$$\|v_s\|_{L^\infty(\Omega)} + \varepsilon^{-1}\|\mu(u)\|_{L^\infty(\Omega)} \le C_0$$

then

$$\begin{aligned}
&\varepsilon^{-1}\|\mu(u)\|_{L^p(K)} + \|L_u^* v_s\|_{L^p(K)} + \|L_u^* J v_s\|_{L^p(K)} \\
&+ \varepsilon\|\nabla_{A,s} v_s\|_{L^p(K)} + \varepsilon\|\nabla_{A,t} v_s\|_{L^p(K)} \\
&\le c\varepsilon^{2/p}\left(\|v_s\|_{L^2(\Omega)} + \varepsilon^{-1}\|\mu(u)\|_{L^2(\Omega)}\right)
\end{aligned} \tag{86}$$

for $2 \le p \le \infty$.

Proof. Consider the functions $u_0, v_0 : \Omega \to \mathbb{R}$ given by

$$u_0 := \frac{1}{2}\left(|v_s|^2 + \frac{\lambda^2}{\varepsilon^2}|\mu(u)|^2\right),$$

$$v_0 := \frac{1}{2}\left(|\nabla_{A,s} v_s|^2 + |\nabla_{A,t} v_s|^2 + \frac{\lambda^4}{\varepsilon^4}|L_u\mu(u)|^2 + \frac{\lambda^2}{\varepsilon^2}|L_u^* v_s|^2 + \frac{\lambda^2}{\varepsilon^2}|L_u^* J v_s|^2\right).$$

We prove that there exists a constant $c_0 > 0$ such that

$$\Delta u_0 \geq v_0 - c_0 u_0. \tag{87}$$

To see this, recall from the proof of Lemma 9.1 that

$$\frac{1}{2}\Delta|\mu(u)|^2 = |L_u^* v_s|^2 + |L_u^* J v_s|^2 + \frac{\lambda^2}{\varepsilon^2}|L_u\mu(u)|^2 - 2\langle \mu(u), \rho(v_s, v_t)\rangle,$$

and hence

$$\begin{aligned}
&\frac{1}{2\varepsilon^2}\Delta(\lambda^2|\mu(u)|^2) \\
&= \frac{\Delta\lambda^2}{2\varepsilon^2}|\mu(u)|^2 + \frac{\lambda^2}{2\varepsilon^2}\Delta|\mu(u)|^2 + \frac{\partial_s\lambda^2}{\varepsilon^2}\partial_s|\mu(u)|^2 + \frac{\partial_t\lambda^2}{\varepsilon^2}\partial_t|\mu(u)|^2 \\
&= \frac{\Delta\lambda^2}{2\varepsilon^2}|\mu(u)|^2 + \frac{2\partial_t\lambda^2}{\varepsilon^2}\langle \mu(u), L_u^* v_s\rangle - \frac{2\partial_s\lambda^2}{\varepsilon^2}\langle \mu(u), L_u^* J v_s\rangle \\
&\quad + \frac{\lambda^2}{\varepsilon^2}|L_u^* v_s|^2 + \frac{\lambda^2}{\varepsilon^2}|L_u^* J v_s|^2 + \frac{\lambda^4}{\varepsilon^4}|L_u\mu(u)|^2 - \frac{2\lambda^2}{\varepsilon^2}\langle \mu(u), \rho(v_s, v_t)\rangle.
\end{aligned}$$

Moreover, by Lemma B.4 and Lemma C.3,

$$\begin{aligned}
&(\nabla_{A,s}\nabla_{A,s} + \nabla_{A,t}\nabla_{A,t})\, v_s \\
&= (\nabla_{A,t}\nabla_{A,s} - \nabla_{A,s}\nabla_{A,t})v_t + \nabla_{A,s}(\nabla_{A,s}v_s + \nabla_{A,t}v_t) - \nabla_{A,t}(\nabla_{A,s}v_t - \nabla_{A,t}v_s) \\
&= -R(v_s, v_t)v_t - \nabla_{v_t}X_\kappa(u) - \nabla_{A,s}(JL_u\kappa) - \nabla_{A,t}(L_u\kappa) \\
&= -R(v_s, v_t)v_t + \frac{\lambda^2}{\varepsilon^2}J\nabla_{v_s}X_{\mu(u)}(u) + \frac{\lambda^2}{\varepsilon^2}(\nabla_{v_s}J)L_u\mu(u) + \frac{2\lambda^2}{\varepsilon^2}\nabla_{v_t}X_{\mu(u)}(u) \\
&\quad + \frac{\lambda^2}{\varepsilon^2}L_uL_u^* v_s - \frac{\lambda^2}{\varepsilon^2}JL_uL_u^* J v_s + \frac{\partial_s\lambda^2}{\varepsilon^2}JL_u\mu(u) + \frac{\partial_t\lambda^2}{\varepsilon^2}L_u\mu(u).
\end{aligned} \tag{88}$$

Hence

$$\begin{aligned}
\frac{1}{2}\Delta|v_s|^2 &= |\nabla_{A,s}v_s|^2 + |\nabla_{A,t}v_s|^2 + \langle v_s, (\nabla_{A,s}\nabla_{A,s} + \nabla_{A,t}\nabla_{A,t})\, v_s\rangle \\
&= |\nabla_{A,s}v_s|^2 + |\nabla_{A,t}v_s|^2 + \frac{\lambda^2}{\varepsilon^2}|L_u^* v_s|^2 + \frac{\lambda^2}{\varepsilon^2}|L_u^* J v_s|^2 \\
&\quad - \frac{3\lambda^2}{\varepsilon^2}\langle \mu(u), \rho(v_s, v_t)\rangle + \frac{\lambda^2}{\varepsilon^2}\langle v_s, (\nabla_{v_s}J)L_u\mu(u)\rangle \\
&\quad - \langle v_s, R(v_s, v_t)v_t\rangle - \frac{\partial_s\lambda^2}{\varepsilon^2}\langle L_u^* J v_s, \mu(u)\rangle + \frac{\partial_t\lambda^2}{\varepsilon^2}\langle L_u^* v_s, \mu(u)\rangle.
\end{aligned}$$

Combining this with the formula for $\Delta(\lambda^2|\mu(u)|^2)/2\varepsilon^2$ we obtain

$$\begin{aligned}\Delta u_0 &= |\nabla_{A,s}v_s|^2 + |\nabla_{A,t}v_s|^2 + \frac{2\lambda^2}{\varepsilon^2}|L_u^*v_s|^2 + \frac{2\lambda^2}{\varepsilon^2}|L_u^*Jv_s|^2 + \frac{\lambda^4}{\varepsilon^4}|L_u\mu(u)|^2 \\ &\quad - \frac{5\lambda^2}{\varepsilon^2}\langle\mu(u),\rho(v_s,v_t)\rangle + \frac{3\partial_t\lambda^2}{\varepsilon^2}\langle\mu(u),L_u^*v_s\rangle - \frac{3\partial_s\lambda^2}{\varepsilon^2}\langle\mu(u),L_u^*Jv_s\rangle \\ &\quad + \frac{\Delta\lambda^2}{2\varepsilon^2}|\mu(u)|^2 + \frac{\lambda^2}{\varepsilon^2}\langle v_s,(\nabla_{v_s}J)L_u\mu(u)\rangle - \langle v_s,R(v_s,v_t)v_t\rangle. \qquad (89)\end{aligned}$$

The first row on the right is bounded below by $2v_0$. Moreover, by assumption, the image of u is contained in the compact set $\{|\mu(x)| \le \varepsilon C_0\}$. Hence the last six terms can be estimated from below by $v_0 - c_0u_0$ for some constant c_0 whenever ε is sufficiently small. Thus we have proved the inequality (87). Hence, by Lemma 9.2, there exist constants $\varepsilon_0 > 0$ and $c_0' > 0$ such that

$$\sup_K u_0 + \int_K v_0 \le c_0' \int_\Omega u_0$$

for every $\varepsilon \in (0,\varepsilon_0]$. Since $|\mu(u)| \le C_0\varepsilon_0$ and zero is a regular value of μ there is an inequality $|L_u\eta| \ge \delta|\eta|$ whenever ε_0 is sufficiently small. Thus we have proved (86) for $p = 2$ as well as

$$\|v_s\|_{L^\infty(K)} + \varepsilon^{-1}\|\mu(u)\|_{L^\infty(K)} \le c_0'\left(\|v_s\|_{L^2(\Omega)} + \varepsilon^{-1}\|\mu(u)\|_{L^2(\Omega)}\right). \qquad (90)$$

Now let us define $u_1 : \Omega \to \mathbb{R}$ by

$$u_1 := \frac{1}{2}|\nabla_{A,s}v_s|^2.$$

We shall prove that there exist positive constants δ_1, c_1, and ε_0 such that

$$\Delta(u_0 + \varepsilon^2u_1) \ge -c_1u_0 \qquad (91)$$

for $0 < \varepsilon \le \varepsilon_0$. We consider the equation

$$\Delta u_1 = |\nabla_{A,s}\nabla_{A,s}v_s|^2 + |\nabla_{A,t}\nabla_{A,s}v_s|^2 + \langle(\nabla_{A,s}\nabla_{A,s} + \nabla_{A,t}\nabla_{A,t})\nabla_{A,s}v_s, \nabla_{A,s}v_s\rangle$$

and use the formula

$$(\nabla_{A,s}\nabla_{A,s} + \nabla_{A,t}\nabla_{A,t})\nabla_{A,s}v_s = I + II + III,$$

where

$$\begin{aligned}
I &:= \left(\nabla_{A,t}\nabla_{A,s} - \nabla_{A,s}\nabla_{A,t}\right)\nabla_{A,t}v_s \\
&= -R(v_s, v_t)\nabla_{A,t}v_s + \frac{\lambda^2}{\varepsilon^2}\nabla_{\nabla_{A,t}v_s}X_\mu(u), \\
II &:= \nabla_{A,t}\left(\nabla_{A,t}\nabla_{A,s} - \nabla_{A,s}\nabla_{A,t}\right)v_s \\
&= -\nabla_{A,t}R(v_s, v_t)v_s + \frac{\lambda^2}{\varepsilon^2}\nabla_{A,t}(\nabla_{v_s}X_\mu(u)) + \frac{\partial_t\lambda^2}{\varepsilon^2}\nabla_{v_s}X_\mu(u), \\
III &:= \nabla_{A,s}\left(\nabla_{A,s}\nabla_{A,s} + \nabla_{A,t}\nabla_{A,t}\right)v_s \\
&= \nabla_{A,s}\Bigg(-R(v_s, v_t)v_t + \frac{\lambda^2}{\varepsilon^2}L_uL_u^*v_s - \frac{\lambda^2}{\varepsilon^2}JL_uL_u^*Jv_s + \frac{\lambda^2}{\varepsilon^2}(\nabla_{v_s}J)X_\mu(u) \\
&\quad + \frac{\lambda^2}{\varepsilon^2}J\nabla_{v_s}X_\mu(u) + \frac{2\lambda^2}{\varepsilon^2}\nabla_{v_t}X_\mu(u) + \frac{\partial_s\lambda^2}{\varepsilon^2}JX_\mu(u) + \frac{\partial_t\lambda^2}{\varepsilon^2}X_\mu(u)\Bigg).
\end{aligned}$$

Here we abbreviate $X_\mu(u) = X_{\mu(u)}(u) = L_u\mu(u)$. The last equality for III follows from (88). Now consider the tensors $\nabla^2 J$ and $\nabla^2 X_\eta$ defined by

$$\begin{aligned}
\nabla^2 J(X, Y, Z) &:= \nabla_X((\nabla_Y J)Z) - (\nabla_{\nabla_X Y}J)Z - (\nabla_Y J)\nabla_X Z, \\
\nabla^2 X_\eta(Y, Z) &:= \nabla_Y(\nabla_Z X_\eta) - \nabla_{\nabla_Y Z}X_\eta
\end{aligned}$$

for $\eta \in \mathfrak{g}$ and $X, Y, Z \in \mathrm{Vect}(M)$. Then

$$\begin{aligned}
\nabla_{A,s}((\nabla_{v_s}J)L_u\mu(u)) &= \nabla^2 J(v_s, v_s, L_u\mu(u)) + (\nabla_{\nabla_{A,s}v_s}J)L_u\mu(u) \\
&\quad + (\nabla_{v_s}J)\nabla_{v_s}X_{\mu(u)}(u) - (\nabla_{v_s}J)L_uL_u^*Jv_s, \\
\nabla_{A,t}(\nabla_{v_s}X_{\mu(u)}(u)) &= \nabla^2 X_{\mu(u)}(v_t, v_s) + \nabla_{\nabla_{A,t}v_s}X_{\mu(u)}(u)\nabla_{v_s}X_{L_u^*v_s}(u), \\
\nabla_{A,s}(R(v_s, v_t)v_t) &= \nabla R(v_s, v_s, v_t, v_t) + R(\nabla_{A,s}v_s, v_t)v_t \\
&\quad + R(v_s, \nabla_{A,s}(Jv_s))v_t) + R(v_s, v_t)\nabla_{A,s}(Jv_s).
\end{aligned}$$

Hence, by a term by term inspection, we obtain an inequality

$$\varepsilon^2\langle\nabla_{A,s}v_s, I + II + III\rangle \geq -cu_0 - v_0$$

for $\varepsilon > 0$ sufficiently small. Note, in particular, that the term $\varepsilon^2\langle\nabla_{A,s}v_s, III\rangle$ contains the two positive summands $\lambda^2|L_u^*\nabla_{A,s}v_s|^2$ and $\lambda^2|L_u^*J\nabla_{A,s}v_s|^2$. Since $\Delta u_0 \geq v_0 - c_0u_0$ the last inequality implies (91) with $c_1 := c + c_0$. Now it follows from (91), (90), Lemma 9.2, and the formula

$$\nabla_{A,t}v_s = -J\nabla_{A,s}v_s - (\nabla_{v_s}J)v_s + \frac{\lambda^2}{\varepsilon^2}L_u\mu(u)$$

that (86) holds for $p = \infty$. The estimate (86) for $2 < p < \infty$ follows by interpolation. $\square$

10 Proof of Theorem D

Theorem 10.1. *Assume (H1) and (H4), let $\bar{B} \in H_2(\bar{M};\mathbb{Z})$ be a nontorsion homology class, and denote $B := \kappa(\bar{B}) \in H_2(M_{\mathrm{G}};\mathbb{Z})$. Then, for every $C > 0$, there exist positive constants ε_0 and c_0 such that for every $\varepsilon \in (0, \varepsilon_0]$ the following holds. If $(u, A) \in \widetilde{\mathcal{M}}^{\varepsilon}_{B,\Sigma}$ satisfies $\|d_A u\|_{L^\infty} \le C$ and $u(P) \subset M^C$ then $(u, A) \in \widetilde{\mathcal{T}}^{\varepsilon}(\widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0))$.*

Proof. Suppose the assertion is false. Then there exist a constant $C > 0$ and sequences $\varepsilon_i \to 0$ and $(u_i, A_i) \in \widetilde{\mathcal{M}}^{\varepsilon_i}_{B,\Sigma}$ such that

$$\|d_{A_i} u_i\|_{L^\infty} \le C, \qquad u_i(P) \subset M^C, \qquad (u_i, A_i) \notin \widetilde{\mathcal{T}}^{\varepsilon_i}(\mathcal{M}^0_{B,\Sigma}(i)).$$

Here ε_i is chosen smaller than the number $\varepsilon_0(i)$ required for the definition of the map $\mathcal{T}^{\varepsilon_i}$. We prove in four steps that there exist an integer $i_0 \in \mathbb{N}$, positive constants c and c_0, and sequences

$$(u_{i0}, A_{i0}) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0 - 1), \qquad \zeta_{i0} = (\xi_{i0}, \alpha_{i0}) \in T_{(u_{i0}, A_{i0})}\mathcal{B}$$

such that

$$(u_i, A_i) = (\exp_{u_{i0}}(\xi_{i0}), A_{i0} + \alpha_{i0}), \qquad \|\zeta_{i0}\|_{1,p,\varepsilon_i} \le c\varepsilon_i^{2/p+1} \tag{92}$$

for every $i \ge i_0$. For i sufficiently large it then follows from Theorem 8.1 that $(u_i, A_i) \in \widetilde{\mathcal{T}}^{\varepsilon_i}(\widetilde{\mathcal{M}}^0_{B,\Sigma}(c_0))$, in contradiction to our assumption.

Step 1. *There exist constants $c > 0$ and $i_0 \in \mathbb{N}$ such that*

$$\varepsilon_i^{-1} \|\mu(u_i)\|_{L^p} + \left\|L^*_{u_i} d_{A_i} u_i\right\|_{L^p} + \left\|L^*_{u_i} J d_{A_i} u_i\right\|_{L^p} + \varepsilon_i \|\nabla_{A_i}{}^* d_{A_i} u_i\|_{L^p} \le c\varepsilon_i^{2/p}$$

for $i \ge i_0$ and $2 \le p \le \infty$.

By the graph construction in Appendix A, it suffices to establish the estimate under the hypothesis that J is independent of $z \in \Sigma$. Namely,

$$L^*_{\tilde{u}_i} d_{A_i} \tilde{u}_i = L^*_{u_i} d_{A_i} u_i, \qquad L^*_{\tilde{u}_i} \tilde{J} d_{A_i} \tilde{u}_i = L^*_{u_i} J d_{A_i} u_i,$$

$$\tilde{\nabla}_{A_i}{}^* d_{A_i} \tilde{u}_i = (\nabla_\Sigma{}^* \mathrm{id}, \nabla_{A_i}{}^* d_{A_i} u_i),$$

where $\tilde{u}_i = (\pi, u_i) : P \to \tilde{M} = \Sigma \times M$, $\tilde{\nabla}_A$ is the connection induced by A on $\tilde{u}_i^* T\tilde{M}/\mathrm{G}$, and $\mathrm{id} \in \Omega^1(\Sigma, T\Sigma)$. Hence we can use the results of Section 9. Since $\|d_{A_i} u_i\|_{L^\infty} \leq C$ and $u_i(\Omega) \subset M^C$, the pair $(\tilde{u}_i, A_i)$ satisfies the hypotheses of Lemma 9.1 and so the sequence ${\varepsilon_i}^{-3/2} \tilde{\mu}(\tilde{u}_i)$ is uniformly bounded. Hence there exists a constant $c_0 > 0$ such that

$$\|d_{A_i} u_i\|_{L^\infty} + {\varepsilon_i}^{-3/2} \|\mu(u_i)\|_{L^\infty} \leq c_0 \tag{93}$$

for every i. This implies that, in local holomorphic coordinates on Σ, the pair $(\tilde{u}_i, A_i)$ satisfies the hypotheses of Lemma 9.3 for i sufficiently large. Hence the estimate holds in local holomorphic coordinates on Σ with u_i replaced by $\tilde{u}_i$. Hence, by a partition of unity argument, it holds globally.

Step 2. *There exists an integer $i_0 \in \mathbb{N}$ and a constant $c > 0$ such that, for every $i \geq i_0$, there exists a unique $\eta_i \in \Omega^0(\Sigma, \mathfrak{g}_P)$ such that*

$$\mu(\exp_{u_i}(JL_{u_i}\eta_i)) = 0, \qquad \|\eta_i\|_{L^\infty} \leq c\,\|\mu(u_i)\|_{L^\infty}.$$

Define $u_i' : P \to M$ and $A_i' \in \mathcal{A}(P)$ by

$$u_i' := \exp_{u_i}(JL_{u_i}\eta_i), \qquad L_{u_i'}^* d_{A_i'} u_i' = 0,$$

so that $d_{A_i'} u_i' \in \Omega^1(\Sigma, H_{u_i'})$, and let $\zeta_i := (JL_{u_i}\eta_i, A_i' - A_i)$. Then there exists a constant $c' > 0$ such that

$$\|\zeta_i\|_{1,p,\varepsilon_i} \leq c' \varepsilon_i^{1+2/p}, \qquad \left\|\bar{\partial}_{J,A_i'}(u_i')\right\|_{L^p} \leq c' \varepsilon_i^{1+2/p}, \qquad \left\|d_{A_i'} u_i'\right\|_{L^\infty} \leq c'$$

for every $i \geq i_0$.

The existence of η_i for large i follows from the implicit function theorem for the map $\eta \mapsto \mu(\exp_{u_i(p)}(JL_{u_i(p)}\eta))$. This sequence satisfies an estimate of the form

$$\|\eta_i\|_{L^p} \leq c_1 \|\mu(u_i)\|_{L^p} \leq c_2 \varepsilon^{1+2/p}$$

for every $i \geq i_0$ and every $p \in [2, \infty]$. Here the constants c_1 and c_2 are independent of i and p, and the second inequality follows from Step 1. For $p = \infty$ there is actually a better estimate (by $\varepsilon^{3/2}$ instead of ε), but we shall not use this here. In the following we suppress the subscript i and write $u, u', A, A', \varepsilon$ instead of $u_i, u_i', A_i, A_i', \varepsilon_i$, respectively. We establish the required estimates in local holomorphic coordinates on Σ. As in Remark 4.1, we write $A' = \Phi'\,ds + \Psi'\,dt$ for some Lie algebra valued functiona Φ' and Ψ', and denote

$$v_s' := \partial_s u' + X_{\Phi'}(u'), \qquad v_t' := \partial_t u' + X_{\Psi'}(u').$$

Then $L_{u'}^* v_s' = L_{u'}^* v_t' = 0$. We assume that the functions $u, u', \Phi, \Psi, \Phi', \Psi'$ are defined on an open set $\Omega \subset \mathbb{C}$ and fix any compact subset $K \subset \Omega$. We must prove the estimates

$$\|(\xi, \varphi, \psi)\|_{1,p,\varepsilon} \le c\varepsilon^{1+2/p}, \quad \|v_s' + Jv_t'\|_{L^p} \le c\varepsilon^{1+2/p}, \quad \|v_s'\|_{L^\infty} + \|v_t'\|_{L^\infty} \le c$$

on the subset K, where

$$\xi := JL_u\eta, \qquad \varphi := \Phi' - \Phi, \qquad \psi := \Psi' - \Psi.$$

Abbreviate $E_i := E_i(u, JL_u\eta)$, $i = 1, 2$. Then

$$\nabla_{A,t}\xi = JL_u\nabla_{A,t}\eta + (\nabla_{v_t}J + \partial_t J)L_u\eta + J\nabla_{v_t}X_\eta(u). \tag{94}$$

Hence, by Lemma C.3,

$$v_t' - E_1 v_t = L_{u'}\psi + E_2\nabla_{A,t}\xi = L_{u'}\psi + E_2 JL_u\nabla_{A,t}\eta + R_t\eta, \tag{95}$$

where

$$R_t\eta := E_2(\nabla_{v_t}J + \partial_t J)L_u\eta + E_2 J\nabla_{v_t}X_\eta(u).$$

Hence

$$d\mu(u')(v_t' - E_1 v_t) = d\mu(u')E_2 JL_u\nabla_{A,t}\eta + d\mu(u')R_t\eta.$$

Since $L_u^* = d\mu(u)J$ we have

$$\begin{aligned} L_u^* L_u\nabla_{A,t}\eta &= (d\mu(u) - d\mu(u')E_2)JL_u\nabla_{A,t}\eta \\ &\quad + d\mu(u')(v_t' - E_1 v_t) - d\mu(u')R_t\eta, \end{aligned}$$

and, since $d\mu(u')v_t' = 0$,

$$\begin{aligned} L_u^* L_u\nabla_{A,t}\eta &= (d\mu(u) - d\mu(u')E_2)JL_u\nabla_{A,t}\eta \\ &\quad + (d\mu(u) - d\mu(u')E_1)v_t + L_u^* Jv_t - d\mu(u')R_t\eta. \end{aligned}$$

It follows that

$$\begin{aligned} \|\nabla_{A,t}\eta\|_{L^p} &\le c_2\left(\|\eta\|_{L^\infty}\|\nabla_{A,t}\eta\|_{L^p} + \|L_u^* Jv_t\|_{L^p} + \|\eta\|_{L^p}\right) \\ &\le c_3\left(\varepsilon^{1+2/p}\|\nabla_{A,t}\eta\|_{L^p} + \varepsilon^{2/p}\right). \end{aligned}$$

If ε is sufficiently small this gives

$$\|\nabla_{A,t}\eta\|_{L^p} \le c_4\varepsilon^{2/p}, \qquad \|\nabla_{A,t}\eta\|_{L^\infty} \le c_4.$$

Here the second inequality follows from a similar argument as the first. Combining these inequalities with (94) we obtain

$$\|\nabla_{A,t}\xi\|_{L^p} \le c_5\varepsilon^{2/p}, \qquad \|\nabla_{A,t}\xi\|_{L^\infty} \le c_5.$$

In order to estimate ψ we apply the operator $L_{u'}^*$ to (95) and use the formula $L_{u'}^*v_t' = 0$ to obtain

$$L_{u'}^*L_{u'}\psi = (L_u^* - L_{u'}^*E_1)v_t - L_u^*v_t - L_{u'}^*E_2JL_u\nabla_{A,t}\eta - L_{u'}^*R_t\eta.$$

Combining this with Step 1 and the estimate for $\nabla_{A,t}\eta$ we obtain

$$\|\psi\|_{L^p} \le c_6\varepsilon^{2/p}, \qquad \|\psi\|_{L^\infty} \le c_6.$$

Hence, by (95),

$$\|v_t' - E_1v_t\|_{L^p} \le c_7\varepsilon^{2/p}, \qquad \|v_t' - E_1v_t\|_{L^\infty} \le c_7.$$

Similarly,

$$\|\nabla_{A,s}\eta\|_{L^p} + \|\varphi\|_{L^p} + \|v_s' - E_1v_s\|_{L^p} \le c_7\varepsilon^{2/p}$$

and

$$\|\nabla_{A,s}\eta\|_{L^\infty} + \|\varphi\|_{L^\infty} + \|v_s' - E_1v_s\|_{L^\infty} \le c_7.$$

Now use (95) again to obtain

$$\begin{aligned} v_s' + J(u')v_t' &= L_{u'}(\varphi - \nabla_{A,t}\eta) + JL_{u'}(\psi + \nabla_{A,s}\eta) \\ &\quad + (E_2JL_u - JL_{u'})\nabla_{A,s}\eta + J(E_2JL_u - JL_{u'})\nabla_{A,t}\eta \\ &\quad + E_1v_s + JE_1v_t + (R_s + JR_t)\eta \\ &= \pi_{u'}\big((E_2JL_u - JL_{u'})\nabla_{A,s}\eta + J(E_2JL_u - JL_{u'})\nabla_{A,t}\eta\big) \\ &\quad + \pi_{u'}\big((JE_1 - E_1J)v_t + (R_s + JR_t)\eta\big). \end{aligned}$$

The second equality uses the fact that $v_s + Jv_t = 0$ and that the 1-form $\bar\partial_{J,A'}(u')$ takes values in $H_{u'}$. It follows that

$$\|v_s' + J(u')v_t'\|_{L^p} \le c_8\varepsilon^{1+2/p}.$$

It remains to show that

$$\|\nabla_{A,s}\varphi\|_{L^p} + \|\nabla_{A,t}\varphi\|_{L^p} + \|\nabla_{A,s}\psi\|_{L^p} + \|\nabla_{A,t}\psi\|_{L^p} \le c\varepsilon^{2/p-1}.$$

To estimate the term $\nabla_{A,t}\psi$ differentiate (95) with respect to t. Then apply the operator $d\mu(u')$ to the resulting expression to eliminate $\nabla_{A,t}\psi$ and obtain an estimate of the form

$$\|\nabla_{A,t}\nabla_{A,t}\eta\|_{L^p} \le c_9\varepsilon^{2/p-1}.$$

Then apply the operator $L^*_{u'}$ to the equation obtained from differentiating (95), and estimate $\nabla_{A,t}\psi$ using the upper bound found for $\nabla_{A,t}\nabla_{A,t}\eta$. The estimate for $\nabla_{A,s}\psi$ is obtained in a similar manner. To estimate $\nabla_{A,s}\varphi$ and $\nabla_{A,t}\varphi$, we begin with the identity

$$v'_s - E_1 v_s = L_{u'}\varphi + E_2 J L_u \nabla_{A,s}\eta + R_s\eta$$

instead of (95) and then follow the same procedure.

Step 3. *There exist an integer* $i_0 \in \mathbb{N}$*, a constant* $c > 0$*, and a sequence* $(u''_i, A''_i) \in \widetilde{\mathcal{M}}^0_{B,\Sigma}$ *such that*

$$u''_i = \exp_{u'_i}(\xi'_i), \qquad \xi'_i \in \Omega^0(\Sigma, H_{u'_i}),$$

and

$$\|\xi'_i\|_{W^{1,p}} + \|A''_i - A'_i\|_{L^p} \le c\varepsilon_i^{1+2/p}, \qquad \left\|d_{A''_i}u''_i\right\|_{L^\infty} \le c,$$

for $i \ge i_0$.

By Step 2,

$$\sup_i \left\|d_{A'_i}u'_i\right\|_{L^\infty} < \infty.$$

Hence the induced maps $\bar{u}'_i : \Sigma \to \bar{M}$ form a sequence of approximate $\bar{J}$-holomorphic curves which satisfy a uniform L^∞-bound on their first derivatives. Hence, by $(H4)$ and Theorem 2.3, there is nearby a true $\bar{J}$-holomorphic curve $\bar{u}''_i : \Sigma \to \bar{M}$ whose $W^{1,p}$-distance to $\bar{u}'_i$ is controlled by the L^p-norm of $\bar{\partial}_{\bar{J}}(\bar{u}'_i)$. Now this $\bar{J}$-holomorphic curve has a unique lift $u''_i : P \to \mu^{-1}(0)$ of the form

$$u''_i = \exp_{u'_i}(\xi'_i), \qquad \xi'_i \in \Omega^0(\Sigma, H_{u'_i}).$$

Let $A''_i \in \mathcal{A}(P)$ be the connection determined by u''_i via $L^*_{u''_i} d_{A''_i} u''_i = 0$. Then

$$\|\xi'_i\|_{W^{1,p}} \le c_1 \left\|\bar{\partial}_{J,A'_i}(u'_i)\right\|_{L^p} \le c_2 {\varepsilon_i}^{1+2/p}.$$

Here the last inequality follows from Step 2. Since $L^*_{u_i'}d_{A_i'}u_i' = L^*_{u_i''}d_{A_i''}u_i'' = 0$ we obtain

$$\|A_i'' - A_i'\|_{L^p} \le c_3\varepsilon_i^{1+2/p}.$$

In particular, these inequalities together give a uniform $W^{1,p}$-bound on the $\bar{J}$-holomorphic curves $\bar{u}_i'' : \Sigma \to \bar{M}$. Hence, by the elliptic bootstrapping techniques for J-holomorphic curves, the sequence $\bar{u}_i''$ satisfies a uniform L^∞ bound on the first derivatives. This proves Step 3.

Unfortunately, the estimate on $A_i'' - A_i'$ in Step 3 is only in the L^p-norm and not in the $W^{1,p}$-norm. A further modification of the pair (u_i'', A_i'') is required to improve this estimate.

Step 4. *There exist an integer $i_0 \in \mathbb{N}$, a constant $c > 0$, and a sequence of gauge transformations $g_i \in \mathcal{G}(P)$ such that the sequence*

$$(u_{i0}, A_{i0}) := g_i^*(u_i'', A_i'') \in \widetilde{\mathcal{M}}^0_{B,\Sigma}$$

satisfies the following. For $i \ge i_0$ the original sequence (u_i, A_i) has the form

$$(u_i, A_i) = (\exp_{u_{i0}}(\xi_i), A_{i0} + \alpha_i)$$

where $\zeta_i := (\xi_i, \alpha_i) \in T_{(u_{i0},A_{i0})}\mathcal{B}$ satisfies (92).

The idea is to choose g_i for large i such that

$$u_i = \exp_{u_{i0}}(\xi_i), \qquad L^*_{u_{i0}}\xi_i = 0, \qquad u_{i0} := g_i^{-1}u_i''.$$

This can be done by using pointwise, for every $p \in P$, the implicit function theorem to obtain the local slice condition. This suffices to obtain the missing estimates for the first derivatives of $g_i^*A_i'' - A_i$. We sketch a proof of this estimate below.

By (93) and Step 2, the distance between u_i and u_i' is unformly bounded by a constant times $\varepsilon_i^{3/2}$ while the distance in the $W^{1,p}$-norm is bounded by a constant times $\varepsilon_i^{2/p}$. By Step 3, the distance between u_i' and u_i'' is bounded in the $W^{1,p}$-norm by a constant times $\varepsilon_i^{1+2/p}$. Hence there exists a sequence of smooth sections $\xi_i \in \Omega^0(\Sigma, u_i{}^*TM/\mathrm{G})$ and a constant $c > 0$ such that

$$u_i'' = \exp_{u_i}(\xi_i), \qquad \|\xi_i\|_{L^\infty} \le c_1\varepsilon_i, \qquad \|\nabla_{A_i}\xi_i\|_{L^p} \le c_1\varepsilon_i^{2/p}. \tag{96}$$

Moreover, the sequence $d_{A_i''}u_i''$ is uniformly bounded in the L^∞-norm and

$$\|A_i'' - A_i\|_{L^p} \le c_1\varepsilon^{2/p}, \qquad \left\|d_{A_i''}u_i'' - E_1(u_i,\xi_i)d_{A_i}u_i\right\|_{L^p} \le c_1\varepsilon^{2/p}. \tag{97}$$

Here the last inequality follows from the identity

$$\begin{aligned} d_{A_i''}u_i'' &= E_1(u_i,\xi_i)d_{A_i}u_i + E_2(u_i,\xi_i)\nabla_{A_i}\xi_i \\ &\quad + E_1(u_i,\xi_i)X_{A_i''-A_i}(u_i) + E_2(u_i,\xi_i)\nabla_{\xi_i}X_{A_i''-A_i}(u_i), \end{aligned}$$

which in turn follows from Lemma C.1. Now, by the inverse function theorem for the map $\mathrm{G}\times\ker L_x^* \to M : (g,\xi)\mapsto g^{-1}\exp_x(\xi)$, there exists a constant $c_2>0$ and (unique) sequences $g_i\in\mathcal{G}(P)$ and $\xi_i''\in\Omega^0(\Sigma, u_i''^*TM/\mathrm{G})$ such that

$$u_i = g_i^{-1}\exp_{u_i''}(\xi_i''), \qquad L_{u_i''}^*\xi_i'' = 0,$$

and

$$\|\xi_i''\|_{L^\infty} \le c_2\varepsilon_i, \qquad \|g_i - \mathbb{1}\|_{L^\infty} \le c_2\varepsilon_i.$$

Define

$$u_{i0} := g_i^{-1}u_i'', \qquad A_{i0} := g_i{}^*A_i'', \qquad \xi_{i0} := g_i^{-1}\xi_i'', \qquad \alpha_{i0} := A_i - A_{i0}.$$

We shall prove that the pair (ξ_{i0},α_{i0}) satisfies (92). To see this note first that

$$u_i = \exp_{u_{i0}}(\xi_{i0}), \qquad A_i = A_{i0} + \alpha_{i0}.$$

The endomorphism $E_1(u_i,\xi_i)g_i^{-1}E_1(u_i'',\xi_i'')$ of $u_i''^*TM$ is ε_i-close to the identity, $d_{A_{i0}}u_{i0} = g_i^{-1}d_{A_i''}u_i''$, and

$$\begin{aligned} &E_1(u_i,\xi_i)(d_{A_i}u_i - E_1(u_{i0},\xi_{i0})d_{A_{i0}}u_{i0}) \\ &= E_1(u_i,\xi_i)d_{A_i}u_i - d_{A_i''}u_i'' + (\mathbb{1} - E_1(u_i,\xi_i)g^{-1}E_1(u_i'',\xi_i''))d_{A_i''}u_i''. \end{aligned}$$

Hence, by (97), there is an estimate

$$\|d_{A_i}u_i - E_1(u_{i0},\xi_{i0})d_{A_{i0}}u_{i0}\|_{L^p} \le c_3\varepsilon_i^{2/p}, \qquad \|d_{A_{i0}}u_{i0}\|_{L^\infty} \le c_3$$

for all i. Hence, by Corollary C.4, there exists a constant $c_4>0$ such that

$$\|\alpha_{i0}\|_{L^p} \le c_4\varepsilon_i^{2/p}, \qquad \|\alpha_{i0}\|_{L^\infty} \le c_4.$$

Next observe that, by Lemma C.1,

$$\begin{aligned} d_{A_i}u_i &= E_1(u_{i0},\xi_{i0})d_{A_{i0}}u_{i0} + E_2(u_{i0},\xi_{i0})\nabla_{A_{i0}}\xi_{i0} \\ &\quad + E_1(u_{i0},\xi_{i0})X_{\alpha_{i0}}(u_{i0}) + E_2(u_{i0},\xi_{i0})\nabla_{\xi_{i0}}X_{\alpha_{i0}}(u_{i0}) \end{aligned}$$

Hence there exists a constant $c_5 > 0$ such that

$$\|\nabla_{A_{i0}}\xi_{i0}\|_{L^p} \le c_5\varepsilon_i^{2/p}, \qquad \|\nabla_{A_{i0}}\xi_{i0}\|_{L^\infty} \le c_5$$

for all i. Thus we have proved that

$$\begin{array}{rcl} \|\xi_{i0}\|_{L^p} + \varepsilon_i\,\|\alpha_{i0}\|_{L^p} + \varepsilon_i\,\|\nabla_{A_{i0}}\xi_{i0}\|_{L^p} & \le & c_6\varepsilon_i^{1+2/p}, \\ \|\xi_{i0}\|_{L^\infty} + \varepsilon_i\,\|\alpha_{i0}\|_{L^\infty} + \varepsilon_i\,\|\nabla_{A_{i0}}\xi_{i0}\|_{L^\infty} & \le & c_6\varepsilon_i \end{array} \tag{98}$$

for all i. It remains to estimate the L^p-norm of the first derivatives of α_{i0}. For this we shall drop the subscript i and write $u, u_0, A, A_0, \xi_0, \alpha_0$ instead of $u_i, u_{i0}, A_i, A_{i0}, \xi_{i0}, \alpha_{i0}$. Moreover, we use local coordinates on Σ as in Step 2 and write

$$\begin{array}{rclrcl} A_0 & := & \Phi_0\,ds + \Psi_0\,dt, & A & := & \Phi\,ds + \Psi\,dt, \\ v_{0s} & := & \partial_s u_0 + L_{u_0}\Phi_0, & v_s & := & \partial_s u + L_u\Phi, \\ v_{0t} & := & \partial_t u_0 + L_{u_0}\Psi_0, & v_t & := & \partial_t u + L_u\Psi, \end{array}$$

and $\varphi_0 := \Phi - \Phi_0$ and $\psi_0 := \Psi - \Psi_0$. Consider the formula

$$\nabla_{A,s}v_s + J\nabla_{A,t}v_s = -(\nabla_{v_t}J)v_s + \frac{\lambda^2}{\varepsilon^2}JL_u\mu(u).$$

By Step 1, we have

$$\|\nabla_{A,s}v_s + J\nabla_{A,t}v_s\|_{L^p} \le c_7\varepsilon^{2/p-1}$$

and hence, by elliptic regularity for the Cauchy-Riemann operator,

$$\|\nabla_{A,s}v_s\|_{L^p} + \|\nabla_{A,t}v_s\|_{L^p} \le c_8\varepsilon^{2/p-1}. \tag{99}$$

Moreover, since $L_{u_0}^*\xi_0 = 0$, it follows from Lemma C.2 that

$$\begin{array}{rcl} L_{u_0}^*\nabla_{A_0,t}\nabla_{A_0,s}\xi_0 & = & \nabla_{A_0,t}(L_{u_0}^*\nabla_{A_0,s}\xi_0) - \rho(v_{0t}, \nabla_{A_0,s}\xi_0) \\ & = & -\nabla_{A_0,t}\rho(v_{0s}, \xi_0) - \rho(v_{0t}, \nabla_{A_0,s}\xi_0). \end{array}$$

and hence

$$\left\|L_{u_0}^*\nabla_{A_0,t}\nabla_{A_0,s}\xi_0\right\| \le c_{10}\varepsilon^{2/p}. \tag{100}$$

Here we use the fact that, by elliptic bootstrapping for $\bar{J}$-holomorphic curves, there is a uniform L^p-bound on $\nabla_{A_0,t}v_{0s}$. Now consider the pointwise inequality

$$\begin{array}{rcl} |\nabla_{A,t}\varphi_0| & \le & c_{11}|L_u^*L_u\nabla_{A,t}\varphi_0| \\ & \le & c_{11}|(L_u^* - L_{u_0}^*E_2^{-1})L_u\nabla_{A,t}\varphi_0| + c_{11}|L_{u_0}^*E_2^{-1}L_u\nabla_{A,t}\varphi_0| \end{array}$$

Since the operator $(L_u^* - L_{u_0}^* E_2^{-1})L_u$ is small, we obtain

$$\|\nabla_{A,t}\varphi_0\|_{L^p} \le c_{12} \left\| L_{u_0}^* E_2^{-1} L_u \nabla_{A,t}\varphi_0 \right\|_{L^p}.$$

Now use Lemma C.5 and the estimates (98), (99), and (100) to obtain

$$\|\nabla_{A,t}\varphi_0\|_{L^p} \le c_{13}\varepsilon^{2/p-1}.$$

The terms $\|\nabla_{A,s}\varphi_0\|_{L^p}$, $\|\nabla_{A,t}\psi_0\|_{L^p}$, and $\|\nabla_{A,s}\varphi_0\|_{L^p}$ are estimated similarly. This proves Step 4.

It follows from Step 4 and Theorem 8.1 that $(u_i, A_i) \in \widetilde{\mathcal{T}}_i^\varepsilon(\widetilde{\mathcal{M}}_{B,\Sigma}^0(c_0)$ for some constant c_0 and i sufficiently large. This contradicts our assumption and hence proves the theorem. □

11 Vortices

In this section we examine the finite energy solutions of (1) over the complex plane $\Sigma = \mathbb{C}$. The equations have the form

$$\begin{aligned} \partial_s u + X_\Phi(u) + J(\partial_t u + X_\Psi(u)) &= 0, \\ \partial_s \Psi - \partial_t \Phi + [\Phi, \Psi] + \mu(u) &= 0, \end{aligned} \tag{101}$$

where $u : \mathbb{C} \to M$ and $\Phi, \Psi : \mathbb{C} \to \mathfrak{g}$. The energy of the triple (u, Φ, Ψ) is given by

$$E(u, \Phi, \Psi) := \int_{\mathbb{C}} \left(|\partial_s u + X_\Phi(u)|^2 + |\mu(u)|^2 \right) dsdt.$$

The vortex equations (101) and the energy are invariant under the action of the gauge group

$$\mathcal{G} := C^\infty(\mathbb{C}, \mathrm{G})$$

by

$$g^*(u, \Phi, \Psi) := (g^{-1}u, g^{-1}\partial_s g + g^{-1}\Phi g, g^{-1}\partial_t g + g^{-1}\Psi g).$$

A solution of (101) is said to be in **radial gauge** if

$$\cos\theta\, \Phi(re^{i\theta}) + \sin\theta\, \Psi(re^{i\theta}) = 0$$

for every $\theta \in \mathbb{R}$ and every sufficiently large $r \ge 0$. It is said to be **bounded** if $\sup_{\mathbb{C}} |\mu(u)| < \infty$.

Proposition 11.1. *Assume* $(H1)$ *and* $(H2)$. *Suppose that* (u, Φ, Ψ) *is a smooth bounded finite energy solution of (101) in radial gauge. Then there exists a* $W^{1,2}$*-function* $x : \mathbb{R}/2\pi\mathbb{Z} \to M$ *and an* L^2*-function* $\eta : \mathbb{R}/2\pi\mathbb{Z} \to \mathfrak{g}$ *such that*

$$\dot{x} + X_\eta(x) = 0, \qquad \mu(x) = 0 \tag{102}$$

and

$$\lim_{r\to\infty} \sup_{\theta\in\mathbb{R}} d(u(re^{i\theta}), x(\theta)) = 0, \qquad \lim_{r\to\infty} \int_0^{2\pi} |\eta(\theta) - \eta_r(\theta)|^2 \, d\theta = 0, \tag{103}$$

where $\eta_r(\theta) := r\cos(\theta)\Psi(re^{i\theta}) - r\sin\theta\Phi(re^{i\theta})$. *Moreover, there exists a constant* $\delta > 0$ *such that*

$$\lim_{r\to\infty} \sup_{\theta\in\mathbb{R}} r^{2+\delta} \Big(|\partial_s u + X_\Phi(u)|^2 + |\mu(u)|^2 \Big) = 0,$$

where $s + it =: re^{i\theta}$,

$$E(u, \Phi, \Psi) = \int_{\mathbb{C}} u^*\omega,$$

and $\sup_{\mathbb{C}}(f \circ u) \le c$, *where* c *is as in hypothesis* $(H2)$ *and* $\mu^{-1}(0) \subset f^{-1}([0, c])$. *If* $(H3)$ *holds then* $E(u, \Phi, \Psi)$ *is an integer multiple of* $\hbar = \tau N$.

Note that the removable singularity theorem for J-holomorphic curves is a corollary of Proposition 11.1 (consider the special case $\mathrm{G} = \{\mathbb{1}\}$ and $M = \bar{M}$). Before entering into the proof we introduce the notion of the local equivariant symplectic action. The definition of this local action functional relies on the following lemma. We identify $S^1 \cong \mathbb{R}/2\pi\mathbb{Z}$.

Lemma 11.2. *Assume* $(H1)$. *Then there exist positive constants* δ *and* c *such that, for every pair of smooth loops* $x : S^1 \to M$ *and* $\eta : S^1 \to \mathfrak{g}$ *such that*

$$\sup_{S^1} |\mu(x)| < \delta,$$

there exists a point $x_0 \in \mu^{-1}(0)$ *and a smooth loop* $g_0 : S^1 \to \mathrm{G}$ *such that*

$$c^{-1} \sup_{S^1} \left|\eta + \dot{g}_0 {g_0}^{-1}\right| \le \ell(x, \eta) := \int_0^{2\pi} |\dot{x} + X_\eta(x)| \, d\theta,$$

and

$$d(x(\theta), g_0(\theta)x_0) \le c\,(|\mu(\theta)| + \ell(x, \eta))$$

for every $\theta \in S^1$.

Proof. Fix a G-invariant and ω-compatible almost complex structure $J \in \mathcal{J}_{\mathrm{G}}(M,\omega)$. If δ is sufficiently small then there exist unique loops $x_0 : S^1 \to \mu^{-1}(0)$ and $\eta_0 : S^1 \to \mathfrak{g}$ such that

$$x(\theta) = \exp_{x_0(\theta)}(\xi_0(\theta)), \qquad \xi_0 := JL_{x_0}\eta_0, \qquad |\xi_0| \le c_1|\mu(x)|.$$

By Lemmata C.1 and C.2, we have

$$\dot{x} + X_\eta(x) = E_1(x_0,\xi_0)(\dot{x}_0 + X_\eta(x_0)) + E_2(x_0,\xi_0)(\nabla\xi_0 + \nabla_{\xi_0}X_\eta(x_0)),$$

$$\nabla\xi_0 + \nabla_{\xi_0}X_\eta(x_0) = JL_{x_0}(\dot{\eta}_0 + [\eta,\eta_0]) + \nabla_{\dot{x}_0+X_\eta(x_0)}(JX_{\eta_0})(x_0).$$

Since the image of JL_{x_0} is the orthogonal complement of the kernel of $d\mu(x_0)$ we deduce that there exists a constant $c_2 > 0$ such that

$$|\dot{x}_0 + X_\eta(x_0)| + |\nabla\xi_0 + \nabla_{\xi_0}X_\eta(x_0)| \le c_2\left|\dot{x} + X_\eta(x)\right|,$$

pointwise for every $\theta \in S^1$. Define $g : \mathbb{R} \to \mathrm{G}$ and $y_0 : \mathbb{R} \to \mu^{-1}(0)$ by

$$\dot{g} + \eta g = 0, \qquad g(0) = \mathbb{1}, \qquad y_0(\theta) := g(\theta)^{-1}x_0(\theta).$$

Then

$$g(\theta + 2\pi) = g(\theta)g(2\pi), \qquad \dot{y}_0 = g^{-1}(\dot{x}_0 + X_\eta(x_0)).$$

Hence $d(y_0(2\pi), x_0(0)) \le 2\pi c_2\ell(x,\eta)$ and so $d(g(2\pi),\mathbb{1}) \le c_3\ell(x,\eta)$. This implies that there exists a path $h : \mathbb{R} \to \mathrm{G}$ such that

$$h(\theta + 2\pi) = h(\theta)g(2\pi), \qquad \sup_{[0,2\pi]}\left|h^{-1}\dot{h}\right| \le c_4\ell(x,\eta).$$

Hence $g_0 := gh^{-1}$ is a loop and

$$\sup_{S^1}\left|\eta + \dot{g}_0{g_0}^{-1}\right| = \sup_{[0,2\pi]}\left|h^{-1}\dot{h}\right| \le c_4\ell(x,\eta).$$

Moreover, with $x_0 := x_0(0)$, we obtain

$$\begin{aligned} d(x(\theta), g_0(\theta)x_0) &\le d(x(\theta), x_0(\theta)) + d(x_0(\theta), g(\theta)h(\theta)^{-1}x_0) \\ &\le d(x(\theta), x_0(\theta)) + d(y_0(\theta), x_0) + d(h(\theta)x_0, x_0) \\ &\le c_1|\mu(x(\theta))| + c_5\ell(x,\eta). \end{aligned}$$

This proves the lemma. □

Let δ be as in Lemma 11.2 and $(x, \eta) : S^1 \to M \times \mathfrak{g}$ be a loop such that $\sup_{S^1} |\mu(x)| < \delta$. Then the **local equivariant symplectic action** of the pair (x, η) is defined by

$$\mathcal{A}(x, \eta) := -\int u^*\omega + \int_0^{2\pi} \langle \mu(x(\theta)), \eta(\theta) \rangle \, d\theta,$$

where $x_0 \in \mu^{-1}(0)$ and $g_0 : S^1 \to \mathrm{G}$ are as in Lemma 11.2, $\xi_0(\theta) \in T_{g_0(\theta)x_0}M$ is the unique small tangent vector such that

$$x(\theta) = \exp_{g_0(\theta)x_0}(\xi_0(\theta)),$$

and $u : [0, 1] \times S^1 \to M$ is defined by

$$u(\tau, \theta) := \exp_{g_0(\theta)x_0}(\tau\xi_0(\theta)).$$

The local action is independent of the choice of x_0 and g_0 so long as the distance between $x(\theta)$ and $g_0(\theta)x_0$ remains sufficiently small.

Lemma 11.3. *Assume (H1). There exist positive constants δ and c such that the following holds. If $(x, \eta) : S^1 \to M$ is a smooth loop such that $\sup_{S^1} |\mu(x)| < \delta$ then*

$$|\mathcal{A}(x, \eta)| \leq c \int_0^{2\pi} \left(|\dot{x} + X_\eta(x)|^2 + |\mu(x)|^2 \right) \, d\theta.$$

Proof. Let $\xi_0 \in C^\infty(S^1, x_0^*TM)$ and $u : [0, 1] \times S^1 \to M$ be as above. Then the local equivariant symplectic action can be expressed in the form

$$\mathcal{A}(x, \eta) = \int_0^1 \int_0^{2\pi} \omega(\partial_\tau u, \partial_\theta u + X_\eta(u)) \, d\theta d\tau.$$

By Lemma 11.2, we have the pointwise inequality

$$|\partial_\tau u| = |\xi_0| = d(x, g_0x_0) \leq c_1 \left(|\mu(x)| + \ell(x, \eta) \right).$$

Moreover, by Lemma C.1,

$$\partial_\theta u + X_\eta(u) = E_1 L_{g_0x_0}(\eta + \dot{g}_0 {g_0}^{-1}) + \tau E_2 \left(\nabla\xi_0 + \nabla_{\xi_0} X_\eta(g_0x_0) \right),$$

where $E_i := E_i(g_0x_0, \tau\xi_0)$ for $i = 1, 2$. With $\tau = 1$ we obtain, by Lemma 11.2,

$$|\nabla\xi_0 + \nabla_{\xi_0} X_\eta(g_0x_0)| \leq c_2 \left(|\dot{x} + X_\eta(x)| + \ell(x, \eta) \right)$$

and this implies

$$|\partial_\theta u + X_\eta(u)| \le c_3 \left(|\dot{x} + X_\eta(x)| + \ell(x,\eta)\right).$$

Hence

$$\begin{aligned}|\mathcal{A}(x,\eta)| &\le c_1 c_3 \int_0^{2\pi} \big(|\mu(x)| + \ell(x,\eta)\big)\big(|\dot{x} + X_\eta(x)| + \ell(x,\eta)\big)\, d\theta \\ &\le c_4 \int_0^{2\pi} \left(|\dot{x} + X_\eta(x)|^2 + |\mu(x)|^2\right)\, d\theta.\end{aligned}$$

This proves the lemma. □

Proof of Proposition 11.1. Let (f, J) be as in $(H2)$ and let (u, Φ, Ψ) be a finite energy solution of (101) in radial gauge. We prove in seven steps that (u, Φ, Ψ) has the properties asserted in the proposition.

Step 1. $\lim_{r\to\infty} r^2\Big(|\partial_s u + X_\Phi(u)|^2 + |\mu(u)|^2\Big) = 0$ *uniformly in* θ.

Abbreviate $v_s := \partial_s u + L_u\Phi$ and $v_t := \partial_t u + L_u\Psi$ as in Section 9. Let

$$e := \frac{1}{2}(|v_s|^2 + |\mu(u)|^2).$$

Then the formula (89) with $\lambda = \varepsilon = 1$ has the form

$$\begin{aligned}\Delta e &= |\nabla_{A,s} v_s|^2 + |\nabla_{A,t} v_s|^2 + 2|L_u^* v_s|^2 + 2|L_u^* J v_s|^2 + |L_u \mu(u)|^2 \\ &\quad - 5\langle \mu(u), \rho(v_s, v_t)\rangle + \langle v_s, (\nabla_{v_s} J) L_u \mu(u)\rangle - \langle v_s, R(v_s, v_t) v_t\rangle.\end{aligned}$$

Since $u(\mathbb{C})$ is contained in a compact subset of M this gives an inequality $\Delta e \ge -c_1 e^2$. Namely, choose $\delta > 0$ such that L_x is injective whenever $|\mu(x)|^2 < \delta$. Then the first term in the secomd row can be estimated from below by $-|L_u\mu(u)|^2/2 - c|v_s|^4$ whenever $e \le \delta$. In case $e \ge \delta$ we can use the inequalities $\Delta e \ge -c(e + e^2)$ and $e \le e^2/\delta$. Now it follows from [MS1, Lemma 4.3.2] that there is a constant $c_2 > 0$ such that

$$|z| - r \ge c_2 \qquad \Longrightarrow \qquad e(z) \le \frac{8}{\pi r^2}\int_{B_r(z)} e.$$

With $r := |z|/2$ this implies $\lim_{|z|\to\infty} |z|^2 e(z) = 0$.

Step 2. *For $R > 0$ sufficiently large, we have*

$$E(u, \Phi, \Psi; \mathbb{C} \setminus B_R) = \mathcal{A}(x_R, \eta_R),$$

where $x_R(\theta) := u(Re^i\theta)$ *and* $\eta_R(\theta) := R\cos\theta\, \Psi(Re^{i\theta}) - R\sin\theta\, \Phi(Re^{i\theta})$.

The energy identity on $B_R = \{|z| \leq R\}$ has the form

$$E(u, \Phi, \Psi; B_R) = \int_{B_R} u^*\omega - \int_0^{2\pi} \langle \mu(x_R(\theta)), \eta_R(\theta)\rangle \, d\theta.$$

For R sufficiently large denote by $u_R : [0,1] \times S^1 \to M$ the function used in the definition of the local symplectic action of (x_R, η_R). Then $u_R(1, \theta) = x_R(\theta)$ and $u_R(0, \theta) = g_R(\theta)x_{R0}$ for some point $x_{R0} \in \mu^{-1}(0)$ and some loop $g_R : S^1 \to \mathrm{G}$. The homotopy class of the connected sum $v_R := u|_{B_R}\#(-u_R)$ (the orientation of u_R is reversed) is independent of R. Hence the number

$$\int {v_R}^*\omega = E(u, \Phi, \Psi; B_R) + \mathcal{A}(x_R, \eta_R)$$

is independent of R. Since $\mathcal{A}(x_R, \eta_R)$ tends to zero as $R \to \infty$ it follows that

$$E(u, \Phi, \Psi; B_R) + \mathcal{A}(x_R, \eta_R) = E(u, \Phi, \Psi)$$

for every sufficiently large number R. This proves Step 2.

Step 3. $\sup_{\mathbb{C}}(f \circ u) \leq c$.

Suppose, by contradiction, that $\sup_{\mathbb{C}}(f \circ u) > c$. Then there exists a regular value a of $f \circ u$ such that $c < a < \sup_{\mathbb{C}}(f \circ u)$. Hence the set $U := \{z \in \mathbb{C} \,|\, f(u(z)) \geq a\}$ is a smooth submanifold of $\mathbb{C}$ with boundary. Since $\mu^{-1}(0) \subset f^{-1}([0, c])$ it follows from Step 1 that there exists a number $R > 0$ such that

$$\sup_{\mathbb{C}\setminus B_R} (f \circ u) < a < \sup_{\mathbb{C}} (f \circ u).$$

Hence U is compact and has a nonempty boundary. By $(H2)$, $\Delta(f \circ u) \geq 0$ in U (see [CGMS]). Hence

$$0 \leq \int_U \Delta(f \circ u) = \int_{\partial U} \frac{\partial(f \circ u)}{\partial \nu} < 0.$$

This contradiction proves Step 3.

Step 4. *Consider equation (101) in polar coordinates* $s + it = e^{\tau + i\theta}$. *Define* $\tilde{u} : \mathbb{R} \times S^1 \to M$ *and* $\tilde{\Phi}, \tilde{\Psi} : \mathbb{R} \times S^1 \to \mathfrak{g}$ *by*

$$\begin{array}{rcl} \tilde{u}(\tau, \theta) & := & u(e^{\tau + i\theta}), \\ \tilde{\Phi}(\tau, \theta) & := & e^\tau \cos\theta \, \Phi(e^{\tau + i\theta}) + e^\tau \sin\theta \, \Psi(e^{\tau + i\theta}), \\ \tilde{\Psi}(\tau, \theta) & := & e^\tau \cos\theta \, \Psi(e^{\tau + i\theta}) - e^\tau \sin\theta \, \Phi(e^{\tau + i\theta}). \end{array}$$

Then $ds \wedge dt = e^{2\tau} d\tau \wedge d\theta$, $\Phi \, ds + \Psi \, dt = \tilde{\Phi} \, d\tau + \tilde{\Psi} \, d\theta$, *and (101) is equivalent to*

$$\tilde{v}_\tau + J \tilde{v}_\theta = 0, \qquad \tilde{\kappa} + e^{2\tau} \mu(\tilde{u}) = 0,$$

where

$$\tilde{v}_\tau := \partial_\tau \tilde{u} + L_{\tilde{u}} \tilde{\Phi}, \qquad \tilde{v}_\theta := \partial_\theta \tilde{u} + L_{\tilde{u}} \tilde{\Psi}, \qquad \tilde{\kappa} := \partial_\tau \tilde{\Psi} - \partial_\theta \tilde{\Phi} + [\tilde{\Phi}, \tilde{\Psi}].$$

The radial gauge condition has the form $\tilde{\Phi}(\tau, \theta) = 0$ *for large* τ. *The energy of the triple* $(\tilde{u}, \tilde{\Phi}, \tilde{\Psi})$ *is given by*

$$E(\tilde{u}, \tilde{\Phi}, \tilde{\Psi}) = \int_{-\infty}^{\infty} \int_0^{2\pi} \left(|\tilde{v}_\tau|^2 + e^{2\tau} |\mu(\tilde{u})|^2 \right) d\theta d\tau.$$

Step 5. *There exist positive constants* c *and* δ *such that, for every* $\tau_0 \geq 0$,

$$\varepsilon(\tau_0) := \int_{\tau_0}^{\infty} \int_0^{2\pi} \left(|\tilde{v}_\tau|^2 + e^{2\tau} |\mu(\tilde{u})|^2 \right) d\theta d\tau \leq c e^{-\delta \tau_0}.$$

By Step 2 and Lemma 11.3, we have

$$\begin{array}{rcl} \varepsilon(\tau) & = & \mathcal{A}(x_{e^\tau}, \eta_{e^\tau}) \\ & \leq & \delta^{-1} \displaystyle\int_0^{2\pi} \left(|\tilde{v}_\tau(\tau, \theta)|^2 + e^{2\tau} |\mu(\tilde{u}(\tau, \theta))|^2 \right) d\theta \\ & = & -\delta^{-1} \varepsilon'(\tau) \end{array}$$

for some constant $\delta > 0$ and every sufficiently large real number τ. Hence there exists a real number τ_0 such that

$$\tau \geq \tau_0 \qquad \Longrightarrow \qquad \varepsilon(\tau) \leq e^{-\delta(\tau - \tau_0)} \varepsilon(\tau_0).$$

Step 6. *There exist positive constants* c *and* δ *such that, for every* $\tau_0 \geq 0$,

$$\sup_{\theta \in \mathbb{R}} \left(|\tilde{v}_\tau(\tau_0, \theta)|^2 + e^{2\tau_0} |\mu(\tilde{u}(\tau_0, \theta))| \right) + \int_{\tau_0}^{\infty} \int_0^{2\pi} e^{4\tau} |\mu(\tilde{u}(\tau, \theta)|^2 \, d\theta d\tau \leq c e^{-\delta \tau_0}.$$

The L^∞ estimate for $\tilde{v}_\tau$ and $e^{2\tau}\mu(\tilde{u})$ follows from Step 5 and (90). The L^2-estimate for $\mu(\tilde{u})$ follows from Step 5 and Lemma 9.3.

Step 7. *There exists a $W^{1,2}$-function $x : S^1 \to \mu^{-1}(0)$ and an L^2-function $\eta : S^1 \to \mathfrak{g}$ such that*

$$\lim_{\tau\to\infty} \sup_{\theta\in\mathbb{R}} d(\tilde{u}(\tau,\theta), x(\theta)) = 0, \qquad \lim_{\tau\to\infty} \int_0^{2\pi} \left|\tilde{\Psi}(\tau,\theta) - \eta(\theta)\right|^2 d\theta = 0,$$

and $\dot{x} + X_\eta(x) = 0$. Moreover, $E(u,\Phi,\Psi) = \int u^*\omega$, and if $(H3)$ holds then $E(u,\Phi,\Psi) \in \mathbb{Z}\hbar$.

By Hölder's inequality and the radial gauge assumption, we have, for $\tau_1 > \tau_0$,

$$\begin{aligned} \left|\tilde{\Psi}(\tau_1,\theta) - \tilde{\Psi}(\tau_0,\theta)\right|^2 &\le \left(\int_{\tau_0}^{\tau_1} e^{2\tau} \left|\mu(\tilde{u}(\tau,\theta))\right| d\tau\right)^2 \\ &\le (\tau_1 - \tau_0) \int_{\tau_0}^{\tau_1} e^{4\tau} \left|\mu(\tilde{u}(\tau,\theta))\right|^2 d\tau. \end{aligned}$$

Hence the existence of the L^2-limit of $\tilde{\Psi}$ follows from Step 6. That $\tilde{u}(\tau,\theta)$ converges uniformly as τ tends to infinity follows from the exponential decay of $\tilde{v}_\tau = \partial_\tau \tilde{u}$ in Step 6. That the limit is a $W^{1,2}$-function and satisfies $\dot{x} + X_\eta(x) = 0$ follows from the fact that $\tilde{v}_\theta = \partial_\theta u + X_{\tilde{\Psi}}(\tilde{u})$ converges (exponentially) to zero as τ tends to infinity. That $E(u,\Phi,\Psi) = \int u^*\omega$ follows from the energy identity in the proof of Step 2 and the L^2-convergence of $\tilde{\Psi}$. That $E(u,\Phi,\Psi)$ is an integer multiple of $\hbar$ (when $(H3)$ holds) follows from the proof of Step 2. $\square$

Every map $u : \mathbb{C} \to M$ that satisfies (103) and (102) determines an equivariant homology class $B = [u] \in H_2(M_{\mathrm{G}};\mathbb{Z})$ as follows. Homotop u to a map $v : D \to M$ such that $v(e^{i\theta}) = g(e^{i\theta})x_0$. Now define a principal bundle $P \to S^2 \cong (\mathbb{Z}_2 \times D)/\sim$ by

$$P := (\mathbb{Z}_2 \times D \times \mathrm{G})/\sim, \qquad (0, e^{i\theta}, h) \sim (1, e^{i\theta}, g(e^{i\theta})h).$$

Then v determines a G-equivariant map $w : P \to M$ by

$$w(0,z,h) := h^{-1}x_0, \qquad w(1,z,h) := h^{-1}v(z).$$

The equivariant homology class of u is defined to be the equivariant homology class of w.

12 Proof of Theorem A

We begin by constructing a $\mathcal{G}$-equivariant smooth function from a suitable open subset of $\mathcal{B} \times P = C^\infty_{\mathrm{G}}(P, M) \times \mathcal{A}(P) \times P$ into a suitable finite dimensional approximation of EG. For positive constants δ and r denote

$$M^\delta := \{x \in M \,|\, |\mu(x)| \le \delta\},$$

$$\mathcal{B}^{\delta,r} := \left\{ (u, A) \in C^\infty_{\mathrm{G}}(P, M) \times \mathcal{A}(P) \,\middle|\, \exists z \in \Sigma \;\sup_{B_r(z)} |\mu \circ u| < \delta \right\}.$$

Throughout we assume that G is a Lie subgroup of $\mathrm{U}(k)$. Then, for every integer $n \ge k$, a finite dimensional approximation of the classifying space of G is the quotient

$$\mathrm{BG}^n := \mathrm{EG}^n/\mathrm{G}, \qquad \mathrm{EG}^n := \mathcal{F}(k, n) := \left\{\Theta \in \mathbb{C}^{n\times k} \,|\, \Theta^*\Theta = \mathbb{1}\right\}.$$

Let $\delta > 0$ be so small that G acts freely on M^δ and choose a smooth G-equivariant classifying map $\theta : M^\delta \to \mathrm{EG}^m$ for some integer $m \ge k$.

Proposition 12.1. *Let $\theta : M^\delta \to \mathrm{EG}^m$ be as above. Then, for every $r > 0$, there exist an integer $n \ge m$ and a map $\Theta : \mathcal{B}^{\delta,r} \times P \to \mathrm{EG}^n$ with the following properties.*

(i) *For $g \in \mathcal{G}$, $h \in \mathrm{G}$, and $(u, A, p) \in \mathcal{B}^{\delta,r} \times P$,*

$$\Theta(g^{-1}u, g^*A, pg(p)^{-1}) = \Theta(u, A, p) = h\Theta(u, A, ph). \tag{104}$$

(ii) *Θ is smooth with respect to the C^0 Banach manifold structure on (the completion of) $\mathcal{B}^{\delta,r} \times P$.*

(iii) *Let $\iota : \mathrm{EG}^m \to \mathrm{EG}^n$ be the obvious inclusion. Then*

$$|\mu(u(p))| < \delta \qquad \Longrightarrow \qquad \Theta(u, A, p) = \iota \circ \theta(u(p)).$$

Proof. Cover Σ by finitely many distinct balls $B_r(z_i)$, $i = 1, \dots, \ell$, and choose points $p_1, \dots, p_\ell \in P$ such that $\pi(p_i) = z_i$. Then, for every $(u, A) \in \mathcal{B}^{\delta,r}$, there exists an $i \in \{1, \dots, \ell\}$ such that $u(p_i) \in M^\delta$. Thus the open set $\mathcal{B}^{\delta,r} \times P$ is contained in the finite union of the following open sets $\mathcal{U}_{ij}$, $i, j = 1, \dots, \ell$.

Choose $\varepsilon > 0$ so small that θ extends to an equivariant function (still denoted by θ) from $M^{\delta+\varepsilon}$ to EG^m and define

$$\begin{aligned}
\mathcal{U}_0 &:= \{(u,A,p)\,|\,|\mu(u(p))| < \delta+\varepsilon\}\,,\\
\mathcal{U}_{ij} &:= \{(u,A,p)\,|\,\pi(p)\in B_r(z_i),\ |\mu(u(p_j))| < \delta+\varepsilon\}\,.
\end{aligned}$$

For every smooth path $\gamma : [0,1] \to \Sigma$ and any two points $p_0 \in \pi^{-1}(\gamma(0))$ and $p_1 \in \pi^{-1}(\gamma(1))$ the holonomy $\rho_A(p_1,\gamma,p_0) \in \mathrm{G}$ of the connection $A \in \mathcal{A}(P)$ is defined by $p_1\rho_A(p_1,\gamma,p_0) := \tilde{\gamma}(1)$, where $\tilde{\gamma} : [0,1] \to P$ is the unique horizontal lift of γ with $\tilde{\gamma}(0) = p_0$. It satisfies

$$\begin{aligned}
\rho_A(p_1g_1,\gamma,p_0g_0) &= {g_1}^{-1}\rho_A(p_1,\gamma,p_0)g_0,\\
\rho_{g^*A}(p_1,\gamma,p_0) &= g(p_1)^{-1}\rho_A(p_1,\gamma,p_0)g(p_0)
\end{aligned}$$

for $g_0, g_1 \in \mathrm{G}$ and $g \in \mathcal{G}$. Hence the map $\mathcal{B} \times \pi^{-1}(\gamma(0)) \to M : (u,A,p_0) \mapsto \rho_A(p_1,\gamma,p_0)^{-1}u(p_1)$ is $\mathcal{G}$-invariant and G-equivariant. Choose a finite sequence of smooth functions $\gamma_{ij} : [0,1] \times B_r(z_i) \to \Sigma$ such that

$$\gamma_{ij}(0,z) = z, \qquad \gamma_{ij}(1,z) = z_j.$$

Then the functions $\Theta_0 : \mathcal{U}_0 \to \mathrm{EG}^m$ and $\Theta_{ij} : \mathcal{U}_{ij} \to \mathrm{EG}^m$, defined by

$$\Theta_0(u,A,p) := \theta(u(p)), \qquad \Theta_{ij}(u,A,p) := \theta\Big(\rho_A\big(p_j,\gamma_{ij}(\cdot,\pi(p)),p\big)^{-1}u(p_j)\Big)$$

for $i,j = 1,\ldots,\ell$, satisfy (104). Now choose a G-invariant smooth function $\beta : M \to [0,1]$ such that $\beta(x) = 1$ for $x \in M^\delta$ and $\beta(x) = 0$ for $x \in M \setminus M^{\delta+\varepsilon/2}$. Define $\rho_j : \bigcup_{i=1}^{\ell}\mathcal{U}_{ij} \to [0,1]$ and $\rho_0 : \mathcal{U}_0 \to [0,1]$ by

$$\begin{aligned}
\rho_0(u,A,p) &:= \frac{\beta(u(p))}{\sqrt{\beta(u(p))^2 + (1-\beta(u(p)))^2}},\\
\rho_j(u,A,p) &:= \frac{\beta(u(p_j))(1-\beta(u(p)))}{\sqrt{(\beta(u(p))^2 + (1-\beta(u(p)))^2)\sum_{k=1}^{\ell}\beta(u(p_k))^2}}
\end{aligned}$$

for $j = 1,\ldots,\ell$. Then ρ_j is smooth with respect to the C^0 Banach manifold structure on (the completion of) $\mathcal{B} \times P$. Moreover,

$$\sum_{j=0}^{\ell}\rho_j(u,A,p)^2 = 1$$

and $\rho_0(u,A,p)=1$ whenever $|\mu(u(p))|\le\delta$. Now choose a finite sequence of smooth functions $\sigma_i:\Sigma\to[0,1]$, $i=1,\dots,\ell$, such that $\operatorname{supp}\sigma_i\subset B_r(z_i)$ and $\sum_{j=1}^{\ell}{\sigma_i}^2=1$. Then the function $\Theta:\mathcal{B}^{\delta,r}\to\mathrm{EG}^{(\ell^2+1)m}$, defined by

$$\Theta(u,A):=\left(\begin{array}{c}\rho_0(u,A,p)\Theta_0(u,A,p)\\ \vdots\\ \sigma_i(\pi(p))\rho_j(u,A,p)\Theta_{ij}(u,A,p)\\ \vdots\end{array}\right),$$

is the required classifying map. □

The integer $n=(\ell+1)m$ in the proof of Proposition 12.1 diverges to infinity as r tends to zero. In general, there is no $\mathcal{G}$-equivariant map from $\mathcal{B}^\delta:=\{(u,A)\in\mathcal{B}\,|\,\min|\mu\circ u|<\delta\}$ to any finite dimensional approximation of EG.

Lemma 12.2. *Assume* $(H1-3)$. *Let* $B\in H_2(M_{\mathrm{G}};\mathbb{Z})$, $(\Sigma,j_\Sigma,\mathrm{dvol}_\Sigma)$ *be a compact Riemann surface,* $\pi:P\to\Sigma$ *be the principal* G*-bundle determined by* B, *and* $\Sigma\to\mathcal{J}_{\mathrm{G}}(M,\omega)$ *be a smooth family of* G *invariant* ω*-compatible almost complex structures on* M *such that each* J_z *agrees with the almost complex structure of* $(H2)$ *outside of a sufficiently large compact subset of* M. *Then for every* $\delta>0$ *there exist positive constants* r *and* ε_0 *such that*

$$\widetilde{\mathcal{M}}^{\varepsilon}_{B,\Sigma}\subset\mathcal{B}^{\delta,r}$$

for $0<\varepsilon\le\varepsilon_0$.

Proof. Suppose the result is false. Then there exist a constant $\delta>0$ and sequences

$$r_i\to0,\qquad\varepsilon_i\to0,\qquad(u_i,A_i)\in\widetilde{\mathcal{M}}^{\varepsilon_i}_{B,\Sigma}$$

such that $(u_i,A_i)\notin\mathcal{B}^{\delta,r_i}$ for every i. This means that, for every $p\in P$, there exists a sequence $p_i\in P$ such that

$$\lim_{i\to\infty}p_i=p,\qquad|\mu(u_i(p_i))|\ge\delta.$$

This contradicts the bubbling argument in Step 5 of the proof of Theorem A below. □

Let $\varepsilon_0 > 0$ be as in Lemma 12.2. For $0 \le \varepsilon \le \varepsilon_0$ we consider the evaluation map

$$\mathrm{ev}^{\varepsilon}_{\mathrm{G}} : \mathcal{M}^{\varepsilon}_{B,\Sigma} \times \Sigma \to M \times_{\mathrm{G}} \mathrm{EG}^n,$$

given by

$$\mathrm{ev}^{\varepsilon}_{\mathrm{G}}([u, A, p]) := [u(p), \Theta(u, A, p)]$$

where $\delta > 0$ is chosen such that $\mathrm{G}_x = \{\mathbb{1}\}$ for every $x \in M^{\delta}$, $r > 0$ is as in Lemma 12.2, and $\Theta : \mathcal{B}^{\delta,r} \times P \to \mathrm{EG}^n$ is the map of Proposition 12.1. Recall that $\mathcal{M}^{0}_{B,\Sigma}$ and $\mathcal{M}^{\varepsilon}_{B,\Sigma}$ have the same dimension.

Proposition 12.3. *For every $c_0 > 0$ there exist positive constants c and ε_0 such that the following holds.*

(i) *For $0 < \varepsilon \le \varepsilon_0$ the map $\mathcal{T}^{\varepsilon} : \mathcal{M}^{0}_{B,\Sigma}(c_0) \to \mathcal{M}^{\varepsilon}_{B,\Sigma}$ is an orientation preserving embedding.*

(ii) *For $0 < \varepsilon \le \varepsilon_0$,*

$$d_{C^1}(\mathrm{ev}^{0}_{\mathrm{G}}, \mathrm{ev}^{\varepsilon}_{\mathrm{G}} \circ (\mathcal{T}^{\varepsilon} \times \mathrm{id})) \le c\varepsilon^{1-2/p},$$

where the C^1-distance is understood on the space of continuously differentiable maps from $\mathcal{M}^{0}_{B,\Sigma}(c_0) \times \Sigma$ to $M \times_{\mathrm{G}} \mathrm{EG}^n$.

Lemma 12.4. *Assume $(H1)$ and $(H4)$ and let $\bar{B} \in H_2(\bar{M}; \mathbb{Z})$ be a nontorsion homology class. For every $p > 2$ and every $c_0 > 0$ there exist positive constants ε_0 and c such that the following holds for every $\varepsilon \in (0, \varepsilon_0]$. Let $I \subset \mathbb{R}$ be an interval and*

$$I \to \widetilde{\mathcal{M}}^{0}_{B,\Sigma}(c_0) : r \mapsto (u_0(r), A_0(r))$$

be a smooth path that satisfies (79). Then every smooth vector field

$$r \mapsto \zeta(r) \in \mathrm{im}\, \left(\mathcal{D}^{\varepsilon}_{(u_0(r),A_0(r))}\right)^*$$

satisfies the inequality

$$\left\|\tilde{\nabla}_r \zeta\right\|_{1,p,\varepsilon} \le c\left(\varepsilon \left\|\tilde{\nabla}_r \mathcal{D}^{\varepsilon}\zeta\right\|_{0,p,\varepsilon} + \left\|\tilde{\nabla}_r \pi_{u_0} \mathcal{D}^{\varepsilon}\zeta\right\|_{L^p} + \varepsilon^{-1}\left\|\zeta\right\|_{1,p,\varepsilon}\right)$$

for $r \in I$, where $\mathcal{D}^{\varepsilon} := \mathcal{D}^{\varepsilon}_{(u_0(r),A_0(r))}$ and π_{u_0} is defined by (9).

Proof. Let $r \mapsto \zeta'(r) \in \mathcal{X}'_{u_0(r)}$ be the smooth path defined by $\zeta = \mathcal{D}^{\varepsilon *}\zeta'$. Then, by Lemmata 4.7, 4.8, and 8.4, we have

$$\begin{aligned}
\left\|\tilde{\nabla}_r\zeta\right\|_{1,p,\varepsilon} &\le \left\|\mathcal{D}^{\varepsilon *}\tilde{\nabla}_r\zeta'\right\|_{1,p,\varepsilon} + \left\|\tilde{\nabla}_r\mathcal{D}^{\varepsilon *}\zeta' - \mathcal{D}^{\varepsilon *}\tilde{\nabla}_r\zeta'\right\|_{1,p,\varepsilon} \\
&\le c_2\left(\varepsilon\left\|\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon *}\tilde{\nabla}_r\zeta'\right\|_{0,p,\varepsilon} + \left\|\pi_{u_0}\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon *}\tilde{\nabla}_r\zeta'\right\|_{L^p}\right) \\
&\quad + \left\|\tilde{\nabla}_r\mathcal{D}^{\varepsilon *}\zeta' - \mathcal{D}^{\varepsilon *}\tilde{\nabla}_r\zeta'\right\|_{1,p,\varepsilon} \\
&\le c_2\left(\varepsilon\left\|\mathcal{D}^{\varepsilon}\tilde{\nabla}_r\zeta\right\|_{0,p,\varepsilon} + \left\|\pi_{u_0}\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon *}\tilde{\nabla}_r\zeta'\right\|_{L^p}\right) \\
&\quad + c_3\left\|\tilde{\nabla}_r\mathcal{D}^{\varepsilon *}\zeta' - \mathcal{D}^{\varepsilon *}\tilde{\nabla}_r\zeta'\right\|_{1,p,\varepsilon} \\
&\le c_2\left(\varepsilon\left\|\tilde{\nabla}_r\mathcal{D}^{\varepsilon}\zeta\right\|_{0,p,\varepsilon} + \left\|\pi_{u_0}\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon *}\tilde{\nabla}_r\zeta'\right\|_{L^p}\right) \\
&\quad + c_2\varepsilon\left\|\mathcal{D}^{\varepsilon}\tilde{\nabla}_r\zeta - \tilde{\nabla}_r\mathcal{D}^{\varepsilon}\zeta\right\|_{0,p,\varepsilon} + c_4\varepsilon^{-1}\left\|\zeta'\right\|_{2,p,\varepsilon} \\
&\le c_2\left(\varepsilon\left\|\tilde{\nabla}_r\mathcal{D}^{\varepsilon}\zeta\right\|_{0,p,\varepsilon} + \left\|\tilde{\nabla}_r\pi_{u_0}\mathcal{D}^{\varepsilon}\zeta\right\|_{L^p}\right) \\
&\quad + c_2\left\|[\pi_{u_0}\mathcal{D}^{\varepsilon}\mathcal{D}^{\varepsilon *}, \tilde{\nabla}_r]\zeta'\right\|_{L^p} + c_5\varepsilon^{-1}\left\|\zeta\right\|_{1,p,\varepsilon} \\
&\le c_2\left(\varepsilon\left\|\tilde{\nabla}_r\mathcal{D}^{\varepsilon}\zeta\right\|_{0,p,\varepsilon} + \left\|\tilde{\nabla}_r\pi_{u_0}\mathcal{D}^{\varepsilon}\zeta\right\|_{L^p}\right) \\
&\quad + c_6\left(\left\|\mathcal{D}^{\varepsilon}\zeta\right\|_{0,p,\varepsilon} + \varepsilon^{-1}\left\|\pi_{u_0}\mathcal{D}^{\varepsilon}\zeta\right\|_{L^p} + \varepsilon^{-1}\left\|\zeta\right\|_{1,p,\varepsilon}\right).
\end{aligned}$$

The last inequality follows as in Step 7 in the proof of Lemma 4.8. Since

$$\left\|\mathcal{D}^{\varepsilon}\zeta\right\|_{0,p,\varepsilon} + \varepsilon^{-1}\left\|\pi_{u_0}\mathcal{D}^{\varepsilon}\zeta\right\|_{L^p} \le c_7\varepsilon^{-1}\left\|\zeta\right\|_{1,p,\varepsilon},$$

the lemma is proved. □

Proof of Proposition 12.3. Let $r \mapsto (u_0(r), A_0(r))$ be as in Lemma 12.4 and $r \mapsto \zeta_\varepsilon(r) = (\xi_\varepsilon(r), \alpha_\varepsilon(r)) \in \mathrm{im}\,(\mathcal{D}^{\varepsilon}_{(u_0(r),A_0(r))})^*$ be as in Theorem 6.1 so that

$$\widetilde{\mathcal{T}}^{\varepsilon}(u_0(r), A_0(r)) = (u_\varepsilon(r), A_\varepsilon(r)) := (\exp_{u_0(r)}(\xi_\varepsilon(r)), A_0(r) + \alpha_\varepsilon(r)).$$

Let $\mathcal{F}^{\varepsilon}_r : \mathcal{X}_{u_0(r)} \to \mathcal{X}'_{u_0(r)}$ be defined by (26). Then $\mathcal{F}^{\varepsilon}_r(\zeta_\varepsilon(r)) = 0$ and hence

$$\mathcal{D}^{\varepsilon}\zeta_\varepsilon = \begin{pmatrix} 0 \\ 0 \\ *F_{A_0(r)} \end{pmatrix} + \Big(\mathcal{F}^{\varepsilon}_r(\zeta_\varepsilon(r)) - \mathcal{F}^{\varepsilon}_r(0) - d\mathcal{F}^{\varepsilon}_r(0)\zeta_\varepsilon(r)\Big).$$

Differentiating this identity with respect to r we find

$$\left\|\tilde{\nabla}_r\mathcal{D}^\varepsilon\zeta_\varepsilon\right\|_{0,p,\varepsilon} \le c\left(\varepsilon + \varepsilon^{-1-2/p}\left\|\zeta_\varepsilon(r)\right\|_{1,p,\varepsilon}\left(\left\|\zeta_\varepsilon(r)\right\|_{1,p,\varepsilon} + \left\|\tilde{\nabla}_r\zeta_\varepsilon(r)\right\|_{1,p,\varepsilon}\right)\right).$$

(See Proposition 5.2.) Hence, by Lemma 12.4,

$$\left\|\tilde{\nabla}_r\zeta_\varepsilon\right\|_{1,p,\varepsilon} \le c'\varepsilon. \tag{105}$$

Since

$$(\partial_r u_\varepsilon, \partial_r A_\varepsilon) = (E_1(u_0,\xi_\varepsilon)\partial_r u_0 + E_2(u_0,\xi_\varepsilon)\nabla_r\xi_\varepsilon, \partial_r A_0 + \partial_r\alpha_\varepsilon),$$

this shows that $\mathcal{T}^\varepsilon : \mathcal{M}^0_{B,\Sigma}(c_0) \to \mathcal{M}^\varepsilon_{B,\Sigma}$ is an orientation preserving embedding. Indeed, it follows that the restriction of $\mathcal{T}^\varepsilon$ to every ball of radius δ is an embedding for δ and ε sufficiently small and hence, by Theorem 6.1, $\mathcal{T}^\varepsilon$ is an embedding for ε sufficiently small. For $\varepsilon \ge 0$ denote

$$\mathcal{P}^\varepsilon_{B,\Sigma}(c_0) := \widetilde{\mathcal{M}}^\varepsilon_{B,\Sigma}(c_0) \times_{\mathcal{G}} P$$

and consider the map $\mathrm{ev}^\varepsilon : \mathcal{P}^\varepsilon_{B,\Sigma}(c_0) \to M$ defined by

$$\mathrm{ev}^\varepsilon([u_\varepsilon, A_\varepsilon, p]) := u_\varepsilon(p).$$

Then it follows from (105) and the inequality $\|\zeta_\varepsilon\|_{2,p,\varepsilon} \le c'\varepsilon^2$ of Theorem 6.1 that

$$d_{C^1}(\mathrm{ev}^0, \mathrm{ev}^\varepsilon \circ (\mathcal{T}^\varepsilon \times \mathrm{id})) \le c''\varepsilon^{1-2/p}.$$

For ε sufficiently small we have $\mathrm{ev}^\varepsilon_{\mathrm{G}} = \theta^\delta \circ \mathrm{ev}^\varepsilon$, where $\theta^\delta : M^\delta \to M \times_{\mathrm{G}} \mathrm{EG}^n$ is given by $\theta^\delta(x) := [x, \theta(x)]$. This proves the proposition. $\square$

Proof of Theorem A. The result is obvious when $\bar{B} = 0$. Moreover, both moduli spaces are empty when $\bar{B}$ is a nonzero torsion class. Hence assume that $\bar{B} \in H_2(\bar{M};\mathbb{Z})$ is a nontorsion homology class, denote by $B \in H_2(M_{\mathrm{G}};\mathbb{Z})$ the corresponding equivariant homology class, fix a compact Riemann surface $(\Sigma, j_\Sigma, \mathrm{dvol}_\Sigma)$, and let $\pi : P \to \Sigma$ be a principal G-bundle whose characteristic class $b \in H_2(\mathrm{BG};\mathbb{Z})$ is the pushforward of B. In the course of the proof it will be necessary to also consider other bundles $\pi' : P' \to \Sigma$ with corresponding equivariant homology clsses $B' \in H_2(M_{\mathrm{G}};\mathbb{Z})$. By $(H2)$, there exists a constant $c > 0$ such that $u(P) \subset M^c := \{x \in M \,|\, |\mu(x)| \le c\}$ for every solution (u, A) of (7) over any Riemann surface. Note that c can be chosen to be a regular value of the function $M \to \mathbb{R} : x \mapsto |\mu(x)|$.

Let $\delta > 0$ such that $\mathrm{G}_x = \{\mathbb{1}\}$ for every $x \in M^\delta$ and let r and ε_0 be as in Lemma 12.2. Fix k points $p_1, \dots, p_k \in P$ such that the points $z_i := \pi(p_i) \in \Sigma$ are pairwise distinct. Choose an integer n, a G-equivariant smooth map $\theta : M^\delta \to \mathrm{EG}^n$, and k smooth classifying maps $\Theta_i : \mathcal{B}^{\delta,r} \to \mathrm{EG}^n$, defined by $\Theta_i(u, A) := \Theta(u, A, p_i)$, where Θ is as in Proposition 12.1. Then

$$\Theta_i(g^{-1}u, g^*A) = g(p_i)^{-1}\Theta_i(u, A)$$

and

$$|\mu(u(p_i))| \leq \delta \qquad \Longrightarrow \qquad \Theta_i(u, A) = \theta(u(p_i))$$

for $i = 1, \dots, k$. For $0 \leq \varepsilon \leq \varepsilon_0$ consider the evaluation maps

$$\mathrm{ev}^\varepsilon_{B,i} : \mathcal{M}^\varepsilon_{B,\Sigma} \to M^c_{\mathrm{G}} := M^c \times_{\mathrm{G}} \mathrm{EG}^n$$

given by $\mathrm{ev}^\varepsilon_{B,i}([u, A]) := [u(p_i), \Theta_i(u, A)]$. Let $\mathrm{ev}^\varepsilon_B : \mathcal{M}^\varepsilon_{B,\Sigma} \to (M^c_{\mathrm{G}})^k$ denote the product map defined by

$$\mathrm{ev}^\varepsilon_B([u, A]) := \left(\mathrm{ev}^\varepsilon_{B,1}([u, A]), \dots, \mathrm{ev}^\varepsilon_{B,k}([u, A])\right).$$

For any subset $I = \{i_1, \dots, i_j\} \subset \{1, \dots, k\}$ such that $i_1 < \dots < i_j$ and any class $B' \in H_2(M_{\mathrm{G}}; \mathbb{Z})$ that descends to $H_2(\bar{M}; \mathbb{Z})$ we consider the evaluation map $\mathrm{ev}^0_{B',I} : \mathcal{M}^0_{B',\Sigma} \to (M^c_{\mathrm{G}})^{|I|}$ given by

$$\mathrm{ev}^0_{B',I}([u, A]) := \left([u(p_{i_1}), \theta(u(p_{i_1}))], \dots, [u(p_{i_j}), \theta(u(p_{i_j}))]\right).$$

Now fix equivariant cohomology classes $\alpha_1, \dots, \alpha_k \in H^*(M_{\mathrm{G}}; \mathbb{Z})$ such that

$$m_i := \deg(\alpha_i) < 2N, \qquad \sum_{i=1}^k \deg(\alpha_i) = \dim\, \mathcal{M}^0_{B,\Sigma}.$$

There is a natural embedding $M^c_{\mathrm{G}} \to M_{\mathrm{G}}$ and we denote by $\alpha^c_i \in H^{m_i}(M^c_{\mathrm{G}}; \mathbb{Z})$ the pullback of α_i under this embedding. Note that M^c_{G} is a compact manifold with boundary. Replacing α_i by some integer multiple of α_i, if necessary, we may assume without loss of generality that, for every i, there exists a compact oriented manifold with boundary Y_i of dimension

$$\dim\, Y_i = \dim\, M^c_{\mathrm{G}} - m_i$$

and a smooth map

$$f_i : (Y_i, \partial Y_i) \to (M^c_{\mathrm{G}}, \partial M^c_{\mathrm{G}})$$

such that the homology class in $H_*(M^c_{\mathrm{G}}, \partial M^c_{\mathrm{G}})$ represented by f_i is Poincaré dual to α^c_i. For $I = \{i_1, \dots, i_j\} \subset \{1, \dots, k\}$ such that $i_1 < \cdots < i_j$ we denote the corresponding product map by

$$Y_I := Y_{i_1} \times \cdots \times Y_{i_j}, \qquad f_I := f_{i_1} \times \cdots \times f_{i_j} : Y_I \to (M^c_{\mathrm{G}})^{|I|}.$$

For $I = \{1, \dots, k\}$ we abbreviate $Y := Y_{\{1,\dots,k\}}$ and $f := f_{\{1,\dots,k\}}$. The functions $f_1, \dots, f_k$ can be chosen such that the following holds.

(H5) *f_i is transverse to $\mu^{-1}(0) \times_{\mathrm{G}} \mathrm{EG}^n$ for every i and f_I is transverse to $\mathrm{ev}^0_{B',I}$ for every subset $I \subset \{1, \dots, \ell\}$ and every equivariant homology class $B' \in H_2(M_{\mathrm{G}}; \mathbb{Z})$.*

Now the notation has been set up and we shall prove Theorem A in five steps. For $0 \le \varepsilon \le \varepsilon_0$ and $B \in H_2(M_{\mathrm{G}}; \mathbb{Z})$ consider the set

$$\mathcal{M}^\varepsilon_{B,\Sigma;f} := \left\{([u, A], y_1, \dots, y_k) \in \mathcal{M}^\varepsilon_{B,\Sigma} \times Y \,|\, \mathrm{ev}^\varepsilon_{B,i}([u, A]) = f_i(y_i)\right\}.$$

Step 1. *The map $\mathrm{ev}^0_B : \mathcal{M}^0_{B,\Sigma} \to (\mu^{-1}(0) \times_{\mathrm{G}} \mathrm{EG}^n)^k$ is a pseudo-cycle.*

The map ev^0_B is the composition $\mathrm{ev}^0_B = \iota^k \circ \overline{\mathrm{ev}}_B$, where the evaluation map $\overline{\mathrm{ev}}_B : \mathcal{M}^0_{B,\Sigma} \to \bar{M}^k$ is given by $\overline{\mathrm{ev}}_B([u, A]) := ([u(p_1)], \dots, [u(p_k)])$ and the embedding $\iota : \bar{M} \to \mu^{-1}(0) \times_{\mathrm{G}} \mathrm{EG}^n$ is given by $\iota([x]) := [x, \theta(x)]$. That $\overline{\mathrm{ev}}_B$ is a pseudo-cycle was proven in [MS1]. Hence ev^0_B is a pseudo-cycle. (see [MS1] for the definitions).

Step 2. *$\mathcal{M}^0_{B,\Sigma;f}$ is a finite set and the number of elements of $\mathcal{M}^0_{B,\Sigma;f}$, counted with appropriate signs, is the Gromov–Witten invariant:*

$$\mathrm{G}W_{\bar{B},\Sigma}(\bar{\alpha}_1, \dots, \bar{\alpha}_k) = \mathrm{ev}^0_B \cdot f = \sum_{([u_0,A_0],y) \in \mathcal{M}^0_{B,\Sigma;f}} \nu^0([u_0, A_0], y).$$

Here the function $\nu^0 : \mathcal{M}^0_{B,\Sigma;f} \to \{\pm 1\}$ denotes the intersection index of the maps ev^0_B and f.

Consider the functions $\varphi_i : X_i \to \mu^{-1}(0) \times_{\mathrm{G}} \mathrm{EG}^n$ given by

$$X_i := f_i^{-1}(\mu^{-1}(0) \times_{\mathrm{G}} \mathrm{EG}^n) \subset Y_i, \qquad \varphi_i := f_i|_{X_i}.$$

Since f_i is transverse to $\mu^{-1}(0) \times_{\mathrm{G}} \mathrm{EG}^n$, X_i is a smooth submanifold of Y_i and φ_i is dual to the cohomology class $\alpha^0_i \in H^*(\mu^{-1}(0) \times_{\mathrm{G}} \mathrm{EG}^n; \mathbb{Z})$ obtained from α^c_i by pullback under the obvious inclusion $\mu^{-1}(0) \times_{\mathrm{G}} \mathrm{EG}^n \to M^c_{\mathrm{G}}$.

The class α_i^0 agrees with the image of the class $\bar{\alpha}_i \in H^*(\bar{M};\mathbb{Z})$ under the homomorphism $H^*(\bar{M};\mathbb{Z}) \cong H^*(\mu^{-1}(0)\times_{\mathrm{G}}\mathrm{EG};\mathbb{Z}) \to H^*(\mu^{-1}(0)\times_{\mathrm{G}}\mathrm{EG}^n;\mathbb{Z})$:

$$\begin{tikzcd}
\alpha_i^c \in & \mathrm{H}^*(M^c \times_{\mathrm{G}} \mathrm{EG}^n;\mathbb{Z}) \arrow[d] & \mathrm{H}^*(M_{\mathrm{G}};\mathbb{Z}) \arrow[l] \arrow[d, "\kappa"] & \ni \alpha_i \\
\alpha_i^0 \in & \mathrm{H}^*(\mu^{-1}(0) \times_{\mathrm{G}} \mathrm{EG}^n;\mathbb{Z}) & \mathrm{H}^*(\bar{M};\mathbb{Z}) \arrow[l] & \ni \bar{\alpha}_i
\end{tikzcd}$$

Hence another representative of the class α_i^0 can be obtained as follows. Let $\bar{\psi}_i : Z_i \to \bar{M}$ be a smooth function, defined on a compact manifold $\bar{Z}_i$ that is dual to $\bar{\alpha}_i$ (replace $\bar{\alpha}_i$ by an integer multiple of $\bar{\alpha}_i$, if necessary). Lift $\bar{\psi}_i$ to a G-equivariant map $\tilde{\psi}_i : Q_i \to \mu^{-1}(0)$, defined on the total space of a principal G-bundle $Q_i \to Z_i$, and consider the induced map

$$\psi_i : Q_i \times_{\mathrm{G}} \mathrm{EG}^n \to \mu^{-1}(0) \times_{\mathrm{G}} \mathrm{EG}^n.$$

It is homologous to φ_i. Let $\varphi := \varphi_1 \times \cdots \times \varphi_k$ and $\psi := \psi_1 \times \cdots \times \psi_k$. Then

$$\mathrm{ev}_B^0 \cdot f = \mathrm{ev}_B^0 \cdot \varphi = \mathrm{ev}_B^0 \cdot \psi = \overline{\mathrm{ev}}_B \cdot \bar{\psi} = \mathrm{GW}_{\bar{B},\Sigma}(\bar{\alpha}_1, \ldots, \bar{\alpha}_k).$$

The first equality follows from the definition of φ, the second from the fact that ev_B^0 is a pseudo-cycle (Step 1) and φ is homologous to ψ, and the third equality follows from the definition of the Gromov–Witten invariants (see [MS1] for example).

Step 3. *The invariant $\Phi_{B,\Sigma}$ can be expressed as the intersection number*

$$\Phi_{B,\Sigma}(\alpha_1 \smile \cdots \smile \alpha_k) = \mathrm{ev}_B^\varepsilon \cdot f$$

for $\varepsilon > 0$ sufficiently small.

The map $f : Y \to (M_{\mathrm{G}}^c)^k$ is dual to the class $\pi_1^*\alpha_1 \smile \cdots \smile \pi_k^*\alpha_k$, where $\pi_i : (M_{\mathrm{G}}^c)^k \to M_{\mathrm{G}}^c$ denotes the projection onto the ith factor. Moreover, $\mathrm{ev}_{B,i}^\varepsilon = \pi_i \circ \mathrm{ev}_B^\varepsilon$. Hence

$$\begin{aligned}
\mathrm{ev}_B^\varepsilon \cdot f &= \int_{\mathcal{M}_{B,\Sigma}^\varepsilon} (\mathrm{ev}_B^\varepsilon)^* \left(\pi_1^*\alpha_1^c \smile \cdots \smile \pi_k^*\alpha_k^c\right) \\
&= \int_{\mathcal{M}_{B,\Sigma}^\varepsilon} (\mathrm{ev}_{B,1}^\varepsilon)^*\alpha_1^c \smile \cdots \smile (\mathrm{ev}_{B,k}^\varepsilon)^*\alpha_k^c \\
&= \int_{\mathcal{M}_{B,\Sigma}^\varepsilon} (\iota^c \circ \mathrm{ev}_{B,1}^\varepsilon)^*\alpha_1 \smile \cdots \smile (\iota^c \circ \mathrm{ev}_{B,k}^\varepsilon)^*\alpha_k \\
&= \int_{\mathcal{M}_{B,\Sigma}^\varepsilon} \mathrm{ev}_{\mathrm{G}}^* \left(\alpha_1 \smile \cdots \smile \alpha_k\right).
\end{aligned}$$

Here $\iota^c : M_{\mathrm{G}}^c \to M_{\mathrm{G}}$ denotes the obvious inclusion. The last equality follows from the fact that $\iota^c \circ \mathrm{ev}_{B,i}^{\varepsilon} : \mathcal{M}_{B,\Sigma}^{\varepsilon} \to M_{\mathrm{G}}$ is homotopic to the evaluation map ev_{G} in the definition of $\Phi_{B,\Sigma}$.

Step 4. *For $\varepsilon > 0$ sufficiently small there is an injective map*

$$\mathcal{T}_{B,\Sigma;f}^{\varepsilon} : \mathcal{M}_{B,\Sigma;f}^{0} \to \mathcal{M}_{B,\Sigma;f}^{\varepsilon}$$

such that

$$\mathcal{T}_{B,\Sigma;f}^{\varepsilon}([u_0, A_0], y_{0,1}, \ldots, y_{0,k}) = ([u_\varepsilon, A_\varepsilon], y_{\varepsilon,1}, \ldots, y_{\varepsilon,k})$$

satisfies

$$(u_\varepsilon, A_\varepsilon) = (\exp_{u_0}(\xi_\varepsilon), A_0 + \alpha_\varepsilon), \qquad \|(\xi_\varepsilon, \alpha_\varepsilon)\|_{2,p,\varepsilon} \le c\varepsilon^{2-2/p},$$

$$\nu^{\varepsilon}([u_\varepsilon, A_\varepsilon], y_{\varepsilon,1}, \ldots, y_{\varepsilon,k}) = \nu^{0}([u_0, A_0], y_{0,1}, \ldots, y_{0,k}).$$

Here $\nu^{\varepsilon} : \mathcal{M}_{B,\Sigma;f}^{\varepsilon} \to \{\pm 1\}$ denotes the intersection index of the maps $\mathrm{ev}_B^{\varepsilon}$ and f (in the transverse case).

Choose $c_0 > 0$ such that $\mathcal{M}_{B,\Sigma;f}^{0} \subset \mathcal{M}_{B,\Sigma}^{0}(c_0)$ and consider the map

$$(\mathrm{ev}_B^{\varepsilon} \circ \mathcal{T}^{\varepsilon}) \times f : \mathcal{M}_{B,\Sigma}^{0}(c_0) \times Y \to M_{\mathrm{G}}^c \times M_{\mathrm{G}}^c.$$

By Proposition 12.3 (ii), this map converges to $\mathrm{ev}_B^0 \times f$ in the C^1-topology as ε tends to zero. By $(H5)$ the map $\mathrm{ev}_B^0 \times f$ is transverse to the diagonal $\Delta \subset M_{\mathrm{G}}^c \times M_{\mathrm{G}}^c$. Hence $(\mathrm{ev}_B^{\varepsilon} \circ \mathcal{T}^{\varepsilon}) \times f$ is transverse to Δ for ε sufficiently small. Moreover, by Theorem 6.1, the image of $\mathcal{M}_{B,\Sigma;f}^{0}$ under $(\mathrm{ev}_B^{\varepsilon} \circ \mathcal{T}^{\varepsilon}) \times f$ is $\varepsilon^{2-2/p}$-close to Δ. Hence, by the implicit function theorem, there is, for ε sufficiently small, a unique injective map

$$\mathcal{M}_{B,\Sigma;f}^{0} \to ((\mathrm{ev}_B^{\varepsilon} \circ \mathcal{T}^{\varepsilon}) \times f)^{-1}(\Delta) \subset \mathcal{M}_{B,\Sigma}^{0}(c_0) \times Y$$

such that the distance between each point and its image is bounded above by a constant times $\varepsilon^{2-2/p}$. Composing this map with

$$\mathcal{T}^{\varepsilon} \times \mathrm{id} : \mathcal{M}_{B,\Sigma}^{0} \times Y \to \mathcal{M}_{B,\Sigma}^{\varepsilon} \times Y$$

we obtain the required map $\mathcal{T}_{B,\Sigma;f}^{\varepsilon}$. By Proposition 12.3 (i), the map $\mathcal{T}_{B,\Sigma;f}^{\varepsilon}$ identifies the two intersection indices.

Step 5. *Assume $\Sigma = S^2$. Then there exists a constant $\varepsilon_0 > 0$ such that the map $\mathcal{T}^{\varepsilon}_{B,\Sigma;f} : \mathcal{M}^0_{B,\Sigma;f} \to \mathcal{M}^{\varepsilon}_{B,\Sigma;f}$ of Step 4 is surjective for $0 < \varepsilon \le \varepsilon_0$.*

Suppose, by contradiction, that there exist sequences $\varepsilon_\nu \to 0$ and

$$([u_\nu, A_\nu], y_{1\nu}, \cdots, y_{k\nu}) \in \mathcal{M}^{\varepsilon_\nu}_{B,S^2;f}$$

such that

$$([u_\nu, A_\nu], y_{1\nu}, \cdots, y_{k\nu}) \notin \operatorname{im} \mathcal{T}^{\varepsilon_\nu}_{B,S^2;f}.$$

Consider the sequence

$$C_\nu := \sup_P \left(|d_{A_\nu} u_\nu| + \varepsilon_\nu^{-1} |\mu(u_\nu)| \right).$$

We prove that C_ν diverges to ∞. Assume otherwise that C_ν is bounded. Then, by Theorem D, there exists a constant $c_1 > 0$ such that $[u_\nu, A_\nu]$ belongs to the image of the map $\mathcal{T}^{\varepsilon_\nu}_{B,S^2} : \mathcal{M}^0_{B,S^2}(c_1) \to \mathcal{M}^{\varepsilon_\nu}_{B,S^2}$ for ν sufficiently large. Write

$$(u_\nu, A_\nu) = \widetilde{\mathcal{T}}^{\varepsilon_\nu}(u_{0\nu}, A_{0\nu}), \qquad (u_{0\nu}, A_{0\nu}) \in \widetilde{\mathcal{M}}^0_{B,S^2}(c_1).$$

Since $\mathcal{M}^0_{B,S^2}(c_1)$ is compact we may assume that the limit

$$(u_0, A_0) = \lim_{\nu \to \infty} (u_{0\nu}, A_{0\nu}) \in \widetilde{\mathcal{M}}^0_{B,S^2}(c_1)$$

exists. Moreover, since Y is compact, we may assume, by passing to a further subsequence if necessary, that the limit

$$(y_1, \ldots, y_k) = \lim_{\nu \to \infty} (y_{1\nu}, \ldots, y_{k\nu})$$

exists. Since $\mathrm{ev}^{\varepsilon_\nu}_B \circ \mathcal{T}^{\varepsilon_\nu}$ converges to ev^0_B in the C^1-topology, and

$$\mathrm{ev}^{\varepsilon_\nu}_B \circ \mathcal{T}^{\varepsilon_\nu}([u_{0\nu}, A_{0\nu}]) = f(y_{1\nu}, \cdots, y_{k\nu})$$

we deduce that $([u_0, A_0], y_1, \ldots, y_k) \in \mathcal{M}^0_{B,S^2;f}$ and, for ν sufficiently large,

$$(\mathcal{T}^{\varepsilon_\nu}([u_{0\nu}, A_{0\nu}]), y_{1\nu}, \cdots, y_{k\nu}) = \mathcal{T}^{\varepsilon_\nu}_{B,S^2;f}([u_0, A_0], y_1, \ldots, y_k).$$

The last assertion follows from the uniqueness part of the implicit function theorem used in the definition of the maps $\mathcal{T}^{\varepsilon_\nu}_{B,S^2;f}$. This contradicts our assumption. Thus we have proved that C_ν diverges to ∞ as claimed.

Now choose a sequence $p_\nu \in P$ such that

$$c_\nu := |d_{A_\nu} u_\nu(p_\nu)| + \varepsilon_\nu^{-1} |\mu(u_\nu(p_\nu))| \to \infty.$$

Passing to a subsequence, if necessary, we may assume that p_ν converges. Denote

$$\mathrm{w} := \lim_{\nu\to\infty} \pi(p_\nu).$$

Moreover, by applying Hofer's trick (see [MS1, Lemma 4.5.3] for example) we may assume that

$$\sup_{B_{r_\nu}(\pi(p_\nu))} \big(|d_{A_\nu} u_\nu| + \varepsilon_\nu^{-1} |\mu(u_\nu)| \big) \le 2c_\nu, \qquad r_\nu c_\nu \to \infty.$$

We distinguish three cases.

Case 1: $c_\nu \varepsilon_\nu \to \infty$.

Case 2: There exists a $\delta > 0$ such that $\delta \le c_\nu \varepsilon_\nu \le \delta^{-1}$ for all ν.

Case 3: $c_\nu \varepsilon_\nu \to 0$.

We shall prove that in Case 1 a nonconstant J-holomorphic sphere in M bubbles off at the point w, in Case 2 a nontrivial solution of the vortex equations (101) bubbles off, and in Case 3 a nonconstant $\bar{J}$-holomorphic sphere in $\bar{M}$ bubbles off. To see this, we choose a local holomorphic coordinate chart $s + it$ on Σ that maps w to zero, identifies a neighbourhood of w with the ball B_{2r}, and identifies the volume form dvol_Σ with the form $\lambda^2 ds \wedge dt$, where $\lambda(0) = 1$. Moreover, we choose a local frame of the bundle P along this coordinate chart. We use the notation of Remark 4.1. Then the sequences $u_\nu : B_{2r} \to M$ and $\Phi_\nu, \Psi_\nu : B_{2r} \to \mathfrak{g}$ satisfy

$$v_{\nu s} + J v_{\nu t} = 0, \qquad \lambda^{-2} \kappa_\nu + \varepsilon_\nu^{-2} \mu(u_\nu) = 0,$$

$$v_{\nu s} := \partial_s u_\nu + X_{\Phi_\nu}(u_\nu), \qquad v_{\nu t} := \partial_t u_\nu + X_{\Psi_\nu}(u_\nu),$$

$$\kappa_\nu := \partial_s \Psi_\nu - \partial_t \Phi_\nu + [\Phi_\nu, \Psi_\nu].$$

Moreover, there is a sequence $\mathrm{w}_\nu := (s_\nu, t_\nu) \to 0$ such that

$$c_\nu = \lambda(\mathrm{w}_\nu)^{-1} |v_{\nu s}(\mathrm{w}_\nu)| + \varepsilon_\nu^{-1} |\mu(u(\mathrm{w}_\nu))| \ge \frac{1}{2} \sup_{B_{r_\nu}(\mathrm{w}_\nu)} \big(\lambda^{-1} |v_{\nu s}| + \varepsilon_\nu^{-1} |\mu(u_\nu)| \big).$$

Let us define $\tilde{u}_\nu : B_{r_\nu c_\nu} \to M$ and $\tilde{\Phi}_\nu, \tilde{\Psi}_\nu : B_{r_\nu c_\nu} \to \mathfrak{g}$ by

$$\tilde{u}_\nu(z) := u(\mathrm{w}_\nu + c_\nu^{-1} z),$$

$$\tilde{\Phi}_\nu(z) := c_\nu^{-1} \Phi_\nu(\mathrm{w}_\nu + c_\nu^{-1} z), \qquad \tilde{\Psi}_\nu(z) := c_\nu^{-1} \Psi_\nu(\mathrm{w}_\nu + c_\nu^{-1} z),$$

and $\tilde{\lambda}_\nu : B_{r_\nu c_\nu} \to (0, \infty)$ and $\tilde{J}_\nu : B_{r_\nu c_\nu} \to \mathcal{J}_{\mathrm{G}}(M, \omega)$ by

$$\tilde{\lambda}_\nu(z) := \lambda(\mathrm{w}_\nu + c_\nu^{-1} z), \qquad \tilde{J}_\nu(z) = J_{\mathrm{w}_\nu + c_\nu^{-1} z}.$$

Then $\tilde{\lambda}_\nu$ converges to 1 in the C^∞-topology and $\tilde{J}_\nu$ converges to J_0 in the C^∞-topology. Moreover,

$$\tilde{v}_{\nu s} + \tilde{J}_\nu \tilde{v}_{\nu t} = 0, \qquad \tilde{\lambda}_\nu^{-2} \tilde{\kappa}_\nu + (c_\nu \varepsilon_\nu)^{-2} \mu(\tilde{u}_\nu) = 0,$$

$$\sup_{B_{r_\nu c_\nu}} \left(\frac{1}{\tilde{\lambda}_\nu} |\tilde{v}_{\nu s}| + \frac{1}{c_\nu \varepsilon_\nu} |\mu(\tilde{u}_\nu)| \right) \le 2 \left(\frac{1}{\tilde{\lambda}_\nu(0)} |\tilde{v}_{\nu s}(0)| + \frac{1}{c_\nu \varepsilon_\nu} |\mu(\tilde{u}_\nu(0))| \right) = 2.$$

Case 1: Suppose that $c_\nu \varepsilon_i$ diverges to infinity. Then, by hypothesis (H3), the curvature $\tilde{\kappa}_\nu$ converges uniformly to zero. Hence, by Uhlenbeck's weak compactness theorem [U, W], we may assume that $\tilde{\Phi}_\nu$ and $\tilde{\Psi}_\nu$ converge in the sup-norm and weakly in $W^{1,p}$. This implies that the sequence $\tilde{u}_i$ is bounded in $W^{1,p}$. Hence, by the usual elliptic bootstrapping argument for pseudoholomorphic curves, it is bounded in $W^{2,p}$ (the lower order terms in the equation have the form $X_{\tilde{\Phi}_i}(\tilde{u}_i)$ and hence are bounded in $W^{1,p}$). Hence there exists a subsequence, still denoted by $\tilde{u}_i$, that converges strongly in $W^{1,p}$ to a J_0-holomorphic curve $\tilde{u} : \mathbb{C} \to M$ with finite energy. Since the sequence $\mu(\tilde{u}_i(0))$ is bounded it follows that $|\partial_s \tilde{u}(0)| = \lim_{\nu \to \infty} |\tilde{v}_{\nu s}(0)| = 1$, and hence $\tilde{u}$ extends to a nonconstant holomorphic sphere in M. This contradicts $(H2)$.

Case 2: Suppose that the sequence $c_\nu \varepsilon_\nu$ is bounded and does not converge to zero. Let us assume, without loss of generality, that $\lim_{\nu \to \infty} c_\nu \varepsilon_\nu = 1$. Then we can use the compactness result of [CGMS] to deduce that, after a suitable gauge transformation and after passing to a further subsequence, the triple $(\tilde{u}_\nu, \tilde{\Phi}_\nu, \tilde{\Psi}_\nu)$ converges to a solution $(\tilde{u}, \tilde{\Phi}, \tilde{\Psi})$ of the vortex equations (101) with finite energy. Moreover,

$$|\partial_s \tilde{u}(0) + X_{\tilde{\Phi}(0)}(\tilde{u}(0))| + |\mu(\tilde{u}(0))| = 1$$

and hence the energy is nonzero. Hence, by Proposition 11.1, we have

$$E(\tilde{u}, \tilde{\Phi}, \tilde{\Psi}) \ge \hbar.$$

Case 3: Suppose that $\lim_{\nu\to\infty} c_\nu\varepsilon_\nu = 0$. Then, by Lemma 9.1,

$$\sup_\nu (c_\nu\varepsilon_\nu)^{-3/2}\|\mu(\tilde{u}_\nu)\|_{L^\infty(K)} + \sup_\nu (c_\nu\varepsilon_\nu)^{-2}\|\mu(\tilde{u}_\nu)\|_{L^2(K)} < \infty$$

for every compact set $K \subset \mathbb{C}$. It follows that the sequence κ_i is uniformly bounded in L^2. Hence, by Uhlenbeck's weak compactness theorem, we may assume that $\tilde{\Phi}_\nu$ and $\tilde{\Psi}_\nu$ converge weakly in $W^{1,2}$ and strongly in L^p, on every compact subset of $\mathbb{C}$. Here p is any fixed real number, say $p > 4$. Hence, the sequence $\tilde{u}_\nu$ is bounded in $W^{1,p}$. Now it follows again from the elliptic bootstrapping analysis for pseudoholomorphic curves that $\tilde{u}_\nu$ is bounded in $W^{2,2}$ and hence has a subsequence that converges strongly in $W^{1,p}$ on every compact subset of $\mathbb{C}$. The limit $(\tilde{u}, \tilde{\Phi}, \tilde{\Psi})$ is a finite energy solution of (8) on $\mathbb{C}$. This solution represents a $\bar{J}$-holomorphic sphere in the quotient $\bar{M}$. Moreover, since $(c_\nu\varepsilon_\nu)^{-1}|\mu(\tilde{u}_\nu(0))| \to 0$ it follows that

$$|\partial_s\tilde{u}(0) + X_{\tilde{\Phi}(0)}\tilde{u}(0)| = 1$$

and hence the resulting holomorphic sphere in $\bar{M}$ is nonconstant. Hence

$$E(\tilde{u}, \tilde{\Phi}, \tilde{\Psi}) \geq \hbar.$$

Thus we have proved in all three cases that

$$\lim_{\nu\to\infty} E_{B_r(\mathrm{w})}(u_\nu, A_\nu) \geq \hbar$$

for every $r > 0$.

This shows that, after passing to a suitable subsequence, bubbling can only take place at finitely many points $\mathrm{w}_1, \dots, \mathrm{w}_\ell \in \Sigma$. On every compact subset of $\Sigma \setminus \{\mathrm{w}_1, \dots, \mathrm{w}_\ell\}$ the sequence $|d_{A_\nu}u_\nu| + \varepsilon_\nu^{-1}|\mu(u_\nu)|$ is uniformly bounded. (As an aside: this is used in the proof of Lemma 12.2.) Hence it follows as in Case 3, that a suitable subsequence in a suitable gauge converges on this complement to a finite energy solution of (8). The limit (u, A) descends to a holomorphic curve

$$\bar{u} : \Sigma \setminus \{\mathrm{w}_1, \dots, \mathrm{w}_\ell\} \to \bar{M}$$

with finite energy. Hence, by the removable singularity theorem for $\bar{J}$-holomorphic curves, it extends to a holomorphic curve on all of Σ, still denoted by $\bar{u}$. The energy of this $\bar{J}$-holomorphic curve satisfies

$$E(\bar{u}) \leq \langle [\bar{\omega}], \bar{B}\rangle - \ell\hbar.$$

By hypothesis (H3), the dimension of the moduli space reduces by at least $2N$ at each bubble. Thus the limit $[u, A]$ belongs to a moduli space $\mathcal{M}^0_{B',S^2}$ of dimension

$$\dim \mathcal{M}^0_{B',S^2} \leq \dim \mathcal{M}^0_{B,S^2} - 2N\ell.$$

If $\{\mathrm{w}_1, \dots, \mathrm{w}_\ell\} \cap \{z_1, \dots, z_k\} = \emptyset$ then the limit curve (u, A) still satisfies $\mathrm{ev}_i([u, A]) \in f_i(Y_i)$ for every i and hence cannot exist, by the transversality condition $(H5)$. In general, denote

$$I := \{i \in \{1, \dots, k\} \,|\, z_i \notin \{\mathrm{w}_1, \dots, \mathrm{w}_\ell\}\}.$$

Then the limit $[u, A]$ satisfies

$$i \in I \qquad \Longrightarrow \qquad \mathrm{ev}_i([u, A]) \in f_i(Y_i).$$

Since the points $z_i \in \Sigma$ are pairwise distinct we have

$$\ell \geq k - |I|$$

and so

$$\dim \mathcal{M}^0_{B',S^2} \leq \dim \mathcal{M}^0_{B,S^2} - 2N(k - |I|) < \sum_{i \in I} \deg(\alpha_i).$$

Here we have used the fact that $\deg(\alpha_i) < 2N$ for each $i \in \{1, \dots, k\} \setminus I$. It follows again from $(H5)$ that such a limit curve cannot exist. Hence our assumption that the map $\mathcal{T}^{\varepsilon_i}_{B,S^2;f}$ were not surjective for every i must have been wrong. This proves the theorem. □

Remark 12.5. A more subtle argument, as in Gromov compactness for pseudoholomorphic spheres, shows that in the higher genus case the limit curve $\bar{u}$ also satisfies

$$\langle c_1(T\bar{M}), [\bar{u}] \rangle \leq \langle c_1(T\bar{M}), \bar{B} \rangle - N\ell,$$

where ℓ denotes the number of points near which bubbling occurs. Here one needs to prove that no energy gets lost and one obtains convergence to a bubble tree that represents the homology class B. With this refined compactness argument one can extend Theorem A to the higher genus case.

Remark 12.6. Assume $(H1)$, $(H2)$, and $(H4)$, but not the monotonicity hypothesis $(H3)$. Suppose that the number $\hbar > 0$ is a lower bound for the energy of the nonconstant $\bar{J}$-holomorphic spheres in $\bar{M}$ as well as for the energy of the nontrivial (that is positive energy) solutions of the vortex equations (101). Let $(\Sigma, \mathrm{dvol}_\Sigma, j_\Sigma)$ be a compact Riemann surface of genus $g > 0$ and suppose that $\bar{B} \in H_2(\bar{M}; \mathbb{Z})$ satisfies

$$0 \le \langle [\bar{\omega}], \bar{B} \rangle < \hbar.$$

Then the moduli space $\mathcal{M}^0_{B,\Sigma}$ is compact and the bubbling argument in the proof of Theorem A together with Proposition 12.3 shows that the map $\mathcal{T}^\varepsilon : \mathcal{M}^0_{B,\Sigma} \to \mathcal{M}^\varepsilon_{B,\Sigma}$ of Theorem 6.1 is a diffeomorphism for $\varepsilon > 0$ sufficiently small. Hence in this case the invariants $\Phi_{B,\Sigma}$ agree with the Gromov–Witten invariants $\mathrm{GW}_{\bar{B},\Sigma}$.

A The graph construction

Let G be a compact Lie group whose Lie algebra $\mathfrak{g} = \mathrm{Lie}(\mathrm{G})$ is equipped with an invariant inner product and (M, ω) be a symplectic manifold with a Hamiltonian G-action generated by a moment map $\mu : M \to \mathfrak{g}$. We denote by $\mathfrak{g} \to \mathrm{Vect}(M) : \eta \mapsto X_\eta$ the infinitesimal action, by $C^\infty_{\mathrm{G}}(M)$ the space of G-invariant smooth functions on M, and by $\mathcal{J}_{\mathrm{G}}(M, \omega)$ the space of G-invariant and ω-compatible almost complex structures on M. We fix a Riemann surface $(\Sigma, \mathrm{dvol}_\Sigma, j_\Sigma)$ and a principal G-bundle $P \to \Sigma$. Given a family of almost complex structures $\Sigma \to \mathcal{J}_{\mathrm{G}}(M, \omega) : z \mapsto J_z$ and a 1-form $T\Sigma \to C^\infty_{\mathrm{G}}(M) : \hat{z} \mapsto H_{\hat{z}}$ we consider the perturbed equations

$$\bar{\partial}_{J,H,A}(u) = 0, \qquad *F_A + \varepsilon^{-2}\mu(u) = 0, \tag{106}$$

where

$$\bar{\partial}_{J,H,A}(u) := \bar{\partial}_{J,A}(u) + X_H(u)^{0,1}.$$

Here the $(0,1)$-form $\bar{\partial}_{J,A}(u) \in \Omega^{0,1}(\Sigma, u^*TM/\mathrm{G})$ is understood with respect to the family of almost complex structures J_z, parametrized by $z \in \Sigma$. Moreover, the Hamiltonian perturbation is defined as follows. Associated to $H \in \Omega^1(\Sigma, C^\infty_{\mathrm{G}}(M))$ is the 1-form $X_H \in \Omega^1(\Sigma, \mathrm{Vect}_{\mathrm{G}}(M, \omega))$ which assigns to every $\hat{z} \in T_z\Sigma$ the Hamiltonian vector field $X_{H,\hat{z}}$ associated to the Hamiltonian function $H_{\hat{z}} : M \to \mathbb{R}$. Thus $\iota(X_{H,\hat{z}})\omega = dH_{\hat{z}}$. The 1-form

$X_H(u) \in \Omega^1(\Sigma, u^*TM/\mathrm{G})$ lifts to an equivariant and horizontal 1-form on P with values in u^*TM, also denoted by $X_H(u)$ and defined by

$$(X_H(u))_p(v) := X_{H,d\pi(p)v}(u(p)).$$

The complex anti-linear part of this 1-form is the Hamiltonian term in the definition of $\bar\partial_{J,H,A}(u)$. In this section we show how to reduce the perturbed equations (106) to (7) via Gromov's graph construction [Gr].

Let us denote by $\alpha_H \in \Omega^1(\Sigma \times M)$ the 1-form associated to H. Thus α_H assigns to every pair of tangent vactors $(\hat z, \hat x) \in T_z\Sigma \times T_xM$ the real number $H_{\hat z}(x)$. Denote

$$\tilde M := \Sigma \times M.$$

The 2-form

$$\tilde\omega := \omega - d\alpha_H + c\,\mathrm{dvol}_\Sigma$$

is a symplectic form on $\tilde M$ whenever the constant c is sufficiently large. Here we have abused notation and denoted by ω the pullback of the 2-form ω on M under the obvious projection $\Sigma \times M \to M$ and likewise for dvol_Σ. To see that $\tilde\omega$ is symplectic for large c, note first that $\tilde\omega$ is a connection form: it is closed and its restriction to each fibre $\{z\} \times M$ is symplectic. The curvature of this connection form is the 2-form

$$\Omega_H\,\mathrm{dvol}_\Sigma := dH + \frac{1}{2}\{H \wedge H\} \in \Omega^2(\Sigma, C^\infty_{\mathrm{G}}(M)).$$

This identity defines the function $\Omega_H : \Sigma \times M \to \mathbb{R}$. Now the top exterior power of $\tilde\omega$ is given by

$$\frac{\tilde\omega^{n+1}}{(n+1)!} = (c - \Omega_H)\frac{\omega^n}{n!} \wedge \mathrm{dvol}_\Sigma,$$

where $\dim M = 2n$. Hence $\tilde\omega$ is nondegenerate whenever $c > \max \Omega_H$. Now consider the almost complex structure $\tilde J$ on $\tilde M$ given by

$$\tilde J(z,x) := \left(\begin{array}{cc} j_\Sigma(z) & 0 \\ J(z,x) \circ X_H(z,x) - X_H(z,x) \circ j_\Sigma(z) & J(z,x) \end{array} \right).$$

Here $J(z,x) := J_z(x)$ and we denote by $X_H(z,x) : T_z\Sigma \to T_xM$ the linear map $\hat z \mapsto X_{H,\hat z}(x)$. Lemma A.2 below shows that $\tilde J$ is compatible with $\tilde\omega$.

Lemma A.1. *Let* $(u, A) \in C^\infty_{\mathrm{G}}(P, M) \times \mathcal{A}(P)$ *and define* $\tilde{u} : P \to \tilde{M}$ *by* $\tilde{u}(p) := (\pi(p), u(p))$. *Then* u *and* A *satisfy (106) if and only if* $\tilde{u}$ *and* A *satisfy*

$$\bar{\partial}_{\tilde{J},A}(\tilde{u}) = 0, \qquad *F_A + \varepsilon^{-2}\tilde{\mu}(\tilde{u}) = 0.$$

Here $\tilde{\mu} : \tilde{M} \to \mathfrak{g}$ *is defined by* $\tilde{\mu}(z, x) := \mu(x)$.

Proof. By definition of $\tilde{J}$ we have $\bar{\partial}_{\tilde{J},A}(\tilde{u}) = (0, \bar{\partial}_{J,H,A}(u))$. Alternatively, we can compute in local holomorphic coordinates $s+it$ on Σ. In such coordinates the Hamiltonian perturbation, the connection A, and the volume form on Σ have the form

$$H = F\,ds + G\,dt, \qquad A = \Phi\,ds + \Psi\,dt, \qquad \mathrm{dvol}_\Sigma = \lambda^2\,ds \wedge dt$$

and the equations (106) have the form

$$\begin{aligned} \partial_s u + L_u\Phi + X_F(u) + J(\partial_t u + L_u\Psi + X_G(u)) &= 0, \\ \partial_s\Psi - \partial_t\Phi + [\Phi, \Psi] + (\lambda/\varepsilon)^2\mu(u) &= 0. \end{aligned}$$

Moreover, the almost complex structure $\tilde{J}$ is given by

$$\tilde{J} := \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ JX_F - X_G & JX_G + X_F & J \end{pmatrix}.$$

This proves the lemma. □

Lemma A.2. *Let* $\hat{z}_i \in T_z\Sigma$ *and* $\hat{x}_i \in T_xM$ *for* $i = 1, 2$. *Then*

$$\tilde{\omega}((\hat{z}_1, \hat{x}_1), (\hat{z}_2, \hat{x}_2)) = \langle \hat{x}_1 + X_{H,\hat{z}_1}(x), \hat{x}_2 + X_{H,\hat{z}_2}(x)\rangle_z + (c - \Omega_H)\langle \hat{z}_1, \hat{z}_2\rangle,$$

where $\langle\cdot,\cdot\rangle_z := \omega(\cdot, J_z\cdot)$.

Proof. Continue the notation of the proof of Lemma A.1. Then the curvature Ω_H is given by

$$\lambda^2\Omega_H = \partial_s G - \partial_t F + \{F, G\},$$

where $\{F, G\} := \omega(X_F, X_G)$ denotes the Poisson bracket on M, and

$$\tilde{\omega} = \omega - dF \wedge ds - dG \wedge dt + (\partial_t F - \partial_s G + c\lambda^2)ds \wedge dt$$

where dF and dG denote the differential on M. Abbreviate $\zeta_i := (\hat{s}_i, \hat{t}_i, \hat{x}_i)$ and $\xi_i := \hat{x}_i + \hat{s}_i X_F + \hat{t}_i X_G$ for $i = 1, 2$. Then

$$\begin{aligned}
\tilde{\omega}(\zeta_1, \tilde{J}\zeta_2) &= \tilde{\omega}(\zeta_1, (-\hat{t}_2, \hat{s}_2, J\xi_2 + \hat{t}_2 X_F - \hat{s}_2 X_G)) \\
&= \omega(\hat{x}_1, J\xi_2 + \hat{t}_2 X_F - \hat{s}_2 X_G) \\
&\quad + \hat{t}_2 dF(\hat{x}_1) + \hat{s}_1 dF(J\xi_2 + \hat{t}_2 X_F - \hat{s}_2 X_G) \\
&\quad - \hat{s}_2 dG(\hat{x}_1) + \hat{t}_1 dG(J\xi_2 + \hat{t}_2 X_F - \hat{s}_2 X_G) \\
&\quad + (\partial_t F - \partial_s G + c\lambda^2)(\hat{s}_1 \hat{s}_2 + \hat{t}_1 \hat{t}_2) \\
&= \omega(\xi_1, J\xi_2) + (c\lambda^2 - \partial_s G + \partial_t F - \{F, G\})(\hat{s}_1 \hat{s}_2 + \hat{t}_1 \hat{t}_2).
\end{aligned}$$

The last identity follows from the fact that $\{F, G\} = dF(X_G) = -dG(X_F)$ and $dF(J\xi_2) = \omega(X_F, J\xi_2)$. $\square$

B Cauchy–Riemann operators

Fix a compact Lie group G, an invariant inner product on the Lie algebra $\mathfrak{g} = \mathrm{Lie}(\mathrm{G})$, a symplectic manifold (M, ω), a Hamiltonian G-action on M generated by a moment map $\mu : M \to \mathfrak{g}$, a compact Riemann surface $(\Sigma, j_\Sigma, \mathrm{dvol}_\Sigma)$, a principal G-bundle $P \to \Sigma$, and a family of G-invariant and ω-compatible almost complex structures $\Sigma \to \mathcal{J}_{\mathrm{G}}(M, \omega) : z \mapsto J_z$. Each almost complex structure determines a Riemannian metric $\langle \cdot, \cdot \rangle_z := \omega(\cdot, J_z \cdot)$ on M and hence a Levi-Civita connection $\nabla = \nabla_z$. The value of z will usually be clear from the context and we shall omit the subscript z. Let $u : P \to M$ be an equivariant smooth map and A be a connection on P. Then A and ∇ determine a connection ∇_A on u^*TM/G given by

$$\nabla_A \xi := \nabla \xi + \nabla_\xi X_A(u)$$

for $\xi \in \Omega^0(\Sigma, u^*TM/\mathrm{G})$. More precisely, we think of A as an equivariant 1-form on P with values in the Lie algebra $\mathfrak{g}$ which identifies the vertical tangent bundle with $\mathfrak{g}$. A section ξ of u^*TM/G lifts to an equivariant section of the bundle $u^*TM \to P$ (also denoted by ξ) and a 1-form $\theta \in \Omega^1(\Sigma, u^*TM/\mathrm{G})$ lifts to an equivariant and horizontal 1-form on P with values in u^*TM (also denoted by θ). In this notation the 1-form $\nabla_A \xi$ is given by

$$(\nabla_A \xi)_p(v) := \nabla_v \xi(p) + \nabla_{\xi(p)} X_{A_p(v)}(u(p))$$

for $v \in T_pP$. In general, ∇_A preserves neither the inner product nor the complex structure on u^*TM/G. More precisely, let $J_u \in \Omega^0(P, \mathrm{End}(u^*TM))$

be given by $J_u(p) := J_{\pi(p)}(u(p)) \in \mathrm{End}(T_{u(p)}M)$. This section is equivariant and hence descend to a complex structure, also denoted by J_u, of the bundle $\mathrm{End}(u^*TM/\mathrm{G}) \to \Sigma$.

Lemma B.1. *The covariant derivative of J_u is given by*

$$\nabla_A J_u = \nabla_{d_A u} J(u) + \dot{J}(u).$$

where $\dot{J}(u) \in \Omega^1(\Sigma, \mathrm{End}(u^*TM/\mathrm{G}))$ is defined by

$$\dot{J}(u)_p(v) := \left.\frac{d}{dt}\right|_{t=0} J_{\gamma(t)}(u(p))$$

for $v \in T_pP$ and a smooth path $\gamma : \mathbb{R} \to P$ such that $\gamma(0) = p$ and $\dot{\gamma}(0) = v$.

Proof. Since J is G-invariant we have $\mathcal{L}_{X_\eta} J = 0$ for every $\eta \in \mathfrak{g}$. This formula can be expressed in the form

$$(\nabla_{X_\eta} J)\xi = \nabla_{J\xi} X_\eta - J\nabla_\xi X_\eta. \tag{107}$$

Using this formula we obtain

$$\begin{aligned}
(\nabla_A J_u)\xi &= \nabla_A(J_u\xi) - J_u(\nabla_A\xi) \\
&= \nabla(J_u\xi) - J_u\nabla\xi + \nabla_{J_u\xi}X_A(u) - J_u\nabla_\xi X_A(u) \\
&= \Big(\nabla_{du}J(u) + \dot{J}(u)\Big)\,\xi + \big(\nabla_{X_A(u)}J(u)\big)\,\xi \\
&= \Big(\nabla_{d_A u}J(u) + \dot{J}(u)\Big)\,\xi
\end{aligned}$$

as claimed. □

It follows from Lemma B.1 that the complex linear part of the connection ∇_A is the connection $\tilde{\nabla}_A$ on u^*TM/G given by

$$\begin{aligned}
\tilde{\nabla}_A\xi &:= \nabla_A\xi - \frac{1}{2}J_u(\nabla_A J_u)\xi \qquad (108) \\
&= \nabla\xi + \nabla_\xi X_A(u) - \frac{1}{2}J_u\Big(\nabla_{d_A u}J(u) + \dot{J}(u)\Big)\,\xi.
\end{aligned}$$

Lemma B.2. *$\tilde{\nabla}_A$ is a Hermitian connection on u^*TM/G.*

Proof. We shall use the identity $\langle (\nabla_{\xi_1} J)\xi_2, \xi_3\rangle +$cyclic $= 0$. By (107), we have

$$\begin{aligned}
&\langle \xi_1, \nabla_{\xi_2} X_\eta\rangle + \langle \nabla_{\xi_1} X_\eta, \xi_2\rangle \\
&= \langle J\xi_1, J\nabla_{\xi_2} X_\eta\rangle + \langle J\nabla_{\xi_1} X_\eta, J\xi_2\rangle \\
&= \langle J\xi_1, \nabla_{\xi_2}(JX_\eta) - (\nabla_{\xi_2} J)X_\eta\rangle + \langle \nabla_{J\xi_1} X_\eta - (\nabla_{X_\eta} J)\xi_1, J\xi_2\rangle \\
&= \langle J\xi_1, \nabla_{\xi_2}(JX_\eta)\rangle + \langle \nabla_{J\xi_1} X_\eta, J\xi_2\rangle - \langle J\xi_1, (\nabla_{\xi_2} J)X_\eta\rangle - \langle \xi_2, (\nabla_{X_\eta} J)J\xi_1\rangle \\
&= \langle J\xi_1, \nabla_{\xi_2}(JX_\eta)\rangle + \langle \nabla_{J\xi_1} X_\eta, J\xi_2\rangle + \langle X_\eta, (\nabla_{J\xi_1} J)\xi_2\rangle \\
&= \langle \nabla_{J\xi_1}(JX_\eta), \xi_2\rangle - \langle J(\nabla_{J\xi_1} X_\eta), \xi_2\rangle - \langle (\nabla_{J\xi_1} J)X_\eta, \xi_2\rangle \\
&= 0
\end{aligned}$$

for $\xi_1, \xi_2 \in T_xM$ and $\eta \in \mathfrak{g}$. Here the penultimate equality follows from the fact that JX_η is a gradient vector field and that $\nabla_{J\xi_1} J$ is skew-adjoint. This shows that $\nabla X_A(u)$ is a 1-form on Σ with values in the bundle of skew-Hermitian endomorphisms of u^*TM/G, and so is $J(\nabla_{d_A u} J)$. Moreover, since

$$d\langle \xi_1, \xi_2\rangle = \langle \nabla\xi_1, \xi_2\rangle + \langle \xi_1, \nabla\xi_2\rangle - \langle \xi_1, J\dot{J}\xi_2\rangle,$$

the operator $\xi \mapsto \nabla\xi - \frac{1}{2}J\dot{J}\xi$ is a Riemannian connection. By (108), this shows that $\tilde{\nabla}_A$ is a Riemannian connection. It follows directly from the definition that $\tilde{\nabla}_A$ preserves the complex structure J_u. □

Lemma B.3. *For every gauge transformation $g \in \mathcal{G}(P)$ and every section $\xi \in \Omega^0(\Sigma, u^*TM/\mathrm{G})$ we have*

$$\nabla_{g^*A}(g^{-1}\xi) = g^{-1}\nabla_A\xi, \qquad \tilde{\nabla}_{g^*A}(g^{-1}\xi) = g^{-1}\tilde{\nabla}_A\xi.$$

Proof. Since the metric $\langle \cdot, \cdot\rangle_z$ is G-invariant for every $z \in \Sigma$ we have

$$\nabla(g^{-1}\xi) = g^{-1}\nabla\xi - \nabla_{g^{-1}\xi} X_{g^{-1}dg}(g^{-1}u).$$

Hence the first identity follows from the fact that $g^*A = g^{-1}dg + g^{-1}Ag$ and that $\nabla_{g^{-1}\xi} X_{g^{-1}\eta g}(g^{-1}x) = g^{-1}\nabla_\xi X_\eta(x)$. The second identity follows from the first and the fact that J_z is G-invariant for every z. □

Lemma B.4. *Suppose that J is independent of $z \in \Sigma$. Then the curvature of the connection ∇_A is the equivariant and horizontal 2-form $F^{\nabla_A} \in \Omega^2(P, \mathrm{End}(u^*TM))$ given by*

$$F^{\nabla_A}(v_1, v_2)\xi = R(d_A u(p)v_1, d_A u(p)v_2)\xi + \nabla_\xi X_{F_A(v_1,v_2)}(u(p))$$

for $v_1, v_2 \in T_pP$ *and* $\xi \in T_{u(p)}M$*, where* $R \in \Omega^2(M, \mathrm{End}(TM))$ *is the Riemann curvature tensor of the metric* $\langle\cdot,\cdot\rangle = \omega(\cdot, J\cdot)$*. This 2-form descends to a 2-form on* Σ *with values in* $\mathrm{End}(u^*TM/\mathrm{G})$*, also denoted by* F^{∇_A}*.*

Proof. Given a map $\mathbb{R}^2 \to M : (s,t) \mapsto u(s,t)$, a vector field $\xi(s,t) \in T_{u(s,t)}M$ along u, and a G-connection $A = \Phi\, ds + \Psi\, dt$, where $\Phi, \Psi : \mathbb{R}^2 \to \mathfrak{g}$, we denote

$$\begin{array}{rclrcl} v_s & := & \partial_s u + X_\Phi(u), & v_t & := & \partial_t u + X_\Psi(u), \\ \nabla_{A,s}\xi & := & \nabla_s\xi + \nabla_\xi X_\Phi(u), & \nabla_{A,t}\xi & := & \nabla_t\xi + \nabla_\xi X_\Psi(u). \end{array}$$

Then the assertion can be restated in the form

$$\nabla_{A,s}\nabla_{A,t}\xi - \nabla_{A,t}\nabla_{A,s}\xi = R(v_s, v_t)\xi + \nabla_\xi X_{\partial_s\Psi - \partial_t\Phi + [\Phi,\Psi]}(u).$$

To prove this we use the identities

$$\begin{array}{rcl} \nabla_{A,s}\nabla_{A,t}\xi - \nabla_{A,t}\nabla_{A,s}\xi & = & \nabla_s\nabla_t\xi - \nabla_t\nabla_s\xi + \nabla_s\nabla_\xi X_\Psi(u) - \nabla_{\nabla_s\xi}X_\Psi(u) \\ & & - \nabla_t\nabla_\xi X_\Phi(u) + \nabla_{\nabla_t\xi}X_\Phi(u) \\ & & + \nabla_{\nabla_\xi X_\Psi}X_\Phi(u) - \nabla_{\nabla_\xi X_\Phi}X_\Psi(u), \\ R(\partial_s u, \partial_t u)\xi & = & \nabla_s\nabla_t\xi - \nabla_t\nabla_s\xi, \\ R(\partial_s u, X_\Psi(u))\xi & = & \nabla_s\nabla_\xi X_\Psi(u) - \nabla_{\nabla_s\xi}X_\Psi(u) - \nabla_\xi X_{\partial_s\Psi}(u), \\ R(X_\Phi(u), X_\Psi(u))\xi & = & -\nabla_\xi X_{[\Phi,\Psi]}(u) + \nabla_{\nabla_\xi X_\Psi}X_\Phi(u) - \nabla_{\nabla_\xi X_\Phi}X_\Psi(u). \end{array}$$

The first and second identities are the definition of the connection ∇_A and the curvature tensor R. The other identities use the equations

$$\nabla_{X_\eta}Z - \nabla_Z X_\eta = [Z, X_\eta] = 0, \qquad \nabla_Z[X_{\eta_1}, X_{\eta_2}] = [\nabla_Z X_{\eta_1}, X_{\eta_2}],$$

for every G-invariant vector field $Z \in \mathrm{Vect}_\mathrm{G}(M)$ and all $\eta, \eta_1, \eta_2 \in \mathfrak{g}$. □

Now consider the Cauchy–Riemann operator

$$D := D_{(u,A)} : \Omega^0(\Sigma, u^*TM/\mathrm{G}) \to \Omega^{0,1}(\Sigma, u^*TM/\mathrm{G})$$

given by

$$D\xi := (\nabla_A\xi)^{0,1} - \frac{1}{2}J(\nabla_\xi J)\partial_{J,A}(u). \tag{109}$$

In the case $\bar\partial_{J,A}(u) = 0$ this operator is the vertical differential of the section $u \mapsto \bar\partial_{J,A}(u)$ of the infinite dimensional vector bundle over the space $C^\infty_\mathrm{G}(P, M)$ with fibre $\Omega^{0,1}(\Sigma, u^*TM/\mathrm{G})$ over u.

In the following we denote the Nijenhuis tensor of J by $N \in \Omega^2(TM, TM)$. It is given by

$$\begin{aligned} N(\xi_1, \xi_2) &= [\xi_1, \xi_2] + J[J\xi_1, \xi_2] + J[\xi_1, J\xi_2] - [J\xi_1, J\xi_2] \\ &= 2J(\nabla_{\xi_2} J)\xi_1 - 2J(\nabla_{\xi_1} J)\xi_2. \end{aligned}$$

Lemma B.5. *The complex linear part of D is the operator $\xi \mapsto (\tilde{\nabla}_A \xi)^{0,1}$. Moreover,*

$$D\xi = (\tilde{\nabla}_A \xi)^{0,1} + \frac{1}{4} N(\xi, \partial_{J,A}(u)) + \frac{1}{2}(J\dot{J}\xi)^{0,1}.$$

Proof. By definition of $\tilde{\nabla}_A$, we have

$$D\xi = (\tilde{\nabla}_A \xi)^{0,1} + \frac{1}{2} J \left(\nabla_{\partial_{J,A}(u)} J(u) + \dot{J}(u)^{0,1} \right) \xi - \frac{1}{2} J(\nabla_\xi J)\partial_{J,A}(u).$$

Hence the formula for $D\xi$ follows from the relation between the Nijenhuis tensor and ∇J. Now this equation shows that the operator $\xi \mapsto D\xi - (\tilde{\nabla}_A \xi)^{0,1}$ is complex anti-linear. □

C Invariant metrics

Let M be a (complete) Riemannian m-manifold. For $v \in T_x M$ and $i, j \in \{1, 2\}$ there exist linear maps

$$E_i(x, v) : T_x M \to T_{\exp_x(v)} M, \qquad E_{ij}(x, v) : T_x M \oplus T_x M \to T_{\exp_x(v)} M$$

characterized by the following conditions. If $x : \mathbb{R} \to M$ is a smooth curve and $v, w : \mathbb{R} \to x^* TM$ are vector fields along x then

$$\begin{aligned} \frac{d}{dt} \exp_{x(t)}(v(t)) &= E_1(x, v)\dot{x} + E_2(x, v)\nabla_t v, \\ \nabla_t (E_1(x, v)w) &= E_{11}(x, v)(w, \dot{x}) + E_{12}(x, v)(w, \nabla_t v) + E_1(x, v)\nabla_t w, \\ \nabla_t (E_2(x, v)w) &= E_{21}(x, v)(w, \dot{x}) + E_{22}(x, v)(w, \nabla_t v) + E_2(x, v)\nabla_t w. \end{aligned}$$

Note that the map $E_{11}(x, v)(w, w')$ is not symmetric in w and w'. It satisfies

$$E_{11}(x, v)(w, w') - E_{11}(x, v)(w', w) = E_2(x, v) R(w, w') v,$$

where $R \in \Omega^2(M, \mathrm{End}(TM))$ denotes the curvature tensor. However,

$$E_{12}(x,v)(w,w') = E_{21}(x,v)(w',w),$$

and $E_{22}(x,v)(w,w')$ is symmetric in w and w'. (See [Ga] for more details.) Now let $\mathrm{G} \times M \to M : (g,x) \mapsto gx$ be a smooth action of a compact Lie group G with infinitesimal action $\mathfrak{g} \to \mathrm{Vect}(M) : \eta \mapsto X_\eta$. Assume that M is equipped with a G-invariant Riemannian metric.

Lemma C.1.

$$\begin{aligned} X_\eta(\exp_x(v)) &= E_1(x,v)X_\eta(x) + E_2(x,v)\nabla_v X_\eta(x), \\ \nabla_{E_i(x,v)w} X_\eta(\exp_x(v)) &= E_{i1}(x,v)(w, X_\eta(x)) + E_{i2}(x,v)(w, \nabla_v X_\eta(x)) \\ &\quad + E_i(x,v)\nabla_w X_\eta(x). \end{aligned}$$

Proof. Since the group action preserves geodesics, $g\exp_x(v) = \exp_{gx}(gv)$. Differentiate this identity with respect to g to obtain the first identity. To prove the second differentiate the first identity covariantly and use the definition of E_i and E_{ij}. For more details see [Ga]. □

For each $x \in M$ denote by $L_x : \mathfrak{g} \to T_xM$ the infinitesimal action, i.e. $L_x\eta = X_\eta(x)$. Given a map $u : \mathbb{R}^2 \to M$, a vector field $\xi : \mathbb{R}^2 \to u^*TM$ along u, a function $\eta : \mathbb{R}^2 \to \mathfrak{g}$, and a G-connection $A = \Phi\, ds + \Psi\, dt$, where $\Phi, \Psi : \mathbb{R}^2 \to \mathfrak{g}$, we denote

$$\begin{aligned} v_s &= \partial_s u + X_\Phi(u), & v_t &= \partial_t u + X_\Psi(u), \\ \nabla_{A,s}\xi &= \nabla_s\xi + \nabla_\xi X_\Phi(u), & \nabla_{A,t}\xi &= \nabla_t\xi + \nabla_\xi X_\Psi(u), \\ \nabla_{A,s}\eta &= \partial_s\eta + [\Phi,\eta], & \nabla_{A,t}\eta &= \partial_t\eta + [\Psi,\eta]. \end{aligned}$$

Define $\rho \in \Omega^2(M, \mathrm{G})$ by

$$\langle \eta, \rho(\xi,\xi')\rangle := \langle \nabla_\xi X_\eta(x), \xi'\rangle = -\langle \nabla_{\xi'} X_\eta(x), \xi\rangle$$

for $\xi, \xi' \in T_xM$ and $\eta \in \mathfrak{g}$.

Lemma C.2. *With the above notation we have*

$$\begin{aligned} \nabla_{A,s}L_u\eta - L_u\nabla_{A,s}\eta &= \nabla_{v_s}X_\eta(u), & \nabla_{A,t}L_u\eta - L_u\nabla_{A,t}\eta &= \nabla_{v_t}X_\eta(u), \\ \nabla_{A,s}L_u^*\xi - L_u^*\nabla_{A,s}\xi &= \rho(v_s,\xi), & \nabla_{A,t}L_u^*\xi - L_u^*\nabla_{A,t}\xi &= \rho(v_t,\xi) \end{aligned}$$

Proof. See [Ga]. □

Now let M^* denote the subset of all points $x \in M$ with finite isotropy subgroup $\mathrm{G}_x := \{g \in \mathrm{G} \,|\, gx = x\}$. Thus $x \in M^*$ if and only if the linear map $L_x : \mathfrak{g} \to T_xM$ is injective. Hence, for every map $u_0 : \mathbb{R}^2 \to M^*$ there exists a unique G-connection $A_0 = \Phi_0\, ds + \Psi_0\, dt$ such that

$$L_{u_0}^* v_{0s} = L_{u_0}^* v_{0t} = 0,$$

where

$$v_{0s} := \partial_s u_0 + L_{u_0}\Phi_0, \qquad v_{0t} := \partial_t u_0 + L_{u_0}\Psi_0.$$

Let $\xi_0 : \mathbb{R}^2 \to u_0^*TM$ be a vector field along u_0, consider the map

$$u(s,t) := \exp_{u_0(s,t)}(\xi_0(s,t)).$$

and abbreviate

$$\varphi_0 := \Phi - \Phi_0, \qquad \psi_0 := \Psi - \Psi_0.$$

Lemma C.3.

$$\begin{aligned} L_u\varphi_0 &= v_s - E_1(u_0,\xi_0)v_{0s} - E_2(u_0,\xi_0)\nabla_{A_0,s}\xi_0, \\ L_u\psi_0 &= v_t - E_1(u_0,\xi_0)v_{0t} - E_2(u_0,\xi_0)\nabla_{A_0,t}\xi_0. \end{aligned}$$

Proof. We compute

$$\begin{aligned} L_u(\Phi - \Phi_0) &= X_\Phi(u) - E_1(u_0,\xi_0)X_{\Phi_0}(u_0) - E_2(u_0,\xi_0)\nabla_{\xi_0}X_{\Phi_0}(u_0) \\ &= \partial_s u - E_1(u_0,\xi_0)\partial_s u_0 - E_2(u_0,\xi_0)\nabla_s\xi_0 \\ &\quad + X_\Phi(u) - E_1(u_0,\xi_0)X_{\Phi_0}(u_0) - E_2(u_0,\xi_0)\nabla_{\xi_0}X_{\Phi_0}(u_0) \\ &= v_s - E_1(u_0,\xi_0)v_{0s} - E_2(u_0,\xi_0)\nabla_{A_0,s}\xi_0. \end{aligned}$$

Here the first equation follows from Lemma C.1, the second from the definition of E_i, and the last from the definitions of v_s and v_{0s}. ☐

In the proof we did not use the fact that $L_{u_0}^* v_{0s} = L_{u_0}^* v_{0t} = 0$. Now suppose $L_{u_0}^*\xi_0 = 0$. Then, by Lemma C.2,

$$L_{u_0}^*\nabla_{A_0,s}\xi_0 = -\rho(v_{0s},\xi_0), \qquad L_{u_0}^*\nabla_{A_0,t}\xi_0 = -\rho(v_{0t},\xi_0).$$

Abbreviating $E_1 = E_1(u_0,\xi_0)$ and $E_2 = E_2(u_0,\xi_0)$ we obtain the following.

Corollary C.4. *If* $L_{u_0}^*\xi_0 = 0$ *then*

$$\begin{aligned} L_{u_0}^* {E_2}^{-1} L_u(\Phi - \Phi_0) &= L_{u_0}^* {E_2}^{-1}(v_s - E_1 v_{0s}) + \rho(v_{0s},\xi_0), \\ L_{u_0}^* {E_2}^{-1} L_u(\Psi - \Psi_0) &= L_{u_0}^* {E_2}^{-1}(v_t - E_1 v_{0t}) + \rho(v_{0t},\xi_0). \end{aligned}$$

Proof. Lemma C.3. □

Lemma C.5. *Assume $L_{u_0}^* \xi_0 = 0$ and abbreviate $E_i := E_i(u_0, \xi_0)$ and $E_{ij} := E_{ij}(u_0, \xi_0)$. Then*

$$\begin{aligned} L_u \nabla_{A,t} \varphi_0 \;&=\; \nabla_{A,t} v_s + \nabla_{X_{\varphi_0}} X_{\psi_0}(u) - \nabla_{v_t} X_{\varphi_0}(u) - \nabla_{v_s} X_{\psi_0}(u) \\ &\quad - E_{11}(v_{0s}, v_{0t}) - E_{12}(v_{0s}, \nabla_{A_0,t}\xi_0) \\ &\quad - E_{21}(\nabla_{A_0,s}\xi_0, v_{0t}) - E_{22}(\nabla_{A_0,s}\xi_0, \nabla_{A_0,t}\xi_0) \\ &\quad - E_1 \nabla_{A_0,t} v_{0s} - E_2 \nabla_{A_0,t} \nabla_{A_0,s} \xi_0, \end{aligned}$$

$$\begin{aligned} L_u \nabla_{A,s} \varphi_0 \;&=\; \nabla_{A,s} v_s + \nabla_{X_{\varphi_0}} X_{\varphi_0}(u) - 2\nabla_{v_s} X_{\varphi_0}(u) \\ &\quad - E_{11}(v_{0s}, v_{0s}) - E_{12}(v_{0s}, \nabla_{A_0,s}\xi_0) \\ &\quad - E_{21}(\nabla_{A_0,s}\xi_0, v_{0s}) - E_{22}(\nabla_{A_0,s}\xi_0, \nabla_{A_0,s}\xi_0) \\ &\quad - E_1 \nabla_{A_0,s} v_{0s} - E_2 \nabla_{A_0,s} \nabla_{A_0,s} \xi_0. \end{aligned}$$

Proof. We only prove the first identity. The proof of the second is similar. By Lemmata C.2 and C.3,

$$\begin{aligned} L_u \nabla_{A,t} \varphi_0 \;&=\; \nabla_{A,t} L_u \varphi_0 - \nabla_{v_t} X_{\varphi^0}(u) \\ &=\; \nabla_{A,t} v_s - \nabla_{A,t}(E_1 v_{0s}) - \nabla_{A,t}(E_2 \nabla_{A_0,s}\xi_0) - \nabla_{v_t} X_{\varphi_0}(u). \end{aligned}$$

Hence, by the definition of E_{ij} and Lemma C.1,

$$\begin{aligned} \nabla_{A,t}(E_1 v_{0s}) \;&=\; \nabla_t(E_1 v_{0s}) + \nabla_{E_1 v_{0s}} X_\Psi(u) \\ &=\; E_{11}(v_{0s}, \partial_t u_0) + E_{12}(v_{0s}, \nabla_t \xi_0) \\ &\quad + E_{11}(v_{0s}, X_{\Psi_0}(u_0)) + E_{12}(v_{0s}, \nabla_{\xi_0} X_{\Psi_0}(u_0)) \\ &\quad + E_1 \nabla_t v_{0s} + \nabla_{E_1 v_{0s}} X_{\psi_0}(u) + E_1 \nabla_{v_{0s}} X_{\Psi_0}(u_0) \\ &=\; \nabla_{E_1 v_{0s}} X_{\psi_0}(u) + E_{11}(v_{0s}, v_{0t}) + E_{12}(v_{0s}, \nabla_{A_0,t}\xi_0) + E_1 \nabla_{A_0,t} v_{0s} \end{aligned}$$

and

$$\begin{aligned} \nabla_{A,t}(E_2 \nabla_{A_0,s}\xi_0) \;&=\; \nabla_t(E_2 \nabla_{A_0,s}\xi_0) + \nabla_{E_2 \nabla_{A_0,s}\xi_0} X_\Psi(u) \\ &=\; E_{21}(\nabla_{A_0,s}\xi_0, \partial_t u_0) + E_{22}(\nabla_{A_0,s}\xi_0, \nabla_t \xi_0) + E_2 \nabla_t \nabla_{A_0,s}\xi_0 \\ &\quad + \nabla_{E_2 \nabla_{A_0,s}\xi_0} X_{\psi_0}(u) + E_{21}(\nabla_{A_0,s}\xi_0, X_{\Psi_0}(u_0)) \\ &\quad + E_{22}(\nabla_{A_0,s}\xi_0, \nabla_{\xi_0} X_{\Psi_0}(u_0)) + E_2 \nabla_{\nabla_{A_0,s}\xi_0} X_{\Psi_0}(u_0) \\ &=\; \nabla_{E_2 \nabla_{A_0,s}\xi_0} X_{\psi_0}(u) + E_{21}(\nabla_{A_0,s}\xi_0, v_{0t}) \\ &\quad + E_{22}(\nabla_{A_0,s}\xi_0, \nabla_{A_0,t}\xi_0) + E_2 \nabla_{A_0,t} \nabla_{A_0,s}\xi_0. \end{aligned}$$

Inserting these two identities into the previous formula we obtain

$$\begin{aligned} L_u \nabla_{A,t}\varphi_0 &= \nabla_{A,t} v_s - \nabla_{v_t} X_{\varphi_0}(u) \\ &\quad -\nabla_{E_1 v_{0s}} X_{\psi_0}(u) - \nabla_{E_2 \nabla_{A_0,s}\xi_0} X_{\psi_0}(u) \\ &\quad -E_{11}(v_{0s}, v_{0t}) - E_{12}(v_{0s}, \nabla_{A_0,t}\xi_0) \\ &\quad -E_{21}(\nabla_{A_0,s}\xi_0, v_{0t}) - E_{22}(\nabla_{A_0,s}\xi_0, \nabla_{A_0,t}\xi_0) \\ &\quad -E_1 \nabla_{A_0,t} v_{0s} - E_2 \nabla_{A_0,t}\nabla_{A_0,s}\xi_0. \end{aligned}$$

Now the result follows from Lemma C.3. □

References

[AB] M.F. Atiyah and R. Bott, The Yang–Mills equations over Riemann surfaces, *Phil. Trans. R. Soc. Lond. A* **308** (1982), 523–615.

[CGS] C. Cieliebak, A.R. Gaio, D.A. Salamon, *J*-homolorphic curves, moment maps and invariants of Hamiltonian group actions, *Int. Math. Res. Notes* **10** (2000), 831–882.

[CGMS] C. Cieliebak, R. Gaio, I. Mundet, D.A. Salamon, Invariants of Hamiltonian group actions, in preparation.

[DK] S.K. Donaldson and P.B. Kronheimer, *The Geometry of Four-Manifolds*, Oxford University Press, 1990.

[DS1] S. Dostoglou and D.A. Salamon, Cauchy-Riemann operators, self-duality, and the spectral flow, in *First European Congress of Mathematics, Volume I, Invited Lectures (Part 1)*, edited by A. Joseph, F. Mignot, F. Murat, B. Prum, R. Rentschler, Birkhäuser Verlag, Progress in Mathematics, **Vol. 119**, 1994, pp. 511–545.

[DS2] S. Dostoglou and D.A. Salamon, Self-dual instantons and holomorphic curves, *Annals of Mathematics* **139** (1994), 581–640.

[Ga] A. R. Gaio, *J*-holomorphic curves and moment maps, PhD thesis, University of Warwick, November 1999.

[GT] D. Gilbarg and N.S. Trudinger, *Elliptic Partial Differential Equations of the Second Order*, Springer 1983.

[Gr] M.Gromov, Pseudo holomorphic curves in symplectic manifolds, *Invent. Math.* **82** (1985), 307–347.

[K] F. Kirwan, *Cohomology of Quotients in Symplectic and Algebraic Geometry*, Princeton University Press, 1984.

[MS1] D. McDuff and D.A. Salamon, *J-holomorphic Curves and Quantum Cohomology*, University Lecture Series, **6**, American Mathematical Society, Providence, RI, 1994; second edition, 2001.

[MS2] D. McDuff and D.A. Salamon, *Introduction to Symplectic Topology*, Oxford University Press, 1995, 2nd edition 1998.

[M] I. Mundet, Yang-Mills-Higgs theory for symplectic fibrations, PhD thesis, Madrid, April 1999.

[RT] Y. Ruan and G. Tian, A mathematical theory of quantum cohomology, *J. Diff. Geom.* **42** (1995), 259–367.

[S] D.A. Salamon, Quantum products for mapping tori and the Atiyah-Floer conjecture, *Amer. Math. Soc. Transl.* **196** (1999), 199–235. Revised in December 2000, http://www.math.ethz.ch∼salamon

[U] K. Uhlenbeck, Connections with L^p-Bounds on Curvature, *Commun. Math. Physics.* **83** (1982), 31-42.

[W] K. Wehrheim, *Uhlenbeck Compactness*, to appear in Birkhäuser Verlag, Basel.